

Colorado

2006 Air Quality Data Report



Colorado Department
of Public Health
and Environment

Air Pollution Control Division

Cover photo

View of Mount Evans, from Highway 119 north of Black Hawk and Central City.

COLORADO AIR QUALITY DATA REPORT

2006



**Colorado Department
of Public Health
and Environment**

Air Pollution Control Division
APCD-TS-B1
4300 Cherry Creek Drive South
Denver, Colorado 80246-1530
(303) 692-3100

August 2007

This report is available electronically at <http://apcd.state.co.us/documents/techdocs.html>

Table of Contents

1.0 Purpose of the Annual Data Report	1
1.1 Symbols and Abbreviations	1
1.2 Description of Monitoring Areas in Colorado	1
1.2.1 Eastern Plains Counties.....	1
1.2.2 Northern Front Range Counties.....	2
1.2.3 Southern Front Range Counties.....	2
1.2.4 Mountain Counties.....	2
1.2.5 Western Counties.....	2
2.0 Criteria Pollutants	6
2.0.1 Exceedance Summary Table.....	6
2.1 Carbon monoxide - Sources.....	7
2.1.1 Carbon monoxide – Standards.....	7
2.1.2 Carbon monoxide – Health Effects.....	7
2.1.3 Carbon monoxide – Emissions	8
2.2 Ozone	8
2.2.1 Ozone – Standards	9
2.2.2 Ozone – Health Effects	9
2.2.3 Ozone – Sources	10
2.3 Sulfur dioxide.....	10
2.3.1 Sulfur dioxide – Standards.....	10
2.3.2 Sulfur dioxide – Health Effects.....	10
2.3.3 Sulfur dioxide – Sources.....	10
2.4 Nitrogen dioxide	11
2.4.1 Nitrogen dioxide – Standards.....	11
2.4.2 Nitrogen dioxide – Health Effects	11
2.4.3 Nitrogen dioxide – Sources.....	11
2.5 Particulate Matter – PM ₁₀	12
2.5.1 Particulate Matter – PM ₁₀ – Standards.....	12
2.5.2 Particulate Matter – PM ₁₀ – Health Effects	13
2.5.3 Particulate Matter – PM ₁₀ – Sources.....	13
2.6 Particulate Matter – PM _{2.5}	14
2.6.1 Particulate Matter – PM _{2.5} – Standards.....	14
2.6.2 Particulate Matter – PM _{2.5} – Health Effects.....	14
2.6.3 Particulate Matter – PM _{2.5} – Sources.....	15
2.7 Lead.....	15
2.7.1 Lead – Standards.....	15
2.7.2 Lead – Health Effects.....	16
2.7.3 Lead – Sources.....	16
3.0 Non-Criteria Pollutants	17
3.1 Visibility	17
3.1.1 Visibility – Standards.....	17
3.1.2 Visibility – Health Effects	17
3.1.3 Visibility – Sources.....	18
3.1.4 Visibility – Monitoring	18
3.1.5 Visibility – Denver Camera	18

3.2 Nitric Oxide	20
3.3 Total Suspended Particulates	20
3.4 Meteorology	20
3.5 Air Toxics	21
3.6 PM _{2.5} Chemical Speciation	21
4.0 Statewide Summaries For Criteria Pollutants	22
4.1 Carbon monoxide	22
4.2 Ozone	23
4.3 Sulfur Dioxide	24
4.4 Nitrogen Dioxide	24
4.5 Particulates – PM ₁₀	25
4.6 Particulates – PM _{2.5}	26
4.7 Lead	27
5.0 National Comparisons For Criteria Pollutants	28
5.1 Carbon monoxide	28
5.2 Ozone	28
5.3 Sulfur Dioxide	29
5.4 Nitrogen Dioxide	29
5.5 Particulates	30
5.6 Lead	31
6.0 Monitoring Results by Area in Colorado	32
6.1 Eastern Plains Counties	32
6.2 Northern Front Range Counties	35
6.3 Southern Front Range Counties	59
6.4 Mountain Counties	65
6.5 Western Counties	70
References	75

Table of Figures

Figure 1 - Monitoring Areas in Colorado	5
Figure 2 - Changes in National Carbon Monoxide Emissions from 1970 – 2005	8
Figure 3 - Changes in National Sulfur Dioxide Emissions from 1970 – 2005	11
Figure 4 - Changes in National Oxides of Nitrogen Emissions from 1970 – 2005	12
Figure 5 - Changes in National PM ₁₀ Emissions from 1970 – 2005	14
Figure 6 - Changes in National PM _{2.5} Emissions from 1990 -- 2005	15
Figure 7 - Changes in National Lead Emissions from 1970 – 2005	16
Figure 8 - Best and Worst Visibility Days for 2006	19
Figure 9 - Statewide Ambient Trends – Carbon Monoxide	22
Figure 10 - Statewide Ambient Trends – Ozone	24
Figure 11 - Statewide Ambient Trends – PM ₁₀	26
Figure 12 - Eastern Plains PM ₁₀ Particulate Graphs	33
Figure 13 - Eastern Plains PM _{2.5} Particulate Graph	33
Figure 14 - Eastern Plains Wind Rose Graph	34
Figure 15 - Northern Front Range PM ₁₀ Particulate Graphs	37 - 38
Figure 16 - Northern Front Range PM _{2.5} Particulate Graphs	39 - 40
Figure 17 - Northern Front Range Lead Graphs	41
Figure 18 - Northern Front Range Carbon Monoxide Graphs	43 - 45
Figure 19 - Northern Front Range Ozone Graphs	46 - 47
Figure 20 - Northern Front Range Nitrogen Dioxide Graphs	48
Figure 21 - Northern Front Range Sulfur Dioxide Graphs	48
Figure 22 - Denver Visibility Data (January 2006 to December 2006)	50
Figure 23 - Denver Visibility Comparison (1997 to 2006)	50
Figure 24 - Fort Collins Visibility Data (January 2006 to December 2006)	51
Figure 25 - Fort Collins Visibility Data (1997 to 2006)	52
Figure 26 - Northern Front Range Wind Roses	53 - 58
Figure 27 - Southern Front Range PM ₁₀ Particulate Graphs	60 - 61
Figure 28 - Southern Front Range PM _{2.5} Particulate Graphs	62
Figure 29 - Southern Front Range Lead Graph	63
Figure 30 - Southern Front Range Carbon Monoxide Graphs	63
Figure 31 - Southern Front Range Ozone Graph	64
Figure 32 - Southern Front Range Wind Rose	64
Figure 33 - Mountain Counties PM ₁₀ Particulate Graphs	66 - 67
Figure 34 - Mountain Counties PM _{2.5} Particulate Graphs	68
Figure 35 - Mountain Counties Lead Graphs	69
Figure 36 - Western Counties PM ₁₀ Particulate Graphs	71 - 72
Figure 37 - Western Counties PM _{2.5} Particulate Graph	73
Figure 38 - Western Counties Carbon Monoxide	74
Figure 39 - Western Counties Wind Roses	74

Table of Tables

Table 1 - Statewide Continuous Monitors In Operation For 2006.....	3
Table 2 - Statewide Particulate Monitors In Operation For 2006	4 - 5
Table 3 - National Ambient Air Quality Standards	6
Table 4 - 2005/2006 Exceedance Summaries	7
Table 5 - Historical Maximum 1-Hr and 8-Hr Carbon Monoxide Concentrations.....	23
Table 6 - Historical Maximum 1-Hour Ozone Concentrations.....	23
Table 7 - Historical Maximum Annual Average Sulfur Dioxide Concentrations.....	24
Table 8 - Historical Maximum Annual Average Nitrogen Dioxide Concentrations	25
Table 9 - Historical Maximum 24-Hour PM ₁₀ Concentrations.....	25
Table 10 - Historical Maximum Quarterly Lead Concentrations	27
Table 11 - 2006 National Ranking of Carbon Monoxide Monitors by 8-Hr Concentrations in ppm..	28
Table 12 - 2006 National Ranking of Ozone Monitors by 1-Hr Concentrations in ppm.....	28
Table 13 - 2006 National Ranking of Ozone Monitors by 8-Hr Concentrations in ppm.....	29
Table 14 - 2006 National Ranking of Sulfur Dioxide Monitors by 24-Hr Concentrations in ppm	29
Table 15 - 2006 National Ranking of Nitrogen Dioxide Monitors by 1-Hr Concentrations in ppm ...	30
Table 16 - 2006 National Ranking of PM ₁₀ Monitors by 24-Hr Maximum Concentrations in µg/m ³	30
Table 17 - 2006 National Ranking of PM _{2.5} Monitors by 24-Hr Maximum Concentrations in µg/m ³	30
Table 18 - 2006 National Ranking of Lead Monitors by 24-Hr Maximum Concentration in µg/m ³ ..	31
Table 19 - Eastern Plains Monitors In Operation For 2006.....	32
Table 20 - Eastern Plains Particulate Values For 2006.....	32
Table 21 - Northern Front Range Particulate Monitors In Operation For 2006	35
Table 22 - Northern Front Range Particulate Values For 2006	36
Table 23 - Northern Front Range TSP and Lead Values For 2006.....	40
Table 24 - Northern Front Range Continuous Monitors In Operation For 2006	42
Table 25 - Northern Front Range Carbon Monoxide Values for 2006	43
Table 26 - Northern Front Range Ozone Values For 2006.....	45
Table 27 - Northern Front Range Oxides of Nitrogen and Sulfur Dioxide Values For 2006	48
Table 28 - Denver Visibility Standard Exceedance Days.....	49
Table 29 - Fort Collins Visibility Standard Exceedance Days	51
Table 30 - Southern Front Range Monitors In Operation For 2006.....	59
Table 31 - Southern Front Range Maximum Particulate Values For 2006.....	60
Table 32 - Southern Front Range TSP and Lead Values For 2006.....	62
Table 33 - Southern Front Range Carbon Monoxide Values For 2006	63
Table 34 - Southern Front Range Ozone Values For 2006.....	64
Table 35 - Mountain Counties Monitors In Operation For 2006	65
Table 36 - Mountain Counties Particulate Values For 2006.....	66
Table 37 - Mountain Counties TSP and Lead Concentrations For 2006	69
Table 38 - Western Counties Monitors In Operation For 2006	70
Table 39 - Western Counties Particulate Values For 2006	71
Table 40 - Western Counties Carbon Monoxide Values For 2006.....	74

1.0 Purpose of the Annual Data Report

The Colorado Department of Public Health and Environment, Air Pollution Control Division (APCD) publishes the Colorado Air Quality Data Report as a companion document to the Colorado Air Quality Control Commission Report to the Public. The Air Quality Data Report addresses changes in ambient air quality measured by Division monitors. The Report to the Public discusses the policies and programs designed to improve and protect Colorado's air quality.

1.1 Symbols and Abbreviations

The following symbols and abbreviations are used through out this report:

- CO – Carbon monoxide
- SO₂ – Sulfur dioxide
- SO_x – Sulfur oxides
- NO_x – Nitrogen oxides
- NO – Nitric oxide
- NO₂ – Nitrogen dioxide
- O₃ – Ozone
- Met – meteorological measurements, wind speed, wind direction, temperature, relative humidity and standard deviation of horizontal wind direction.
- TSP – Total suspended particulates
- PM₁₀ – Particulate matter less than 10 microns in aerometric diameter.
- PM_{2.5} – Particulate matter less than 2.5 microns in aerometric diameter.
- Pb – Lead
- ppm – parts per million – this is used with gaseous pollutants.
- µg/m³ – micrograms per cubic meter – this is used with particulate pollutants.

1.2 Description of Monitoring Areas in Colorado

The state has been divided into five multi-county areas that are generally based on topography. The areas are: the Eastern Plains; the Northern Front Range; the Southern Front Range; the Mountain Counties and the Western Counties. These divisions are a somewhat arbitrary grouping of monitoring sites with similar characteristics.

The Eastern Plains Counties consist of those counties east of the I-25 corridor. These counties are generally rolling agricultural plains below 6000 feet.

The Front Range used in this definition is defined by the counties along or associated with the I-25 corridor not strictly by the Continental Divide. A division using the Continental Divide would place Leadville with the same counties as Colorado Springs and Denver. Leadville as the highest city in the U.S. has more in common with Breckenridge and Aspen than Denver or Colorado Springs.

The Mountain Counties are generally those on the west side of the Continental Divide and the Western Counties are the ones adjacent to the Utah border. Other divisions can and have been made, but these five divisions seemed appropriate for this report. Figure 1 shows the boundaries of these areas.

1.2.1 Eastern Plains Counties

The Air Pollution Control Division has only monitored for particulates and meteorology in the Eastern Plains Counties. The Eastern Plains Counties do not have the pollution sources that can generate health impacting concentrations of the criteria pollutants.

The Division has monitored for particulates in the communities along I-76, I-70 and along US Highway 50. The only monitors still in operation are in Lamar. The other monitors were discontinued after a review of the data showed that levels of particulates were well below the standard and were declining.

1.2.2 Northern Front Range Counties

The Northern Front Range Counties are those along the urbanized I-25 corridor from the Colorado/Wyoming border to just south of the city of Castle Rock. This area has the majority of the population in the state. It also has the majority of the monitors, with the Denver-metro area being the most heavily monitored. The remaining monitors are located in or near Fort Collins, Greeley, Longmont and Boulder.

1.2.3 Southern Front Range Counties

The Southern Front Range Counties are those along the urbanized I-25 corridor from south of the city of Castle Rock to the southern Colorado border. The cities with monitoring in the area include Colorado Springs, Pueblo, Cripple Creek, Cañon City and Alamosa. Alamosa is included because it shares more in common with the other cities in this group than it does with the mountain counties. Colorado Springs is the only city in the area that is monitored for carbon monoxide and ozone; the other cities are only monitored for particulates. In the past the APCD has conducted particulate monitoring in both Walsenburg and Trinidad. The monitoring in those cities was discontinued after a review of the data showed that levels of particulates were below the standard and were declining.

1.2.4 Mountain Counties

The Mountain Counties are those counties along the west side of the Continental Divide. The cities are usually located in tight mountain valleys where nighttime temperature inversions trap any pollution near the ground. Their primary monitoring concern is with particulate pollution from wood burning and road sanding. These communities range from Steamboat Springs in the north to Pagosa Springs in the south and include Vail, Silverthorne and Breckenridge in the I-70 corridor; Aspen, Leadville, Crested Butte, Mt. Crested Butte, and Gunnison in the central mountains.

1.2.5 Western Counties

The Western Counties generally contain smaller towns located in fairly broad river valleys. Grand Junction is the only large city in the area and the only location that monitors for carbon monoxide on the western slope. The other Western Slope monitors are located in the cities of Parachute, Delta, Durango and Telluride. These locations monitor only for particulates.

Table 1 - Statewide Continuous Monitors In Operation For 2006
X - Monitors continued in 2006 A – Monitors added in 2006
D – Monitors discontinued in 2006

County	Site Name	Location	CO	SO ₂	NO _x	O ₃	Met
Eastern Plains Counties							
Prowers	Lamar - POE	7100 Hwy 50					X
Northern Front Range Counties							
Adams	Commerce City	7101 Birch St.					X
	Welby	3174 E. 78 th Ave.	X	X	X	X	X
Arapahoe	Highland Res.	8100 S. University Blvd.				X	X
Boulder	Boulder	1405½ S. Foothills Hwy.				X	
	Longmont	440 Main St.	X				
Denver	Auraria Lot R	12 th St. & Auraria Parkway					X
	Denver CAMP	2105 Broadway	X	X	X	X	X
	Denver Carriage	2325 Irving St.	D			X	X
	Denver NJH	14 th Ave. & Albion St.	D				
	DESCI Building	1901 13 th Ave. (Visibility)					
	Firehouse #6	1300 Blake St.	X				
Douglas	Chatfield Res.	11500 N. Roxborough Pk. Rd.				X	X
Jefferson	Arvada	9101 W. 57 th Ave.	D			X	X
	NREL	2229 Old Quarry Rd.				X	
	Rocky Flats	16600 W. Hwy. 128				X	X
		9901 Indiana St.					X
	Welch	12400 W. Hwy. 285				X	X
Larimer	Fort Collins	708 S. Mason St.	X			X	X
		300 Remington St. (Visibility)					
		4407 S. College Ave.	X				
		3416 LaPorte Ave.				X	
Weld	Greeley	905 10 th Ave.	X				
		3101 35 th Ave.				X	
Southern Front Range Counties							
El Paso	Colorado Springs	1098 Glenn Ave.	D				
		USAFA Rd. 640				X	
		690 W. Hwy. 24	X				
	Manitou Springs	101 Banks Pl.				X	
Teller	Cripple Creek	2 nd St. & Warren Ave.					D
Western Counties							
Mesa	Grand Junction	645¼ Pitkin Ave.	X				X

Table 2 - Statewide Particulate Monitors In Operation For 2006

X - Monitors continued in 2006 A – Monitors added in 2006

D – Monitors discontinued in 2006 H – Hourly particulate monitor S – Chemical Speciation

County	Site Name	Location	TSP	Pb	PM ₁₀	PM _{2.5}
Eastern Plains Counties						
Elbert	Elbert	Wright-Ingraham Inst.				X
Prowers	Lamar	100 2 nd St.			X	
		104 Parmenter St.			X	
Northern Front Range Counties						
Adams	Brighton	22 S. 4 th Ave.			D	
	Commerce City	7101 Birch St.	D	D	X	X/H/S
	Globeville	5400 Washington St.	D	D		
	Welby	3174 E. 78 th Ave.			X/H	
Arapahoe	Arapahoe Comm. College	6190 S. Santa Fe Dr.				X
Boulder	Longmont	350 Kimbark St.			X	X/H
	Boulder	2440 Pearl St.			X	X
		2102 Athens St.				H
Denver	Denver CAMP	2105 Broadway	D	D	X/H	X/H
	Denver NJH	14 th Ave. & Albion St.				H
	Denver Visitor Center	225 W. Colfax Ave.			X	
	Lowry	8100 Lowry Blvd.			D	
	Denver Animal Shelter	678 S. Jason St.	A	A	A/H	A
	Swansea Elementary Sch.	4650 Columbine St.				X
Douglas	Chatfield Reservoir	11500 Roxborough Rd.				X/H
Larimer	Fort Collins	251 Edison St.			X	X
Weld	Greeley	1516 Hospital Rd.			X	X/H
	Platteville	1004 Main St.				X/S
Southern Front Range Counties						
Alamosa	Alamosa	208 Edgemont Blvd.			X	
		425 4 th St.			X	
El Paso	Colorado Springs	3730 Meadowlands			D	D
		101 W. Costilla St.	D	D	X	X/S
Fremont	Cañon City	128 Main St.			X	
Pueblo	Pueblo	211 E. D St.			X	X
Teller	Cripple Creek	209 Bennett Ave.			D	
Mountain Counties						
Archuleta	Pagosa Springs	309 Lewis St.			X	X
Gunnison	Crested Butte	603 6 th St.			X	
	Gunnison	221 N. Wisconsin St.			D	
	Mt. Crested Butte	19 Emmons Rd.			X	X
Lake	Leadville	510 Harrison St.	D	D		
Pitkin	Aspen	120 Mill St.			X/H	
Routt	Steamboat Springs	136 6 th St.			X	
Summit	Breckenridge	501 N. Park Ave.			X	

Table 2 - Statewide Particulate Monitors In Operation For 2006 (continued)


X - Monitors continued in 2006 A – Monitors added in 2006

D – Monitors discontinued in 2006 H – Hourly particulate monitor S – Chemical Speciation

County	Site Name	Location	TSP	Pb	PM ₁₀	PM _{2.5}
Western Counties						
Delta	Delta	560 Dodge St.			X	D
Garfield	Parachute	100 E. 2 nd St.			X	
	Rifle	144 E. 3 rd Ave.			X	
	New Castle	402 W. Main St.			X	
	Silt – Bell Ranch	512 Owens Dr.			X	
	Silt – Daley Ranch	884 County Rd. 327			X	
	Silt – Cox Ranch	5933 County Rd. 233			X	
	Glenwood Springs	109 8 th St.			X	
La Plata	Durango	1060 2 nd Ave.			D	
		56 Davidson Creek Rd.			D	
		1235 Camino del Rio			X	
		117 Cutler Dr.			D	
Mesa	Grand Junction	650 South Ave.			X	X/H/S
		645 ¼ Pitkin Ave.			H	
San Miguel	Telluride	333 W. Colorado Ave.			X	D

Figure 1

Monitoring Areas in Colorado



The pin symbols on the map show the approximate location of the monitors in Colorado.

2.0 Criteria Pollutants

The criteria pollutants are those for which the federal government has established ambient air quality standards in the Federal Clean Air Act and its amendments. There are six criteria pollutants. They are carbon monoxide, ozone, sulfur dioxide, nitrogen dioxide, lead and particulate matter. The standards for criteria pollutants are established to protect the most sensitive members of society. These are usually defined as those with respiratory problems, the very young and the infirm. The concentrations of each standard for the criteria pollutants are discussed in each section and a summary is presented in Table 3.

Table 3 - National Ambient Air Quality Standards¹

Pollutant	Averaging Time	Concentration
Carbon Monoxide (CO)		
Primary	1-hour*	35 ppm
Primary	8-hour*	9 ppm
Ozone (O₃)		
Primary	8-hour**	0.08 ppm
Secondary	Same as primary	
Nitrogen Dioxide (NO₂)		
Primary	Annual arithmetic mean	0.053 ppm
Secondary	Same as primary	
Sulfur Dioxide (SO₂)		
Primary	Annual arithmetic mean	0.03 ppm
Primary	24-hour*	0.14 ppm
Secondary	3-hour*	0.5 ppm
Particulate (PM₁₀)		
Primary	Annual arithmetic mean****	50 µg/m ³
Primary	24-hour***	150 µg/m ³
Particulate (PM_{2.5})		
Primary	Annual arithmetic mean****	15 µg/m ³
Primary	24-hour*****	65 µg/m ³
Lead (Pb)		
Primary	Calendar quarter	1.5 µg/m ³

* This concentration is not to be exceeded more than once per year.

** The 8-hour Ozone standard is set at 0.08 ppm as the 3-year average of the annual 4th maximum 8-hour average concentration.

*** The 24-hour standard is attained when the expected number of exceedances for each calendar year, averaged over three years, is less than or equal to one.

**** The annual arithmetic mean standard is a 3-year average.

***** The 24-hour PM_{2.5} standard is based on the three-year average of the 98th percentile.

2.0.1 Exceedance Summary Table

Table 4 is a summary of the number of exceedances of the ambient air quality standards for Colorado for 2005 and 2006. This table does not show sites in violation of the standard. That determination is based on a multi-year average for both PM₁₀ and for ozone. A discussion of the ozone standard is given in Section 2.2.1. The PM₁₀ standard is discussed in section 2.5.1.

In 2006, there were two monitors that recorded levels where the 4th maximum concentration was greater than the level of the 8-hour ozone standard. There were three monitors that recorded exceedances of the 24-hour PM₁₀ standard. None of the other criteria pollutant standards were exceeded.

Table 4 - 2005/2006 Exceedance Summaries

Location	2005	2006	
	PM10	Ozone*	PM10
Mt. Crested Butte	X		
Breckenridge	X		
Chatfield Reservoir		X	
Rocky Flats - N		X	
Alamosa ASC			X
Alamosa Municipal			X
Grand Junction			X

* - The ozone exceedances listed are those where the 4th maximum 8-Hr concentration for the year is greater than 0.085 ppm.

2.1 Carbon monoxide - Sources

Carbon monoxide is a colorless and odorless gas, formed when carbon in fuel is not burned completely. It is a component of motor vehicle exhaust, which contributes about 60 percent of all carbon monoxide emissions nationwide. Non-road vehicles account for the remaining carbon monoxide emissions from transportation sources. High concentrations of carbon monoxide generally occur in areas with heavy traffic congestion. In cities, as much as 85 percent of all carbon monoxide emissions may come from automobile exhaust. Other sources of carbon monoxide emissions include industrial processes, non-transportation fuel combustion, and natural sources such as wildfires. Peak carbon monoxide concentrations typically occur during the colder months of the year when carbon monoxide automotive emissions are greater and nighttime inversion conditions (where air pollutants are trapped near the ground beneath a layer of warm air) are more frequent.²

2.1.1 Carbon monoxide – Standards

The U.S. Environmental Protection Agency (EPA) has developed two national standards for carbon monoxide. They are 35 ppm averaged over a 1-hour period and 9 ppm averaged over an 8-hour period. These values are not to be exceeded more than once in a given year at any given location. A location will violate the standard with a second exceedance of either standard in a calendar year. The EPA directive requires that comparison with the carbon monoxide standards will be made in integers. Fractions of 0.5 or greater are rounded up, thus, actual concentrations of 9.5 ppm and 35.5 ppm or greater are necessary to exceed the 8-hour and 1-hour standards, respectively.³

2.1.2 Carbon monoxide – Health Effects

Carbon monoxide affects the central nervous system by depriving the body of oxygen. It enters the body through the lungs, where it combines with hemoglobin in the red blood cells. Normally, hemoglobin carries oxygen from the lungs to the cells. The oxygen attached to the hemoglobin is exchanged for the carbon dioxide generated by the cell's metabolism. The carbon dioxide is then carried back to the lungs where it is exhaled from the body. Hemoglobin binds approximately 240 times more readily with carbon monoxide than with oxygen. In the presence of carbon monoxide the distribution of oxygen is reduced throughout the body. Blood laden with carbon monoxide can weaken heart contractions with the result of lowering the volume of blood distributed to the body. It can significantly reduce a healthy person's ability to do manual tasks, such as working, jogging and walking. A life-threatening situation can exist for patients with heart disease when these people are unable to compensate for the oxygen loss by increasing the heart rate.²

The EPA has concluded that the following groups may be particularly sensitive to carbon monoxide exposures: angina patients, individuals with other types of cardiovascular disease, persons with chronic obstructive pulmonary disease, anemic individuals, fetuses and pregnant women. Concern also

exists for healthy children because of increased oxygen requirements that result from their higher metabolic rate.²


Carbon monoxide is exhausted from the body at varying rates, depending on physiological and external factors. The general guideline is that 20 to 40 percent is lost from the system after 2 to 3 hours following exposure.² The severity of health effects depends on both the concentration and the length of exposure because it takes time to remove it from the blood stream.

2.1.3 Carbon monoxide – Emissions

In Denver, the APCD estimates that 86 percent of the carbon monoxide emissions are from automotive sources. An estimated 3 percent of Denver's carbon monoxide emissions are from woodburning stoves and fireplaces. The remainder originates from aircraft, locomotives, construction equipment, power plants and space heating.⁴ These numbers are similar to the nationwide emissions.⁴

The percentage of carbon monoxide emissions contributed from various sources has not changed appreciably since 1970.⁴ What has changed is the amount of carbon monoxide emitted by these sources (Figure 2). In 1970, the total nationwide carbon monoxide emissions were approximately 197 million tons, in 2005, this had been reduced to 87 million tons.⁵

Figure 2 – Changes in National Carbon Monoxide Emissions from 1970 – 2005⁵



2.2 Ozone

“Ozone (O₃) is a gas composed of three oxygen atoms. It is not usually emitted directly into the air, but at ground-level is created by a chemical reaction between oxides of nitrogen (NO_x) and volatile organic compounds (VOC) in the presence of sunlight. Ozone has the same chemical structure whether it occurs miles above the earth or at ground-level and can be "good" or "bad," depending on its location in the atmosphere.

In the earth's lower atmosphere, ground-level ozone is considered "bad." Motor vehicle exhaust and industrial emissions, gasoline vapors, and chemical solvents as well as natural sources emit NO_x and VOC that help form ozone. Ground-level ozone is the primary constituent of smog. Sunlight and hot weather cause ground-level ozone to form in harmful concentrations in the air. As a result, it is known as a summertime air pollutant. Many urban areas tend to have high levels of "bad" ozone, but even rural areas are also subject to increased ozone levels because wind carries ozone and pollutants that form it hundreds of miles away from their original sources.

In the troposphere, the air closest to the Earth's surface, ground-level or "bad" ozone is a pollutant that is a significant health risk, especially for children with asthma. It also damages crops, trees and other vegetation. It is a main ingredient of urban smog.

The stratosphere, or "good" ozone layer extends upward from about 6 to 30 miles and protects life on Earth from the sun's harmful ultraviolet (UV) rays. This natural shield has been gradually depleted by man-made chemicals like chlorofluorocarbons (CFCs). A depleted ozone shield allows more UV from the sun to reach the ground, leading to more cases of skin cancer, cataracts, and other health problems.”⁶

2.2.1 Ozone – Standards

In July 1997, the U.S. Environmental Protection Agency established a new ozone standard. The reasons for these changes were:

“ . . . to provide protections for children and other at-risk populations against a wide range of ozone induced health effects, including decreased lung function (primarily in children active outdoors), increased respiratory symptoms (particularly in highly sensitive individuals), hospital admissions and emergency room visits for respiratory causes (among children and adults with pre-existing respiratory disease such as asthma), inflammation of the lung and possible long-term damage to the lungs.”⁷

“The 1-hour primary standard of 0.12 ppm was replaced by an 8-hour standard at a level of 0.08 ppm with a form based on the 3-year average of the annual 4th-highest daily maximum 8-hour average ozone concentration measured at each monitor within an area.”⁷

The 8-hour averaging time is more directly associated with health effects of concern at lower ozone concentrations than is the former 1-hour averaging time. Therefore, the 8-hour standard was felt to be more appropriate for a human health-based standard than the 1-hour standard.⁷ The EPA is currently in the process of re-examining the 8-hour standard. It may be lower in the future.

2.2.2 Ozone – Health Effects

Exposure to ozone has been linked to a number of health effects, including significant decreases in lung function, inflammation of the airways, and increased respiratory symptoms, such as cough and pain when taking a deep breath. Exposure can also aggravate lung diseases such as asthma, leading to increased medication use and increased hospital admissions and emergency room visits. Active children are the group at highest risk from ozone exposure because they often spend a large part of the summer playing outdoors. Children are also more likely to have asthma, which may be aggravated by ozone exposure. Other at-risk groups include adults who are active outdoors (e.g., some outdoor workers) and individuals with lung diseases such as asthma and chronic obstructive pulmonary disease. In addition, long-term exposure to moderate levels of ozone may cause permanent changes in lung structure, leading to premature aging of the lungs and worsening of chronic lung disease. Ozone also affects vegetation and ecosystems, leading to reductions in agricultural crop and commercial forest yields, reduced growth and survivability of tree seedlings, and increased plant susceptibility to disease, pests, and other environmental stresses (e.g., harsh weather). In long-lived species, these effects may become evident only after several years or even decades and may result in long-term effects on forest ecosystems. Ground level ozone injury to trees and plants can lead to a decrease in the natural beauty of our national parks and recreation areas.⁸

The on going review of the ozone standard (by the EPA and others) also highlighted concerns with ozone effects on vegetation for which the 1-hour ozone standard did not provide adequate protection. These effects can include reduction in agricultural and commercial forest yields, reduced growth and decreased survivability of tree seedlings, increased tree and plant susceptibility to disease, pests and other environmental stresses and potential long-term effects on forests and ecosystems.⁸

2.2.3 Ozone – Sources

Ozone is not emitted directly from a source, as are other pollutants, but forms as a secondary pollutant. Its precursors are certain reactive hydrocarbons and nitrogen oxides, which react chemically in sunlight to form ozone. The main sources for these reactive hydrocarbons are automobile exhaust, gasoline, oil storage and transfer facilities, industrial paint solvents, degreasing agents, cleaning fluids and ink solvents. High temperature combustion combines nitrogen and oxygen in the air to form oxides of nitrogen. Vegetation can also emit reactive hydrocarbons such as terpenes from pine trees, for example.⁹

Although some ozone is produced all year, the highest concentrations usually occur in the summer. The stagnant air and intense sunlight on hot, bright summer days provide the conditions for the precursor chemicals to react and form ozone. The ozone produced under these stagnant summer conditions remains as a coherent air mass and can be transported many miles from its point of origin.

2.3 Sulfur dioxide

Sulfur dioxide is a colorless gas with a pungent odor. It is detectable by smell at concentrations of about 0.5 to 0.8 ppm. It is highly soluble in water. In the atmosphere, sulfur oxides and nitric oxides are converted to “acid rain.”¹⁰

2.3.1 Sulfur dioxide – Standards

There are two primary standards for sulfur dioxide. The first is a long-term, one-year arithmetic average not to exceed 0.03 ppm. The second is a short-term, 24-hour average where concentrations are not to exceed 0.14 ppm more than once per year. The secondary standard is a 3-hour average not to exceed 0.5 ppm more than once per year.⁹

2.3.2 Sulfur dioxide – Health Effects


High concentrations of sulfur dioxide can result in temporary breathing impairment for asthmatic children and adults who are active outdoors. Short-term exposures of asthmatic individuals to elevated sulfur dioxide levels during moderate activity may result in breathing difficulties that can be accompanied by symptoms such as wheezing, chest tightness, or shortness of breath. Other effects that have been associated with longer-term exposures to high concentrations of sulfur dioxide, in conjunction with high levels of particulate matter, include aggravation of existing cardiovascular disease, respiratory illness, and alterations in the lungs’ defenses. The subgroups of the population that may be affected under these conditions include individuals with heart or lung disease, as well as the elderly and children.¹⁰

Together, sulfur dioxide and oxides of nitrogen are the major precursors to acidic deposition (acid rain), which is associated with the acidification of soils, lakes, and streams and accelerated corrosion of buildings and monuments. Sulfur dioxide also is a major precursor to PM_{2.5}, which is a significant health concern, and a main contributor to poor visibility.¹¹

2.3.3 Sulfur dioxide – Sources

Nationwide, over 65 percent of sulfur dioxide released to the air, or more than 13 million tons per year, comes from electric utilities, especially those that burn coal. Other sources of sulfur dioxide are industrial facilities that derive their products from raw materials like metallic ore, coal, and crude oil, or that burn coal or oil to produce process heat. Examples are petroleum refineries, cement manufacturing, and metal processing facilities. Also, locomotives, large ships, and some nonroad diesel equipment currently burn high sulfur fuel and release sulfur dioxide emissions to the air in large quantities.¹¹

Figure 3 - Changes in National Sulfur Dioxide Emissions from 1970 – 2005¹²



2.4 Nitrogen dioxide

In its pure state, nitrogen dioxide is a reddish-brown gas with a characteristic pungent odor. It is corrosive and a strong oxidizing agent. As a pollutant in ambient air, however, it is virtually colorless and odorless. Nitrogen dioxide can be an irritant to the eyes and throat. Oxides of nitrogen (nitric oxide and nitrogen dioxide) are formed when the nitrogen and oxygen in the air are combined in high temperature combustion.

2.4.1 Nitrogen dioxide – Standards

The annual standard for nitrogen dioxide is 0.053 ppm expressed as an annual arithmetic mean (average).¹³ “Pomona, CA, was the last U.S. city to record an exceedance of the nitrogen dioxide annual standard in 1989”¹⁴


2.4.2 Nitrogen dioxide – Health Effects

Elevated concentrations of nitrogen dioxide cause respiratory distress, degradation of vegetation, clothing and visibility, and increased acid deposition. Nitrate aerosols, which result from nitric oxide and nitrogen dioxide combining with water vapor in the air, have been consistently linked to Denver's visibility problems.

2.4.3 Nitrogen dioxide – Sources

About 44 percent of the emissions of nitrogen dioxide in the Denver area come from large combustion sources such as power plants. Almost 33 percent comes from motor vehicles, 15 percent from space heating, 3 percent from aircraft and 5 percent from miscellaneous off-road vehicles. Minor sources include fireplaces and woodstoves and high temperature combustion processes used in industrial work.¹⁵

Figure 4 - Changes in National Oxides of Nitrogen Emissions from 1970 – 2005¹⁶



2.5 Particulate Matter – PM₁₀

Particle pollution is a mixture of microscopic solids and liquid droplets suspended in air. This pollution, also known as particulate matter, is made up of a number of components, including acids (such as nitrates and sulfates), organic chemicals, metals, soil or dust particles, and allergens (such as fragments of pollen or mold spores).

The size of particles is directly linked to their potential for causing health problems. Small particles, less than 10 micrometers in diameter, pose the greatest problems. The smallest particles can get deep into your lungs, and some may even get into your bloodstream. Exposure to such particles can affect both your lungs and your heart. Larger particles are of less concern, although they can irritate your eyes, nose, and throat.

Small particles of concern include "fine particles" (such as those found in smoke and haze), which are 2.5 micrometers in diameter or less; and "coarse particles" (such as those found in wind-blown dust), which have diameters between 2.5 and 10 micrometers.¹⁷

2.5.1 Particulate Matter – PM₁₀ – Standards

In July 1987, EPA promulgated National Ambient Air Quality Standards for particulates with an aerodynamic diameter of 10 microns or less (PM₁₀). This is a size that can be inhaled into the bronchial and alveolar regions of the lungs. The standard has two forms, a 24-hour standard of 150 $\mu\text{g}/\text{m}^3$ and an annual arithmetic mean standard of 50 $\mu\text{g}/\text{m}^3$.¹⁸

1. The 24-hour standard is attained when the expected number of exceedances for each calendar year, averaged over three years, is less than or equal to one. The estimated number of exceedances is computed quarterly using available data and adjusting for missing sample days.
2. The annual arithmetic mean standard is attained when the annual mean, averaged over three years is less than or equal to the level of the standard. Each annual mean is computed from the average of each quarter in the year, with adjustments made for missing sample days.
3. In both cases, a data recovery of 75 percent is needed for each calendar quarter to be considered a valid quarter of data.

The 24-hour standard was modified in by EPA in July 1997, but was subsequently nullified back to this form in May 1999 due to a challenge in the courts.

2.5.2 Particulate Matter – PM₁₀ – Health Effects

According to American Lung Association's paper The Perils of Particulates:

“The health risk from an inhaled dose of particulate matter depends on the size and concentration of the particulate. Size determines how deeply the inhaled particulate will penetrate into the respiratory tract where they can persist and cause respiratory damage. Particles less than 10 microns in diameter are easily inhaled deep into the lungs. In this range, larger particles tend to deposit in the tracheobronchial region and smaller ones in the alveolar region. Particulates deposited in the alveolar region can remain in the lungs for long periods because the alveoli have a slow mucociliary clearance system.”¹⁹


“Fine particulate pollution does not affect the health of exposed persons with equal severity. Certain subgroups of people potentially exposed to air pollutants can be identified as potentially ‘at risk’ from adverse health effects of air borne pollutants. There is very strong evidence that asthmatics are much more sensitive (i.e., respond with symptoms at relatively low concentrations) to the effects of particulates than the general healthy population.”²⁰

The welfare effects of particulate exposure may be the most widespread of all the pollutants. Because of the potential for extremely long-range transport of fine particles and chemical reactions that occur, no place on earth has been spared from the particulate pollution generated by urban and rural sources. The effects of particulates range from visibility degradation to climate changes and vegetation damage. General soiling, commonly thought to be just a nuisance, can have long-term adverse effects on building paints and other materials. Acid deposition as particulates can be detected in the most remote areas of the world.

2.5.3 Particulate Matter – PM₁₀ – Sources

Most anthropogenic (manmade) particulates are in the 0.1 to 10 micron diameter range. Particles larger than 10 microns are usually due to “fugitive dust”. Fugitive dust is wind-blown sand and dirt from roadways, fields and construction sites that contain large amounts of silica (sand-like) materials. Anthropogenic particulates are created during the burning of fuels associated with industrial processes or heating. These particulates include fly ash (from power plants), carbon black (from automobiles and diesel engines) and soot (from fireplaces and woodstoves). The PM₁₀ particulates from these sources contain a large percentage of elemental and organic carbon. These types of particles play a role in both visual haze and health issues.²⁰ Figure 5 shows the changes in national PM₁₀ emissions from 1970 through 2005.

Figure 5 - Changes in National PM₁₀ Emissions from 1970 – 2005²¹



2.6 Particulate Matter – PM_{2.5}

According to the Environmental Protection Agency’s Latest Findings on National Air Quality: 2000 Status and Trends, Particulate Matter, “PM_{2.5} is composed of a mixture of particles directly emitted into the air and particles formed in the air by the chemical transformation of gaseous pollutants. The principle types of secondary pollutants are ammonium sulfate and ammonium nitrate formed in the air from gaseous emissions of sulfur dioxide and oxides of nitrogen, reacting with ammonia. The main source of sulfur dioxide is combustion of fossil fuels in boilers and the main source of oxides of nitrogen are the combustion of fossil fuels in boilers and mobile sources. Some secondary particles are also formed from semi-volatile organic compounds which are emitted from a wide range of combustion sources.”

2.6.1 Particulate Matter – PM_{2.5} – Standards

In 1997, the EPA added new fine particle standards, PM_{2.5}, to the existing PM₁₀ standards. EPA added an annual PM_{2.5} standard set at a concentration of 15 micrograms per cubic meter (µg/m³) and a 24-hour PM_{2.5} standard set at 65 µg/m³. The annual component of the standard was set to provide protection against typical day-to-day exposures as well as longer-term exposures, while the daily component protects against more extreme short-term events.

Areas will be considered in compliance with the annual PM_{2.5} standard when the 3-year average of the annual arithmetic mean PM_{2.5} concentrations, from single or multiple community-oriented monitors, is less than or equal to 15 µg/m³. The 24-hour PM_{2.5} standard is based on the 98th percentile of 24-hour PM_{2.5} concentrations in a year (averaged over 3 years). The change to a percentile based standard from a second maximum based standard was designed to eliminate the effect of anomalously high concentrations. In addition, this change is an attempt to focus more on the true health effects of the pollutant. The 24-hour standard was lowered by EPA on September 20, 2006 to 35 µg/m³. This new standard did not become effective until December 17, 2006.

2.6.2 Particulate Matter – PM_{2.5} – Health Effects

The health effects of PM_{2.5} are not just a function of their size, 1/20th the size of a human hair, which allows them to be breathed deeply into the alveoli the lungs, but of their composition. These particles can remain in the lungs for a long time and cause a great deal of damage to the lung tissue. They can reduce lung function as well as cause or aggravate respiratory problems. They can increase the long-term risk of lung cancer or lung diseases such as emphysema or pulmonary fibrosis.²²


2.6.3 Particulate Matter – PM_{2.5} – Sources

Figure 6 shows the nationwide changes in emissions of PM_{2.5} particulates from 1990 through 2006.

The primary source of fine particles emitted directly into the air is carbonaceous material from combustion. Secondary particles, are another large source of “fine” particulates. Secondary particles are those that are created in the atmosphere by chemical reactions of gaseous pollutants and water vapor to form a semi-solid particle.²¹

Particles less than 2.5 microns in diameter, or PM_{2.5}, are the major contributors to visibility problems because of their ability to scatter or absorb light. In Denver, the effects of this particulate pollution can be seen as the “Brown Cloud” or more appropriately, the “Denver Haze” because it is frequently neither brown nor an actual cloud.

Figure 6 - Changes in National PM_{2.5} Emissions from 1990 -- 2005²³



2.7 Lead

Lead is a metal found naturally in the environment as well as in manufactured products. The major sources of lead emissions have historically been motor vehicles (such as cars and trucks) and industrial sources. Due to the phase out of leaded gasoline, metals processing is the major source of lead emissions to the air today. The highest levels of lead in air are generally found near lead smelters. Other stationary sources are waste incinerators, utilities, and lead-acid battery manufacturers.²⁴

2.7.1 Lead – Standards

The current federal standard for lead is a calendar quarter (3-month) average concentration not to exceed 1.5 micrograms of lead per cubic meter of air ($\mu\text{g}/\text{m}^3$). This standard was established to maintain blood lead concentrations below 30 micrograms per deciliter ($\mu\text{g}/\text{dL}$) due to exposure to atmospheric lead concentrations.²⁵ In the future, the focus on lead monitoring will shift to ensure that stationary sources do not create violations of the standard in localized areas. Colorado had at least one such source in the Denver area that was the subject of monitoring. This source ceased operation in August of 2006. The Historical Lead Comparison graphs show data back to 1990. The concentrations recorded at most of the monitoring sites are approaching the limits of detection for ambient lead. The last violation of the lead standard in Colorado was the first quarter of 1980.


2.7.2 Lead – Health Effects

Exposure to lead occurs mainly through inhalation of air and ingestion of lead in food, water, soil, or dust. It accumulates in the blood, bones, and soft tissues and can adversely affect the kidneys, liver, nervous system, and other organs. Excessive exposure to lead may cause neurological impairments such as seizures, mental retardation, and behavioral disorders. Even at low doses, lead exposure is associated with damage to the nervous systems of fetuses and young children, resulting in learning deficits and lowered IQ. Recent studies also show that lead may be a factor in high blood pressure and subsequent heart disease. Lead can also be deposited on the leaves of plants, presenting a hazard to grazing animals and humans through ingestion.²⁶

2.7.3 Lead – Sources

“Because of the phase-out of leaded gasoline, lead emissions and concentrations decreased sharply during the 1980s and early 1990s. Emissions of lead decreased 96 percent over the 24-year period 1980–2004. These large reductions in long-term lead emissions from transportation sources have changed the nature of the ambient lead problem in the United States. Because industrial processes are now responsible for all violations of the lead NAAQS, the lead monitoring strategy currently focuses on emissions from these point sources.”²⁷ Figure 7 shows the decline in lead emissions in the past 36 years.

Figure 7 - Changes in National Lead Emissions from 1970 – 2005²⁷



3.0 Non-Criteria Pollutants

Non-criteria pollutants are those pollutants for which there are no current national ambient air quality standards. These include but are not limited to pollutants that impair visibility, total suspended particulates, nitric oxide and air toxics. Meteorological measurements of wind speed, wind direction, temperature and humidity are also included in this group.

3.1 Visibility

Visibility is unique among air pollution effects in that it involves human perception and judgment. It has been described as the maximum distance that an object can be perceived against the background sky. Visibility also refers to the clarity with which the form and texture of distant, middle and near details can be seen as well as the sense of the trueness of their apparent coloration. As a result, measures of visibility serve as surrogates of human perception. There are several ways to measure visibility but none of them tell the whole story or completely measure visibility as human beings experience it.

3.1.1 Visibility – Standards

The Colorado Air Quality Control Commission established a visibility standard in 1990 for the Front Range cities from Fort Collins to Colorado Springs. The standard, an atmospheric extinction of 0.076 per kilometer, was based on the public's definition of unacceptable amounts of haze as judged from slides of different haze levels taken in the Denver area. At the standard, 7.6 percent of the light in a kilometer of air is blocked. The standard applies from 8 A.M. to 4 P.M. each day, during those hours when the relative humidity is less than 70 percent. Visibility, along with meteorology and concentrations of other pollutants for which National Ambient Air Quality Standards exist, is used to determine the need for mandatory woodburning and voluntary driving restrictions.

There is no quantitative visibility standard for Colorado's pristine and scenic rural areas. However, in the 1977 amendments to the Federal Clean Air Act, Congress added Section 169a²⁸ and established a national visibility goal that created a qualitative standard of “the prevention of any future and the remedying of any existing, impairment of visibility in mandatory Class I federal areas which impairment results from manmade air pollution”. The implementation of Section 169a has led to federal requirements to protect visual air quality in large national parks and wilderness areas.²⁹ Colorado has 12 of these Class I areas. Federal and state law prohibits visibility impairment in national parks and wildernesses due to large stationary sources of air pollution.

3.1.2 Visibility – Health Effects

Visual air quality is an element of public welfare. Specifically, it is an important aesthetic, natural and economic resource of the state of Colorado. The worth of visibility is difficult to measure; yet good visibility is something that people undeniably value. Impaired visibility can affect the enjoyment of a recreational visit to a scenic mountain area. Similarly, people prefer to have clear views from their homes and offices. These concerns are often reflected in residential property values and office rents. Any loss in visual air quality may contribute to corresponding losses in tourism and usually make an area less attractive to residents, potential newcomers and industry.

There is increasing information that shows a correlation between ambient concentrations of particulate matter and respiratory illnesses. Some researchers believe this link may be strongest with concentrations of fine particles, which also contribute to visibility impairment. In July 1997, the EPA developed a National Ambient Air Quality Standard for particulate matter less than 2.5 microns in diameter (PM_{2.5}). See the section 2.6 for more information on PM_{2.5}. Any control strategies to lower ambient concentrations of fine particulate matter for health reasons will also improve visibility.

3.1.3 Visibility – Sources

The cause of visibility impairment in Colorado is most often fine particles in the 0.1 to 2.5 micrometer size range (one micrometer is a millionth of a meter). Light passing from a vista to an observer is either scattered away from the sight path or absorbed by the atmospheric fine particulate. Sunlight entering the pollution cloud may be scattered into the sight path adding brightness to the view and making it difficult to see elements of the vista. Sulfate, nitrate, elemental carbon and organic carbon are the types of particulate matter most effective at scattering and/or absorbing light. The man-made sources of these particulates include woodburning, electric power generation, industrial combustion of coal or oil, and emissions from cars, trucks and buses.

Visibility conditions vary considerably across the state. Usually, visibility in Colorado is among the best in the country. Our prized western vistas exist due to unique combinations of topography and scenic features. Air in much of the West contains low humidity and minimal levels of visibility-degrading pollution. Nevertheless, visibility problems occur periodically throughout the state. Woodburning haze is a concern in several mountain communities each winter. Denver has its “Brown Cloud.” Even the national parks, monuments, and wilderness areas shows pollution-related visibility impairment on occasion due to regional haze, the interstate or even regional-scale transport of visibility-degrading pollution.

The visibility problems across the state have raised public concern and spurred research. The goal of Colorado's visibility program is to protect visual air quality where it is presently good and improve visibility where it is degraded.

3.1.4 Visibility – Monitoring

There are several ways to measure visibility. Currently, the Division uses camera systems to provide qualitative visual documentation of a view. Transmissometers and nephelometers are used to measure the atmosphere's ability to attenuate light quantitatively.

A visibility site was installed in Denver in late 1990 using a long-path transmissometer. Visibility in the downtown area is monitored using a receiver located near Cheesman Park at 1901 E. 13th Avenue and a transmitter located on the roof of the Federal Building at 1929 Stout Street. This instrument directly measures light extinction, which is proportional to the ability of atmospheric particles and gases to attenuate image-forming light as it travels from an object to an observer. The visibility standard is stated in units of atmospheric extinction. Days when the visibility is affected by rain, snow or high relative humidity are termed “excluded” (as shown in Figures 21 and 23) and are not counted as violations of the visibility standard. In September 1993, a transmissometer and nephelometer were purchased by the city of Fort Collins to monitor visibility.

In Colorado, several agencies of the federal government, in cooperation with regional and nationwide state air pollution organizations, also monitor visibility in a number of national parks and wilderness areas “Class I” areas, either individually or jointly through the Inter-agency Monitoring of PROtected VIsual Environments (IMPROVE) monitoring program. The goals of the monitoring programs are to establish background visibility levels, identify trends of deterioration or improvement, to identify suspected sources of visibility impairment and to track regional haze. Visibility and the atmospheric constituents that cause visibility degradation are characterized with camera systems, transmissometers and extensive fine-particle chemical composition measurements by the monitoring network. There are currently monitoring sites in Rocky Mountain National Park, Mesa Verde National Park, Weminuche Wilderness, Mount Zirkel Wilderness, Great Sand Dunes National Monument and Maroon Bells/Snowmass Wilderness. These data are not contained in this report, but are available at this web site address: <http://vista.cira.colostate.edu/improve/>

3.1.5 Visibility – Denver Camera

The Division operates a web-based camera that can be viewed by clicking on the “Live Image” tab on the left side of the screen at the Air Pollution Control Division's web site <http://apcd.state.co.us/psi/main.html>. There is a great deal of other information available from this site in

addition to the image from the visibility camera. The Front Range Air Quality Forecast, Air Quality Advisory, Monitoring Reports and Open Burning Forecast are also available.

The images in Figure 8 show the visibility on one of the “Best” and “Worst” days in 2006. The “Best “ visibility day was December 10, 2006. The “Worst” visibility day was December 19, 2006.

Figure 8 - Best and Worst Visibility Days for 2006



These two pictures are images made by the web camera at the visibility monitor located at 1901 E. 13th Avenue in Denver. These images are centered on the Federal Building at 1929 Stout Street. The difference in these two pictures is the brightness and detail that can be seen in the image on the top as compared to the image on the bottom. Look specifically at the edges of the downtown buildings and the area on the horizon at the right edge of the picture.

3.2 Nitric Oxide

Nitric oxide is the most abundant of the oxides of nitrogen emitted from combustion sources. There are no known adverse health effects at normal ambient concentrations. However, nitric oxide is the precursor, is involved in the reaction of nitrogen dioxide, nitric acid, nitrates and ozone, all of which have demonstrated adverse health effects.³⁰ There are no federal or state standards for nitric oxide.

3.3 Total Suspended Particulates

Total suspended particulates (TSP) were first monitored in Colorado in 1960 at 414 14th St. in Denver. This location monitored particulates until 1988. The Adams City and Gates total suspended particulate monitors began operation in 1964 and the Denver CAMP monitor at 2105 Broadway began operating in 1965. Either the Federal EPA or the City of Denver operated these monitors until the mid-1970s when daily operation was taken over by the Colorado Department of Public Health and Environment.

Particulate monitoring expanded to more than 70 locations around the state by the early 1980s. The primary standards for total suspended particulates were 260 $\mu\text{g}/\text{m}^3$ as a 24-hour sample and 75 $\mu\text{g}/\text{m}^3$ as an annual geometric mean. On July 1, 1987, with the promulgation of the PM_{10} standards, the old particulate standards were eliminated. Until December 2006 the Division operated six TSP samplers to measure lead. On January 1, 2007 the number of lead monitoring sites were reduced to one, at the Denver Municipal Animal Shelter located at 678 S. Jason Street. The reason for the change in the number of lead monitors is that the ambient concentrations of lead have been reduced dramatically and federal monitoring requirements have been changed.

3.4 Meteorology

The Air Pollution Control Division takes a limited set of meteorological measurements at 18 locations around the state. These measurements include wind speed, wind direction, temperature, standard deviation of horizontal wind direction and some monitoring of relative humidity. Relative humidity measurements are also taken in conjunction with the two visibility monitors. The humidity data are not summarized in this report since they are used primarily to validate the visibility measurements taken at the specific locations. The Division does not collect precipitation measurements. The wind speed, wind direction and temperature measurements are collected primarily for air quality forecasting and air quality modeling. The instruments are on ten-meter towers and the data are stored as hourly averages.

The wind roses displayed in this report are placed on a background map that shows the approximate location of the meteorological site. The wind roses are based on the direction that the wind is blowing from. Another way of visualizing a wind rose is to picture yourself standing in the center of the plot and facing into the wind. The wind direction is broken down in the 16 cardinal directions (i.e. N, NNE, NE, ENE, E, ESE, SE, SSE, S, etc). The wind speed is broken down in six categories. The graphs in this report use 1-3 mph, 4-5 mph, 7-11 mph, 12-14 mph, 15-38 mph and greater than 38 mph. The length of each arm of the wind rose represents the percentage of time the wind was blowing from that direction at that speed. The longer the arm the greater percentage of time the wind is blowing from that direction. A review of the wind rose in Figure 26, for example, shows that in Arvada the majority of the winds come from the west and west-northwest and that these winds are generally in the 1-3 mph and 4-6 mph ranges.

3.5 Air Toxics

Toxic air pollutants, or air toxics, are those pollutants that cause or may cause cancer or other serious health effects, such as reproductive effects or birth defects. Air toxics may also cause adverse environmental and ecological effects. EPA is required to reduce air emissions of 188 air toxics listed in the Clean Air Act. Examples of toxic air pollutants include benzene, found in gasoline; perchloroethylene, emitted from some dry cleaning facilities; and methylene chloride, used as a solvent by a number of industries. Most air toxics originate from man-made sources, including mobile sources (e.g., cars, trucks, construction equipment) and stationary sources (e.g., factories, refineries, power plants), as well as indoor sources (e.g., some building materials and cleaning solvents). Some air toxics are also released from natural sources such as volcanic eruptions and forest fires.³¹

People exposed to toxic air pollutants at sufficient concentrations may experience various health effects including cancer and damage to the immune system, as well as neurological, reproductive (e.g., reduced fertility), developmental, respiratory and other health problems. In addition to exposure from breathing air toxics, risks also are associated with the deposition of toxic pollutants onto soils or surface waters, where they are taken up by plants and ingested by animals and eventually magnified up through the food chain. Like humans, animals may experience health problems due to air toxics exposure.

The APCD currently monitors for air toxics in Grand Junction as part of EPA's National Air Toxics Trend Stations. The data from this study will be presented in a separate report.

3.6 PM_{2.5} Chemical Speciation

Chemical speciation analysis is conducted on some PM_{2.5} filters. These analyses are conducted for several elements and chemical compounds, which can cause serious health effects, premature deaths, visibility degradation and regional haze. There are two broad categories of PM_{2.5}: primary and secondary particles. Primary PM_{2.5} particles are those emitted directly to the air from crushed geologic materials to carbonaceous particles from incomplete combustion (see section 2.6.3 for more information on PM_{2.5} sources). Secondary PM_{2.5} is formed from gases that combine in the atmosphere through chemical processes and form liquid aerosol droplets. If the PM_{2.5} pollution needs to be controlled it is important to know the composition of PM_{2.5} particles so that the appropriate sources can be targeted for control.

Numerous health effects studies have correlated negative health effects to the total mass concentration of PM_{2.5} in ambient air.¹⁸ However, it has not yet been determined if the health correlation is to total mass concentration or to concentrations of specific chemical species in the PM_{2.5} mix. When the EPA promulgated the NAAQS for PM_{2.5} in 1997 the compliance (mass) monitoring part of the network was established first. Mass concentrations from the compliance network are used to determine attainment of the NAAQS. EPA soon supplemented the PM_{2.5} network with chemical speciation monitoring to provide information on the chemical composition of PM_{2.5}. The main purposes are to identify sources, develop implementation plans to reduce PM_{2.5} pollution and support health effects research.

Colorado began chemical speciation monitoring at the Commerce City site in February 2001 at the state's only speciation trend network site. Four other chemical speciation sites were established in 2001 in the following areas: Colorado Springs, Durango, Grand Junction and Platteville. The Durango site was closed in September 2003. The Colorado Springs site was removed on December 31, 2006. It will moved to the Denver Municipal Animal Shelter location in 2007. Each air filter is analyzed for gravimetric mass, 48 elemental concentrations (sodium through lead), organic (four types) and elemental carbon and five ions (ammonium, sodium, potassium, sulfate and nitrate.) Selected filters can also be analyzed for semi volatile organics and microscopic analyses. The results of these samples can be obtained from the Air Pollution Control Division upon request.

4.0 Statewide Summaries For Criteria Pollutants

4.1 Carbon monoxide


Carbon monoxide concentrations have dropped dramatically from the early 1970s. This change can be seen in both the concentrations measured and the number of monitors that exceeded the level of the 8-hour standard. In 1975, 9 of the 11 state-operated monitors exceeded the 8-hour standard. In 1980, 13 of the 17 state-operated monitors exceeded the 8-hour standard. Since 1996 none of the state-operated monitors have recorded a violation of the 8-hour standard. In 2006 the highest statewide 2nd maximum 8-hour concentration was a 3.3 ppm recorded at the Greeley, 905 10th Avenue. monitor.

Figure 9, shows the trend of the statewide average for the second maximum 1-hour and 8-hour concentrations for carbon monoxide for the periods from 1970 to 2006.

Two important points to note are:

1. Before 1989 the average 2nd maximum 8-hour concentration for all state-operated carbon monoxide monitors was greater than the 8-hour standard of 9.5 ppm.
2. In the last 5 years the downward trend in concentrations has continued, but at a slower rate. The statewide average 8-hour concentration is now about one quarter of the standard or 2.4 ppm.

Figure 9 - Statewide Ambient Trends – Carbon Monoxide²⁸



The trend in the 1-hour average carbon monoxide concentrations statewide has fallen more dramatically than the 8-hour concentrations. The maximum 1-hour concentration ever recorded at any of the state-operated monitors was a 79.0 ppm recorded at the Denver CAMP monitor in 1968. In 2006, the maximum 1-hour concentration recorded was 9.3 ppm recorded at the Denver Firehouse #6 monitor. The 1-hour annual maximum concentrations have declined from more than twice the standard in the late 1960s to about one quarter of the standard in 2006. Table 5 presents the historical maximum values.

Table 5 - Historical Maximum 1-Hr and 8-Hr Carbon Monoxide Concentrations³²

1-Hour ppm	Location	Date	Number of Annual Exceedances	8-Hour ppm	Location	Date	Number of Annual Exceedances
79.0	CAMP	11-20-68	13	48.1	CAMP	12-21-73	133
70.0	CAMP	11-21-74	15	33.9	CAMP	12-28-65	197
67.0	CAMP	12-21-73	21	33.4	CAMP	12-04-81	42
65.0	CAMP	12-21-73	21	33.2	CAMP	12-23-71	188
64.9	NJH-W	11-16-79	15	33.1	CAMP	11-20-68	98
2006 Maximum Carbon Monoxide Concentration							
9.3	Firehouse #6	02-23-06	0	3.4	Greeley	12-08-06	0

4.2 Ozone

A complete analysis of the trend in ozone values over time is more complex than the simple linear regression used for this report since it must deal with variations in meteorological conditions from year to year. However, Figure 10 Statewide Ambient Trends, shows that the second maximum 1-hour ozone concentrations have declined since 1985. The linear regression trend is not as clear for the 8-hour average ozone concentrations, but over the past 20 years it is essentially flat. According to the Denver Early Action Ozone Compact, February 2004 the high values seen in 2003 were the result of “Anomalously high temperatures and anomalously low mixing heights. . .”.


The Division conducted a detailed analysis of the ozone trends as a part of the Denver Early Action Ozone Compact, February 2004. That report concluded that there had been a decline in the daily 8-hour concentrations of 1.2 percent per year for the period from 1993 through 2003. The full report is available on the web at http://apcd.state.co.us/documents/eac/Denver_EAC-WOEv4.pdf.

Table 6 lists the five highest 1-hour ozone concentrations recorded in Colorado. Ozone monitoring began in 1972 at the Denver CAMP station and eight exceedances of the 1-hour standard were recorded that year. However, data before 1975 are not included because quality assurance and maintenance records are no longer available. In addition, a review of the ozone data before 1975 shows several values that are questionable because of time of day, time of year and inconsistencies with other monitors in the area.

Table 6 - Historical Maximum 1-Hour Ozone Concentrations³³

1-Hour ppm	Monitor	Date
0.223	Welby	03-03-78
0.197	Arvada	07-28-75
0.186	Children’s Asthmatic Research Institute and Hospital, 21 st Ave. & Julian St.	09-17-76
0.184	Arvada	06-30-76
0.182	Welby	08-05-75
2006 Maximum Ozone Concentration		
0.112	Welch	07-29-06

Figure 10 - Statewide Ambient Trends – Ozone



4.3 Sulfur Dioxide

The concentrations of sulfur dioxide in Colorado have never been a major health concern since we have few industries that burn large amounts of coal. The concern in Colorado with sulfur dioxide has been associated with acid deposition and its effects on the mountain lakes and streams. Historically the maximum annual concentration recorded by APCD monitors was 0.018 ppm in 1979 at the Denver CAMP monitor. The annual standard is 0.030 ppm. Since 1990, the annual average at the Denver CAMP monitor has declined from a high in 1992 of 0.010 ppm to 0.003 ppm in 2004.

Figure 21 shows both the declining trend in sulfur dioxide readings as well as the generally low concentrations of sulfur dioxide recorded at the APCD’s monitors. This same trend is evident, although not as pronounced, in the 3-hour and 24-hour averages as well.

Table 7 - Historical Maximum Annual Average Sulfur Dioxide Concentrations³⁴

Annual Average ppm	Monitor	Date
0.018	CAMP	1979
0.013	CAMP	1980
0.013	CAMP	1981
0.013	CAMP	1983
0.012	CAMP	1978
2006 Maximum Sulfur Dioxide Concentration		
0.0029	CAMP	2006

4.4 Nitrogen Dioxide

Colorado exceeded the nitrogen dioxide standard in 1977 at the Denver CAMP monitor. Concentrations have shown a gradual decline for the past 20 years. However, for the trend for the past ten years in the annual average has been nearly flat.

Figure 20 shows that levels have declined at the Welby monitor over the past ten years the annual average at the Denver CAMP monitor has shown little to no change at all. The cause of this is most likely

due to an increase in the number of vehicles and increased power generation associated with the increases in population in the Denver-metro area.

Table 8 - Historical Maximum Annual Average Nitrogen Dioxide Concentrations³⁵

Annual Average ppm	Monitor	Date
0.052	CAMP	1975
0.052	CAMP	1976
0.052	CAMP	1979
0.052	CAMP	1973
0.051	CAMP	1977
2006 Maximum Nitrogen Dioxide Concentration		
0.029	CAMP	2006

4.5 Particulates – PM₁₀

Particulate matter 10 microns and smaller (PM₁₀) data have been collected in Colorado since 1985. The samplers were modified in 1987 to conform to the requirements of the new standard when it was established in July of 1987. Therefore, annual trends are only valid back to July 1987.

Since 1988, the state has had at least one monitor exceed the level of the 24-hour PM₁₀ standard (150 µg/m³) every year except 2004. By contrast, no monitor with at least 75 percent data recovery has exceeded the level of the annual standard (50 µg/m³). As seen in the graph on page 26 there is a great deal more variation in the 24-hour maximum values than in the annual averages.

The data contained in Table 9-the Historical Maximum values table, include those concentrations that are the result of exceptional events. There have been several of these events documented in Colorado since PM₁₀ monitoring began in 1988. In general, in order to qualify for exclusion, a value (or values) has to be associated with a regional natural phenomenon. One such event was the large wind and dust storm that occurred on March 31, 1999 when monitors from Steamboat Springs to Telluride reported high PM₁₀ concentrations. Similar exceptional events have been documented in Lamar and Alamosa. These events are not included in NAAQS determinations, not because they are without any health risk but because they are natural and are not controllable or predictable.


Table 9 - Historical Maximum 24-Hour PM₁₀ Concentrations³⁶

24-Hour Maximum µg/m ³	Monitor	Date
412	Alamosa	04-10-91
306	Cripple Creek	12-27-95
262	Pagosa Springs	12-29-94
236	Aspen	02-22-91
235	Cripple Creek	02-11-97
2006 Maximum PM₁₀ Concentration		
424	Alamosa ASC	02-10-06

Most of these high values are natural events, high winds and dry conditions, that result in blowing dust. This includes the high value from Alamosa – ASC in 2006.

Figure 11-Statewide Ambient Trends shows a decline in both the 24-hour and the Annual average concentrations since 1987. This graph has been modified from previous years in that the exceptional events have been excluded from the trend data. The 412 $\mu\text{g}/\text{m}^3$ in 1991 occurred at the Alamosa – ASC monitor and may have been a high wind event as well. The overall trend remains the same whether the 1991 value is included or not. The trend in the 24-hour concentrations over the past three years is increasing but the trend in the annual average concentrations has continued to decline.

Figure 11 - Statewide Ambient Trends – PM₁₀



4.6 Particulates – PM_{2.5}

Monitoring for PM_{2.5} in Colorado began with the establishment of sites in Denver, Grand Junction, Steamboat Springs, Colorado Springs, Greeley, Fort Collins, Platteville, Boulder, Longmont and Elbert County in 1999. Additional sites were established nearly every month until full implementation of the base network was achieved in July of 1999. In 2004, there were 20 PM_{2.5} monitoring sites in Colorado. Thirteen of the 20 sites were selected based on the population of the metropolitan statistical areas. This is a federal selection criterion that was developed to protect the public health in the highest population centers. In addition, there were seven special purpose-monitoring (SPM) sites. These sites were selected due to historically elevated concentrations of PM₁₀ or because citizens or local governments had concerns of possible high PM_{2.5} concentrations in their communities. All SPM sites were removed as of December 31, 2006 due to low concentrations and a lack of funding.

Only one site in Colorado has exceeded the level of the 24-hour standard and no sites have exceeded the level of the annual standard. The Denver CAMP site exceeded the 24-hour level of the standard twice in 2001. The exceedances occurred on Thursday, February 15, 2001 (68.4 $\mu\text{g}/\text{m}^3$) and Saturday, February 17, 2001 (68.0 $\mu\text{g}/\text{m}^3$). The 24-hour standard was lowered by EPA on September 20, 2006 to 35 $\mu\text{g}/\text{m}^3$. This new standard did not become effective until December 17, 2006. Since the previous standard of 65 $\mu\text{g}/\text{m}^3$ applied for most of 2006 it has been used throughout this report. The EPA will use the 2004-2006 PM_{2.5} data to compare sites to the new 24-hour standard. Several sites have exceeded the level of the new standard. However, no sites have violated the 3-year average for the standard.

4.7 Lead

In Colorado the last violation of the federal lead standard occurred in the first quarter of 1980 at the Denver CAMP monitor. Since then, the concentrations recorded at all monitors have shown a steady decline, to the point where now all monitors are regularly at or near the minimum detectable limits of analysis. This decline is the direct result of the use of unleaded gasoline and replacement of older cars with newer ones that do not require leaded gasoline. The reduction in atmospheric lead shows what pollution control strategies can accomplish.

Table 10 - Historical Maximum Quarterly Lead Concentrations³⁷

Quarterly Maximum $\mu\text{g}/\text{m}^3$	Monitor	Date
3.47	Denver CAMP, 2105 Broadway	1 st Qtr 1979
3.40	Denver, 414 14 th St.	4 th Qtr 1969
3.03	Denver, 414 14 th St.	1 st Qtr 1973
3.03	Denver CAMP, 2105 Broadway	4 th Qtr 1978
3.02	Denver, 414 14 th St.	4 th Qtr 1972
2006 Maximum Quarterly Lead Concentration		
0.16	Globeville	1 st Qtr 2006

5.0 National Comparisons For Criteria Pollutants

5.1 Carbon monoxide

According to the Environmental Protection Agency's emissions trends report: Between 1990 and 2006, national average ambient carbon monoxide concentrations decreased 62 percent. Total carbon monoxide emissions decreased 38 percent (excluding wildfires and prescribed burning) for the same period. This improvement in air quality occurred despite a 32-percent increase in vehicle miles traveled during the 10-year period."³⁸

Table 11 - 2006 National Ranking of Carbon Monoxide Monitors by 8-Hr Concentrations in ppm³⁹

Nationwide (406 monitors)					Colorado (13 Monitors)				
National Rank	City/Area	Max	2 nd Max	# >9.5	Nat'l Rank	City/Area	Max	2 nd Max	# >9.5
1	Portland, OR	14.1	7.5	1	51	Greeley	3.4	3.3	0
2	Calexico, CA	9.8	6.6	1	52	CAMP	3.4	3.1	0
3	Birmingham, AL	9.6	9.5	2	76	Firehouse #6	3.1	2.6	0
4	Weirton, WV	8.5	3.9	0	86	Carriage	3.0	3.0	0
5	Anchorage, AK	6.5	6.1	0	102	NJH-E	2.9	2.5	0

5.2 Ozone

Over the past 30 years, EPA, in conjunction with state and local agencies, has instituted various programs to reduce oxides of nitrogen (NO_x) and volatile organic compounds (VOC) emissions that contribute to ozone formation. These emission reductions occurred at the same time the nation's economy, energy consumption, and population were growing. For example, between 1970 and 2003, gross domestic product increased approximately 176 percent; vehicle miles traveled (VMT), 155 percent; energy consumption, 45 percent; and population, 39 percent, whereas emissions of NO_x and VOCs decreased approximately 25 percent and 54 percent, respectively. The ratio of NO_x and VOC emissions to population has also dropped since 1970.⁴¹

This year, both the 1-hour and the 8-hour ozone national rankings have been included. The fourth maximum value is included in the 8-hour table because that is the value that is compared to the standard. The ozone standard is set at 0.08 ppm as the 3-year average of the annual 4th maximum 8-hour average concentration.

Table 12 - 2006 National Ranking of Ozone Monitors by 1-Hr Concentrations in ppm⁴⁰

Nationwide (1,189 Monitors)					Colorado (15 Monitors)				
National Rank	City/Area	Max	2 nd Max	Days >0.125	National Rank	City/Area	Max	2 nd Max	Days >0.125
1	Glendora, CA	0.175	0.155	11	253	Welch	0.112	0.098	0
2	Perris, CA	0.169	0.155	12	287	Ft. Collins	0.109	0.106	0
3	Riverhead, NY	0.168	0.146	3	308	NREL	0.109	0.100	0
4	Burbank, CA	0.166	0.152	6	335	Carriage	0.108	0.090	0
5	Upland, CA	0.166	0.151	14	432	Chatfield Res.	0.103	0.102	0

Table 13 - 2006 National Ranking of Ozone Monitors by 8-Hr Concentrations in ppm⁴¹

Nationwide (1,189 Monitors)					Colorado (15 Monitors)				
National Rank	City/Area	Max	4 th Max	Days >0.085	National Rank	City/Area	Max	4 th Max	Days >0.085
1	Crestline, CA	0.142	0.111	59	204	Ft Collins - West	0.097	0.087	6
2	Redlands, CA	0.135	0.125	36	223	Rocky Flats - N	0.096	0.090	5
3	Upland, CA	0.131	0.112	25	226	Welch	0.096	0.081	2
4	Narragansett, RI	0.130	0.081	3	253	NREL	0.094	0.083	2
5	Riverhead, NY	0.130	0.101	5	324	Chatfield Res.	0.092	0.086	4

5.3 Sulfur Dioxide

“Nationally, average sulfur dioxide ambient concentrations have decreased 54 percent from 1983 to 2002 and 39 percent over the more recent 10-year period of 1993 to 2002. Sulfur dioxide emissions decreased 33 percent from 1983 to 2002 and 31 percent from 1993 to 2002. Reductions in sulfur dioxide concentrations and emissions since 1990 are due, in large part, to controls implemented under EPA’s Acid Rain Program beginning in 1995.”⁴²

Table 14 - 2006 National Ranking of Sulfur Dioxide Monitors by 24-Hr Concentrations in ppm⁴³

Nationwide (508 Monitors)					Colorado (2 Monitors)				
National Rank	City/Area	Max	2 nd Max	#>0.14	Nat'l Rank	City/Area	Max	2 nd Max	#>0.14
1	Hawaii Volcanoes NP	0.353	0.173	6	373	CAMP	0.009	0.009	0
2	Tahlequah, OK	0.322	0.023	1	395	Welby	0.007	0.006	0
3	Easton, PA	0.156	0.147	2					
4	Gibson Co, IN	0.098	0.086	0					
5	Alcoa, TN	0.098	0.080	0					

5.4 Nitrogen Dioxide

“Since 1983, monitored levels of nitrogen dioxide have decreased 21 percent. These downward trends in national nitrogen dioxide levels are reflected in all regions of the country. Nationally, average nitrogen dioxide concentrations are well below the NAAQS and are currently at the lowest levels recorded in the past 20 years. All areas of the country that once violated the NAAQS for nitrogen dioxide now meet that standard. Over the past 20 years, national emissions of oxides of nitrogen have declined by almost 15 percent. The reduction in emissions for oxides of nitrogen presented here differs from the increase in oxides of nitrogen emissions reported in previous editions of this report. In particular, this report’s higher estimate of oxides of nitrogen emissions in the 1980s and early 1990s reflects an improved understanding of emissions from real-world driving. While overall oxides of nitrogen emissions are declining, emissions from some sources such as nonroad engines have actually increased since 1983. These increases are of concern given the significant role oxides of nitrogen emissions play in the formation of ground-level ozone (smog) as well as other environmental problems like acid rain and nitrogen loadings to water bodies described above. In response, EPA has proposed regulations that will significantly control oxides of nitrogen emissions from nonroad diesel engines.”⁴⁴

Table 15 - 2006 National Ranking of Nitrogen Dioxide Monitors by 1-Hr Concentrations in ppm⁴⁵

Nationwide (424 Monitors)					Colorado (2 Monitors)				
National Rank	City/Area	1-hr Max	2 nd Max	Ann. Avg.	National Rank	City/Area	1-hr Max	2 nd Max	Ann. Avg.
1	Albuquerque, NM	0.284	0.267	0.013	5	CAMP	0.169	0.136	0.029
2	Saint Louis, MO	0.256	0.245	0.009	63	Welby	0.090	0.090	0.019
3	Thomaston, CT	0.203	0.075	0.012					
4	Cleveland, OH	0.175	0.075	0.014					
5	Denver - CAMP	0.169	0.136	0.029					

5.5 Particulates

The monitors recording the three highest PM₁₀ concentrations, in the nation, are located in Owens Valley, California. These levels are associated with the high winds that blow across the dry bed of Owens Lake. In the past seven years monitors in area have recorded levels in excess of 20,000 µg/m³ as a 24-hour average.⁴⁷

Table 16 - 2006 National Ranking of PM₁₀ Monitors by 24-Hr Maximum Concentrations in µg/m³⁴⁶

Nationwide (1,136 Monitors)					Colorado (41 Monitors)				
National Rank	City/Area	1 st Max	2 nd Max	Annual Mean	National Rank	City/Area	1 st Max	2 nd Max	Annual Mean
1	Olancho, CA	8,299	2371	64	19	Alamosa - ASC	424*	158	21.6
2	Lee Vining, CA	4,300	1,915	88	35	Alamosa Municipal	289*	213	25.2
3	Keeler, CA	2,100	756	33	62	Aspen	174*	125	34.2
4	Maricopa, AZ	1,079	794	231	118	Lamar - 100 2nd	136	127	24.0
5	El Paso, TX	712	559	65	140	Fort Collins	130	55	20.7

* - These have been classified as natural events by the APCD. They are the result of high winds and blowing dust.

“PM_{2.5} concentrations can reach unhealthy levels even in areas that meet the annual standard. In 2005 there were 67.6 million people in the U. S. living in counties with levels above the National Ambient Air Quality Standard for PM_{2.5}. Most metropolitan areas had fewer unhealthy PM_{2.5} days in 2005 compared to the average from the previous 3 years, which reflects the improvements observed in 2005.”²⁴

Table 17 - 2006 National Ranking of PM_{2.5} Monitors by 24-Hr Maximum Concentrations in µg/m³⁴⁷

Nationwide (1,217 Monitors)					Colorado (19 Monitors)				
National Rank	City/Area	1 st Max	2 nd Max	Annual Mean	National Rank	City/Area	1 st Max	2 nd Max	Annual Mean
1	Keeler, CA	193	22	6.1	222	Platteville	43	35	9.4
2	Nogales, AZ	103	56	15.6	385	CAMP	38	31	8.5
3	Liberty, PA	101	82	19.0	433	Commerce City	37	28	9.9
4	Atlantic City, NJ	91	45	12.4	435	Swansea	37	35	9.0
5	Fresno, CA	87	65	17.6	492	Arapahoe CC	36	25	7.6

5.6 Lead

The statistic used to track ambient lead air quality is the maximum quarterly mean concentration for each year. “Because of the phase-out of leaded gasoline, lead emissions and concentrations decreased sharply during the 1980s and early 1990s. The 2002 average air quality concentration for lead is 94 percent lower than in 1983. Emissions of lead decreased 93 percent over the 21-year period 1982 to 2002. These large reductions in long-term lead emissions from transportation sources have changed the nature of the ambient lead problem in the United States. Because industrial processes are now responsible for all violations of the lead NAAQS, the lead monitoring strategy currently focuses on emissions from these point sources. Today, the only violations of the lead NAAQS occur near large industrial sources such as lead smelters and battery manufacturers. Various enforcement and regulatory actions are being actively pursued by EPA and the states for cleaning up these sources.”⁴⁸

Table 18 - 2006 National Ranking of Lead Monitors by 24-Hr Maximum Concentration in $\mu\text{g}/\text{m}^3$ ⁴⁹

Nationwide (220 Monitors)					Colorado (6 Monitors)				
National Rank	City/Area	24-hr Max	Max Qtr	Qtrs >1.5	National Rank	City/Area	24-hr Max	Max Qtr	Qtrs >1.5
1	Herculanium, Mo	26.93	1.82	2	33	Globeville	0.87	0.16	0
2 (5)	Iron County, Mo	9.14	1.27	0	101	Commerce City	0.06	0.02	0
3 (6)	Knoxville, Tn	9.00	1.82	1	123	CAMP	0.04	0.02	0
4 (8)	Muncie, In	5.83	0.83	0	124	Leadville	0.04	0.02	0
5 (15)	Tampa, Fl	3.70	0.83	0	140	Animal Shelter	0.03	0.01	0

In Table 18 the ranking is by highest city or area monitored. The reason for this is that six of the top fifteen monitors are located in Herculanium, Mo, three monitors are located in Muncie, IN and two are located in Iron County, Mo. The rank by monitor is shown in parentheses.

6.0 Monitoring Results by Area in Colorado

6.1 Eastern Plains Counties

The Eastern Plains Counties are those east of the urbanized I-25 corridor. Historically, there have been a number of communities that were monitored for particulates. In the northeast along the I-76 corridor, the communities of Sterling, Brush and Fort Morgan have been monitored. Along the I-70 corridor only the community of Limon has been monitored for particulates. In the southeast, the US-50/Arkansas River corridor, the communities of La Junta and Rocky Ford have been monitored in the past, but like the other communities that have been monitored on the Eastern Plains, the monitoring was discontinued when the concentrations were shown to be below the standard.

Currently, there are two PM₁₀ monitors in Lamar and a background PM_{2.5} monitor in Elbert County at the Wright-Ingraham Institute. The Lamar monitors have recorded exceedances in the 24-hour PM₁₀ standard. These have been associated with high winds and dry conditions that can occur in springtime. The Elbert County monitor operates as a background PM_{2.5} monitor. This monitor provides baseline PM_{2.5} readings away from any influence of man made particulate sources.

Table 19 - Eastern Plains Monitors In Operation For 2006
 X - Monitors continued in 2006 A – Monitors added in 2006
 D – Monitors discontinued in 2006 H – Hourly particulate monitor

Site Name	Location	PM ₁₀	PM _{2.5}	Met
Elbert				
Elbert	Wright-Ingraham Inst		X	
Prowers				
Lamar	100 2 nd St.	X		
	104 Parmenter St.	X		
Lamar Port of Entry	7100 US Hwy 50			X

Table 20 - Eastern Plains Particulate Values For 2006

Location	PM ₁₀ (µg/m ³)		PM _{2.5} (µg/m ³)	
	Annual Average	24-hour Maximum	Annual Average	24-hour Maximum
Elbert				
Wright-Ingraham Inst			(4.11)	20.2
Prowers				
100 2 nd St.	24.0	136		
104 Parmenter St.	19.5	116		

() indicates <75 percent data recovery in one or more quarters.

Figure 12 - Eastern Plains PM₁₀ Particulate Graphs




Figure 13 - Eastern Plains PM_{2.5} Particulate Graph





Figure 14 - Eastern Plains Wind Rose Graph
 Lamar Port of Entry, 7100 US Hwy



50 © 2000 Microsoft Corp. and/or its suppliers. All rights reserved.

6.2 Northern Front Range Counties

The Northern Front Range Counties are those along the urbanized I-25 corridor from the Colorado/Wyoming border to just south of the city of Castle Rock. This area has the majority of the larger cities in the state. The majority of monitors are located in the Denver-metro area and the rest are located in or near Fort Collins, Greeley, Longmont and Boulder.

Table 21 - Northern Front Range Particulate Monitors In Operation For 2006

X - Monitors continued in 2006 A – Monitors added in 2006

D – Monitors discontinued in 2006 H – Hourly particulate monitor S – Chemical Speciation

Site Name	Location	TSP	Pb	PM ₁₀	PM _{2.5}
Adams					
Brighton	22 S. 4 th Ave.			D	
Commerce City	7101 Birch St.	D	D	X	X/H/S
Globeville	5400 Washington St.	D	D		
Welby	3174 E. 78 th Ave.			X/H	
Arapahoe					
Arapahoe Community Coll.	6190 S. Santa Fe Dr.				X
Boulder					
Boulder	2440 Pearl St.			X	X
	2102 Athens St.				H
Longmont	350 Kimbark St.			X	X/H
Denver					
Denver CAMP	2105 Broadway	D	D	X/H	X/H
Denver - NJH	14 th Ave. & Albion St.				H
Denver Visitor Center	225 W. Colfax Ave.			X	
Lowry	8100 Lowry Blvd.			D	
Swansea Elementary School	4650 Columbine St.				X
Denver Animal Shelter	678 S. Jason St.	A	A	A/H	A
Douglas					
Chatfield Reservoir	11500 Roxborough Pk. Rd.				X/H
Larimer					
Fort Collins	251 Edison St.			X	X
Weld					
Greeley	1516 Hospital Rd.			X	X/H
Platteville	1004 Main St.				X/S


Table 22 - Northern Front Range Particulate Values For 2006

Site Name	PM ₁₀ (µg/m ³)		PM _{2.5} (µg/m ³)	
	Annual Average	24-hour Maximum	Annual Average	24-hour Maximum
Adams				
Brighton	22.6	46		
Commerce City (Continuous Monitor)	35.0	115	9.89	36.7
			7.76	22.9
Welby (Continuous Monitor)	27.8	50		
	27.4	83		
Arapahoe				
Arapahoe Community Coll.			7.64	36.1
Boulder				
Boulder, 2440 Pearl St.	17.4	34	6.72	22.0
Boulder, 2102 Athens St. (Continuous Monitor)			(5.78)	24.2
Longmont (Continuous Monitor)	20.0	37	7.98	30.5
			(9.32)	20.4
Denver				
Denver CAMP (Continuous Monitor)	27.3	59	8.46	37.8
	(23.5)	85	13.05	43.4
Denver – NJH (Continuous Monitor)			9.77	39.6
Denver Visitor Center	24.0	79		
Lowry	20.2	55		
Swansea Elementary School			8.99	36.7
Denver Animal Shelter (Continuous Monitor)	25.2	52		
	(27.8)	88		
Douglas				
Chatfield Reservoir (Continuous Monitor)			6.18	19.7
			(6.32)	23.0
Larimer				
Fort Collins	20.7	130	7.30	28.8
Weld				
Greeley (Continuous Monitor)	22.4	55	8.21	33.8
			6.84	23.4
Platteville			9.37	42.6


() Indicates less than 75 percent data for one or more quarters.

Figure 15 - Northern Front Range PM₁₀ Particulate Graphs


Brighton, 22 4th Ave. - Ambient Trends - PM10




Commerce City, 7101 Birch St. - Ambient Trends - PM10




Welby, 3174 E. 78th Ave. - Ambient Trends - PM10



Boulder, 2440 Pearl St. - Ambient Trends - PM10



Longmont, 350 Kimbark St. - Ambient Trends - PM10



Denver CAMP, 2150 Broadway - Ambient Trends - PM10






Figure 15 - Northern Front Range PM₁₀ Particulate Graphs (continued)


Denver Gates/Animal Shelter - Ambient Trends - PM10




Denver Visitor Center, 225 W. Colfax Ave - Ambient Trends PM10



Denver Lowry, 8100 Lowry Blvd - Ambient Trends PM10



Fort Collins, 251 Edison St. - Ambient Trends - PM10



Greeley, 1516 Hospital Rd. - Ambient Trends - PM10




Figure 16 - Northern Front Range PM_{2.5} Particulate Graphs









Figure 16 - Northern Front Range PM_{2.5} Particulate Graphs (continued)




Table 23 - Northern Front Range TSP and Lead Values For 2006

Site Name	Location	TSP (µg/m ³)		Lead (µg/m ³)	
		Annual Geometric Mean	24-hour Maximum	Maximum Quarter	24-hour Maximum
Adams					
Adams	Commerce City	74.8	167	0.015	0.063
	Globeville	81.6	168	0.158	0.873
Denver					
Denver	Denver CAMP	66.3	146	0.016	0.044
	Denver Animal	51.2	109	0.095	0.026

() indicates less than 75 percent data for one or more quarters.

Figure 17 - Northern Front Range Lead Graphs







Table 24 - Northern Front Range Continuous Monitors In Operation For 2006
X - Monitors continued in 2006 A – Monitors added in 2006 D – Monitors discontinued in 2006

Site Name	Location	CO	SO ₂	NO _x	O ₃	Met
Adams						
Commerce City	7101 Birch St.					X
Welby	3174 E. 78 th Ave.	X	X	X	X	X
Arapahoe						
Highland Res.	8100 S. University Blvd.				X	X
Boulder						
Boulder	1405½ S. Foothills Hwy.				X	
Longmont	440 Main St.	X				
Denver						
Auraria Lot R	12 th St. & Auraria Parkway					X
Denver - CAMP	2105 Broadway	X	X	X	X	X
Denver - Carriage	2325 Irving St.	D			X	X
Denver - NJH	14 th Ave. & Albion St.	D				
Firehouse #6	1300 Blake St.	X				
Douglas						
Chatfield Reservoir	11500 N. Roxborough Pk. Rd.				X	X
Jefferson						
Arvada	9101 W. 57 th Ave.	D			X	X
NREL	2229 Old Quarry Rd.				X	
Rocky Flats - N	16600 W. Hwy. 128				X	X
Rocky Flats - SE	9901 Indiana St.					X
Welch	12400 W. Hwy. 285				X	X
Larimer						
Fort Collins	708 S. Mason St.	X			X	X
	4407 S. College Ave.	X				
	3416 LaPorte Ave.				X	
Weld						
Greeley	905 10 th Ave.	X				
	3101 35 th Ave.				X	

Table 25 - Northern Front Range Carbon Monoxide Values for 2006

Site Name	Location	CO 1-hour Avg. (ppm)		CO 8-hour Avg. (ppm)	
		Max	2 nd Max	Max	2 nd Max
Adams					
Welby	3174 E. 78 th Ave.	3.8	3.8	2.6	2.5
Boulder					
Longmont	440 Main St.	3.9	2.8	2.2	1.8
Denver					
Denver - CAMP	2105 Broadway	6.4	4.6	3.4	3.1
Denver - Carriage	2325 Irving St.	3.9	3.5	3.0	3.0
Denver - NJH	14 th Ave. & Albion St.	4.4	3.9	2.9	2.5
Firehouse #6	1300 Blake St.	9.3	5.7	3.1	2.6
Jefferson					
Arvada	9101 W. 57 th Ave.	3.6	3.5	2.0	2.0
Larimer					
Fort Collins	708 S. Mason St.	4.4	3.9	2.8	2.7
	4407 S. College Ave.	3.2	3.0	1.9	1.8
Weld					
Greeley	905 10 th Ave.	6.4	5.6	3.4	3.3

Figure 18 - Northern Front Range Carbon Monoxide Graphs




Figure 18 - Northern Front Range Carbon Monoxide Graphs (continued)









Figure 18 - Northern Front Range Carbon Monoxide Graphs (continued)




Table 26 - Northern Front Range Ozone Values For 2006

Site Name	Location	Ozone 1-hour Avg. (ppm)		Ozone 8-hour Avg. (ppm)	
		Maximum	2 nd Maximum	Maximum	4 th Maximum
Adams					
Welby	3174 E. 78 th Ave.	0.089	0.087	0.081	0.069
Arapahoe					
Highland Res.	8100 S. University Blvd.	0.097	0.095	0.085	0.081
Boulder					
Boulder	1405½ S. Foothills Hwy.	0.099	0.098	0.087	0.082
Denver					
Denver - CAMP	2105 Broadway	0.085	0.079	0.071	0.062
Carriage	2325 Irving St.	0.108	0.090	0.092	0.072
Douglas					
Chatfield Res.	11500 Roxborough Park Rd.	0.103	0.102	0.092	0.086
Jefferson					
Arvada	9101 W. 57 th Ave.	0.099	0.094	0.083	0.082
NREL	2229 Old Quarry Rd.	0.109	0.100	0.094	0.083
Rocky Flats	16600 W. Hwy 128	0.104	0.103	0.094	0.090
Welch	12400 W. Hwy 285	0.112	0.098	0.096	0.081
Larimer					
Fort Collins	708 S. Mason St.	0.100	0.094	0.084	0.078
	3416 LaPorte Ave.	0.109	0.106	0.097	0.087
Weld					
Greeley	3101 35 th Ave.	0.103	0.100	0.090	0.082

Figure 19 - Northern Front Range Ozone Graphs









Figure 19 - Northern Front Range Ozone Graphs (continued)








Table 27 - Northern Front Range Oxides of Nitrogen and Sulfur Dioxide Values For 2006

Site Name	Location	Nitrogen Dioxide	Nitric Oxide	Sulfur Dioxide		
		Annual Avg. (ppm)	Annual Avg. (ppm)	3-hour 2 nd Max (ppm)	24-hour 2 nd Max (ppm)	Annual Avg. (ppm)
Adams						
Welby	3174 E. 78 th Ave.	0.0192	0.026	0.025	0.006	0.0020
Denver						
Denver CAMP	2105 Broadway	0.0294	0.035	0.022	0.009	0.0029

Figure 20 - Northern Front Range Nitrogen Dioxide Graphs




Figure 21 - Northern Front Range Sulfur Dioxide Graphs




Table 28 - Denver Visibility Standard Exceedance Days
(Transmissometer Data)
January 2006 – December 2006

Month	Days	EX POOR	POOR	FAIR	GOOD	Missing	(>70% RH)
January	31					31	
February	28					28	
March	31					31	
April	30					30	
May	31					31	
June	30		6	2	1	21	
July	31	1	15	10	3		2
August	31	1	8	9	13		
September	30	3	11	9	6		1
October	31		10	9	8		4
November	30	3	4	14	7		2
December	31	2	4	9	11		5
Totals	365	10	58	62	49	172	14

Figure 22 shows that 47 percent or of the visibility data for 2006 was listed as “Missing.” The data from January through June 22, 2006 were removed after the annual data review. The review showed a series of inconsistent baseline values that did not become apparent until the instrument was returned from its annual calibration in June 2006. Further comparisons with PM_{2.5} samples, meteorological measurements and visual analysis raised additional questions regarding the validity of the data for the same period. The Division has instituted procedural changes to reduce the possibility of this much data loss in the future.

Figure 22 - Denver Visibility Data (January 2006 to December 2006)




Figure 23 - Denver Visibility Comparison (1997 to 2006)




Table 29 - Fort Collins Visibility Standard Exceedance Days
(Transmissometer Data)
January 2006 – December 2006

Month	Days	EX POOR	POOR	FAIR	GOOD	Missing	(>70% RH)
January	31		4	7	19		1
February	28		6	7	8	6	1
March	31		6	7	7	5	6
April	30		5	15	10		
May	31		4	16	9	2	
June	30	1	10	10	2	7	
July	31		21	7		3	
August	31	1	11	8	7	4	
September	30	1	13	9	3	4	
October	31		9	10	1	11	
November	30		8	5	9	7	1
December	31		6	2	2	14	7
Totals	365	3	103	103	77	63	16

The transmissometer was removed from the site for its annual servicing and calibration from June 7 through August 10. The other cause for missing data was normal maintenance of the instrument.

Figure 24 - Fort Collins Visibility Data (January 2006 to December 2006)




Figure 25 shows that since 1997 the Fort Collins has averaged 176 days per year where the visibility was either “Fair” or “Good” and only 97 days where the visibility was either “Poor” or “Ex Poor”. The missing days are lost due to either high relative humidity (greater than 70 percent) or machine maintenance.

Figure 25 - Fort Collins Visibility Data (1997 to 2006)





Figure 26 - Northern Front Range Wind Roses
 Arvada, 9101 W. 57th Ave.



Auraria, Parking Lot R





Figure 26 - Northern Front Range Wind Roses (continued)
 Chatfield Reservoir, 11500 N. Roxborough Pk. Rd.



Commerce City, 7101 Birch St.





Figure 26 - Northern Front Range Wind Roses (continued)
 Denver CAMP, 2105 Broadway



Denver Carriage, 2325 Irving St.





Figure 26 - Northern Front Range Wind Roses (continued)
Fort Collins, 708 S. Mason St.



Highland Reservoir, 8100 S. University Blvd.





Figure 26 - Northern Front Range Wind Roses (continued)
 Rocky Flats-N, 16600 W. Hwy. 128



Rocky Flats- SE, 9901 Indiana St.






Figure 26 - Northern Front Range Wind Roses (continued)
 Welby, 3174 E. 78th Ave.



Welch, 12400 W. Hwy. 285



6.3 Southern Front Range Counties

The Southern Front Range Counties are those along the urbanized I-25 corridor from south of the city of Castle Rock to the southern Colorado border. The cities with monitoring in the area are Colorado Springs, Pueblo, Cripple Creek, Cañon City and Alamosa. These last three cities are not strictly in the Front Range I-25 corridor but fit better with those cities than they do the Mountain Counties. Colorado Springs is the only city in the area that is monitored for carbon monoxide and ozone by the APCD. The other cities are only monitored for particulates. In the past the APCD has conducted particulate monitoring in both Walsenburg and Trinidad but that monitoring was discontinued in 1979 and 1985 respectively.

Table 30 - Southern Front Range Monitors In Operation For 2006

X - Monitors continued in 2006 A – Monitors added in 2006

D – Monitors discontinued in 2006 H – Hourly particulate monitor S – Chemical Speciation

Site Name	Location	CO	O ₃	TSP	Pb	PM ₁₀	PM _{2.5}	Met
Alamosa								
Alamosa	359 Edgemont Blvd.					X		
	425 4 th St.					X		
El Paso								
Colorado Springs	1098 Glenn St.	D						
	3730 Meadowlands					D	D	
	101 W. Costilla St.			D	D	X	X/S	
	USAF Rd. 640		X					
	690 W. Hwy. 24	X						
Manitou Springs	101 Banks Pl.		X					
Fremont								
Cañon City	128 Main St.					X		
Pueblo								
Pueblo	211 E. D St.					X	X	
Teller								
Cripple Creek	209 Bennett Ave.					D		
	Warren Ave. & 2 nd St							D

Table 31 - Southern Front Range Maximum Particulate Values For 2006

Site Name	Location	PM ₁₀ (µg/m ³)		PM _{2.5} (µg/m ³)	
		Annual Average	24-Hr Maximum	Annual Average	24-Hr Maximum
Alamosa					
Alamosa	208 Edgemont Blvd.	21.6	424		
	425 4 th St.	(25.2)	289		
El Paso					
Colorado Springs	3730 Meadowlands	21.4	53	6.71	19.0
	101 W. Costilla St.	22.6	56	8.14	19.1
Fremont					
Cañon City	128 Main St.	16.9	54		
Pueblo					
Pueblo	211 E. D St.	23.2	58	(8.72)	19.2
Teller					
Cripple Creek	209 Bennett Ave.	15.6	46		

() Indicates less than 75 percent data for one or more quarters.

Figure 27 - Southern Front Range PM₁₀ Particulate Graphs







Figure 27 - Southern Front Range PM₁₀ Particulate Graphs (continued)


Colorado Springs, 3730 Meadowlands - Ambient Trends - PM10




Colorado Springs, 101 W. Costilla St. - Ambient Trends - PM10



Canon City - Ambient Trends - PM10



Pueblo - Ambient Trends - PM10



Cripple Creek, 209 Bennet Ave. - Ambient Trends - PM10




Figure 28 - Southern Front Range PM_{2.5} Particulate Graphs






Table 32 - Southern Front Range TSP and Lead Values For 2006

Site Name	Location	TSP (µg/m ³)		Lead (µg/m ³)	
		Annual Geometric Mean	24-Hr Maximum	Maximum Quarter	24-Hr Maximum
El Paso					
Colorado Springs	101 W. Costilla St.	48.3	134	0.006	0.014

Figure 29 - Southern Front Range Lead Graph




Table 33 - Southern Front Range Carbon Monoxide Values For 2006

Site Name	Location	CO 1-hour Avg. (ppm)		CO 8-hour Avg. (ppm)	
		Maximum	2 nd Maximum	Maximum	2 nd Maximum
El Paso					
Colorado Springs	1098 Glenn St.	4.4	3.6	2.0	1.8
	690 Hwy. 24	4.3	4.0	2.6	2.4

Figure 30 - Southern Front Range Carbon Monoxide Graphs





Table 34 - Southern Front Range Ozone Values For 2006

Site Name	Location	Ozone 1-hour Avg. (ppm)		Ozone 8-hour Avg. (ppm)	
		Maximum	2 nd Maximum	Maximum	4th Maximum
El Paso					
Colorado Springs	USAFA Rd. 640	0.080	0.079	0.073	0.072
Manitou Springs	101 Banks Pl.	0.089	0.089	0.079	0.076

Figure 31 - Southern Front Range Ozone Graph





Figure 32 - Southern Front Range Wind Rose
Cripple Creek, Warren Ave. & 2nd St.



6.4 Mountain Counties

The Mountain Counties are generally the towns near the Continental Divide. They are mostly small towns in tight mountain valleys. Their primary monitoring concern is with particulate pollution from wood burning and road sanding. These communities range from Steamboat Springs in the north, to Silverthorne, Vail and Breckenridge in the I-70 corridor, Aspen, Leadville, Crested Butte, Mt. Crested Butte and Gunnison in the central mountains to Telluride in the southwest.

Table 35 - Mountain Counties Monitors In Operation For 2006

X - Monitors continued in 2006 A – Monitors added in 2006

D – Monitors discontinued in 2006 H – Hourly particulate monitor

Site Name	Location	TSP	Pb	PM ₁₀	PM _{2.5}
Archuleta					
Pagosa Springs	309 Lewis St.			X	X
Gunnison					
Crested Butte	603 6 th St.			X	
Gunnison	211 Wisconsin Ave.			D	
Mt. Crested Butte	19 Emmons Loop			X	X
Lake					
Leadville	510 Harrison St.	D	D		
Pitkin					
Aspen	120 Mill St.			X/H	
Routt					
Steamboat Springs	136 6 th St.			X	
Summit					
Breckenridge	501 N. Park Ave.			X	

Table 36 - Mountain Counties Particulate Values For 2006

Site Name	Location	PM ₁₀ (µg/m ³)		PM _{2.5} (µg/m ³)	
		Annual Average	24-Hr Maximum	Annual Average	24-Hr Maximum
Archuleta					
Pagosa Springs	309 Lewis St.	19.8	122	4.58	7.0
Gunnison					
Crested Butte	603 6 th St.	27.1	100		
Gunnison	211 Wisconsin Ave.	17.4	55		
Mt. Crested Butte	19 Emmons Loop	29.1	120	6.42	21.7
Pitkin					
Aspen	120 Mill St.	(17.4)	57		
	(Continuous Monitor)	17.4	58		
Routt					
Steamboat Springs	136 6 th St.	23.4	87		
Summit					
Breckenridge	501 N. Park Ave.	16.7	86		

() Indicates less than 75 percent data for one or more quarters.

Figure 33 - Mountain Counties PM₁₀ Particulate Graphs




Figure 33 - Mountain Counties PM₁₀ Particulate Graphs (continued)








Figure 34 - Mountain Counties PM_{2.5} Particulate Graphs






Table 37 - Mountain Counties TSP and Lead Concentrations For 2006

Site Name	Location	TSP ($\mu\text{g}/\text{m}^3$)		Lead ($\mu\text{g}/\text{m}^3$)	
		Annual Geometric Mean	24-Hr Maximum	Maximum Quarter	24-Hr Maximum
Lake					
Leadville	510 Harrison St.	26.2	129	0.016	0.044

Figure 35 - Mountain Counties Lead Graphs



6.5 Western Counties

The Western Counties are generally smaller towns in fairly broad river valleys. Grand Junction is the only large city in the area and the only location that monitors for carbon monoxide on the western slope. The other locations monitor only for particulates. They are located in Parachute, Delta, Durango and Pagosa Springs.

Table 38 - Western Counties Monitors In Operation For 2006

X - Monitors continued in 2006 A – Monitors added in 2006

D – Monitors discontinued in 2006 H – Hourly particulate monitor S – Chemical Speciation

Site Name	Location	CO	PM ₁₀	PM _{2.5}	Met
Delta					
Delta	560 Dodge St.		X	D	
Garfield					
Parachute	100 E. 2 nd Ave.		X		
Rifle	144 E. 3 rd Ave.		X		
New Castle	402 W. Main St.		X		
Silt – Bell Ranch	512 Owens Dr.		X		
Silt – Daley Ranch	884 County Rd. 327		X		
Silt – Cox Ranch	5933 County Rd. 233		X		
Glenwood Springs	109 8 th St.		X		
La Plata					
Durango	1060 2 nd Ave.		D		
	56 Davidson Creek Rd.		D		
	1235 Camino Del Rio		X		
	117 Cutler Dr.		D		
Mesa					
Grand Junction	650 South Ave.		X	X/H	
	645¼ Pitkin Ave.	X	H		X/S
San Miguel					
Telluride	333 W. Colorado Ave.		X	D	

Table 39 - Western Counties Particulate Values For 2006

Site Name	Location	PM ₁₀ (µg/m ³)		PM _{2.5} (µg/m ³)	
		Annual Average	24-Hr Maximum	Annual Average	24-Hr Maximum
Delta					
Delta	560 Dodge St.	24.3	53	7.10	18.0
Garfield					
Parachute	100 E. 2 nd Ave.	30.2	76		
Rifle	144 E. 3 rd Ave.	29.5	67		
New Castle	402 W. Main St.	23.0	53		
Silt – Bell Ranch	512 Owens Dr.	11.4	28		
Silt – Daley Ranch	884 County Rd. 327	9.7	20		
Silt – Cox Ranch	5933 County Rd. 233	13.6	26		
Glenwood Springs	106 8 th St.	14.6	31		
La Plata					
Durango	1060 2 nd Ave.	15.2	46		
	56 Davidson Creek Rd.	(15.5)	39		
	1235 Camino Del Rio	18.9	47		
	117 Cutler Dr.	(14.4)	34		
Mesa					
Grand Junction	650 South Ave.	30.2	98	9.70	28.5
(Continuous Monitor)				(9.33)	18.2
(Continuous Monitor)	645¼ Pitkin Ave.	34.2	174		
San Miguel					
Telluride	333 W. Colorado Ave.	18.9	69	(4.14)	16.3

() Indicates less than 75 percent data for one or more quarters.

Figure 36 - Western Counties PM₁₀ Particulate Graphs





Figure 36 - Western Counties PM₁₀ Particulate Graphs (continued)








Figure 37 - Western Counties PM_{2.5} Particulate Graph






Table 40 - Western Counties Carbon Monoxide Values For 2006

Site Name	Location	CO 1-hour Avg.(ppm)		CO 8-hour Avg.(ppm)	
		Maximum	2 nd Maximum	Maximum	2 nd Maximum
Mesa					
Grand Junction	645¼ Pitkin Ave.	3.0	2.8	1.9	1.7

Figure 38 - Western Counties Carbon Monoxide Graph





Figure 39 - Western Counties Wind Roses
Grand Junction, 645¼ Pitkin Ave.



References

- 1 National Primary and Secondary Ambient Air Quality Standards, Title 40 Code of Federal Regulation, Pt. 50. 1999 ed.
- 2 United States Environmental Protection Agency – Air Trends, Carbon Monoxide, March 1, 2006
- 3 National Primary and Secondary Ambient Air Quality Standards for Carbon Monoxide, Title 40 Code of Federal Regulations, Pt. 50.8. 1999 ed.
- 4 Air Pollution Control Division, Colorado Implementation Plan for Carbon Monoxide Support Document, Chapter #6, Colorado Department of Public Health and Environment, October, 1994.
- 5 United States Environmental Protection Agency Air Emissions Trends – Carbon Monoxide http://www.epa.gov/air/airtrends/2006/emissions_summary_2005.html
- 6 United States Environmental Protection Agency Ground Level Ozone, <http://www.epa.gov/air/airtrends/ozone.html>
- 7 Environmental Protection Agency, National Ambient Standards for Ozone: Final Rule, Title 40 Code of Federal Regulations Pt. 50, July 18, 1997.
- 8 United States Environmental Protection Agency – Air Trends, Ozone, March 1, 2006.
- 9 National Primary and Secondary Ambient Air Quality Standards for Sulfur Dioxide, Title 40 Code of Federal Regulations, Pt. 50.5. 1999 ed.
- 10 United States Environmental Protection Agency – Air Trends, Sulfur Dioxide, March 1, 2006.
- 11 United States Environmental Protection Agency – Basic Information – Six Common Pollutants Sulfur Dioxide, March 6, 2007
- 12 United States Environmental Protection Agency Air Emissions Trends – Sulfur Dioxide http://www.epa.gov/air/airtrends/2006/emissions_summary_2005.html
- 13 Environmental Protection Agency Strategies and Air Standards Division, Preliminary Assessment of Health and Welfare Effects Associated with Nitrogen Oxides for Standards Setting Purposes, U.S. Government Printing Office, Washington: October 1981, pp i-iii.
- 14 Environmental Protection Agency, Aerometric Information Retrieval System, Quick Look Report, Nitrogen Dioxide, June 2007.
- 15 Air Pollution Control Division, Mobile Sources Vehicles Travel Projections, January 1997
- 16 United States Environmental Protection Agency – Air Trends, Nitrogen Dioxide, March 1, 2006.
- 17 AirNow - Particulate Pollution and Your Health, What is particulate pollution, www.airnow.gov/index.cfm/actionparticle.airborne

-
- 18 National Primary and Secondary Ambient Air Quality Standards for Particulate Matter, Title 40 Code of Federal Regulation, Pt. 50.7. 1999 ed.
- 19 The Perils of Particulates, American Lung Association, New York, New, York, March 1994.
- 20 United States Environmental Protection Agency – Air Trends, Particulate Matter, May 4, 2004.
- 21 United States Environmental Protection Agency Air Emissions Trends – PM₁₀
http://www.epa.gov/air/airtrends/2006/emissions_summary_2005.html
- 22 Air Quality – Particulate Matter, www.fort-collins.co.us/airquality/particulate-matter.php,
April 18, 2002.
- 23 United States Environmental Protection Agency Air Emissions Trends – PM_{2.5}
http://www.epa.gov/air/airtrends/2006/emissions_summary_2005.html
- 24 United States Environmental Protection Agency – Basic Information – Six Common Pollutants Lead, March 6, 2007
- 25 National Primary and Secondary Ambient Air Quality Standards for Lead, Title 40 Code of Federal Regulation, Pt. 50.7. 1999 ed.
- 26 United States Environmental Protection Agency – Air Trends, Lead, May 4, 2004.
- 27 United States Environmental Protection Agency Air Emissions Trends – Lead
http://www.epa.gov/air/airtrends/2006/emissions_summary_2005.html
- 28 Clean Air Act as amended in 1977, section 169a (42 USC 7491)
- 29 Visibility Protection for Federal Class I Areas, codified at Title 40 Code of Federal Regulations, Pt. 51.300-309.
- 30 Environmental Protection Agency, Air Quality Criteria for Oxides of Nitrogen, December, 1982, AP 600/8-82-029aF.
- 31 United States Environmental Protection Agency – Air Trends, Toxic Air Pollutants, May 4, 2004.
- 32 Environmental Protection Agency, Air Quality System, Quick Look Report, Carbon Monoxide, June 2007.
- 33 Environmental Protection Agency, Air Quality System, Quick Look Report, Ozone, June 2007.
- 34 Environmental Protection Agency, Air Quality System, Quick Look Report, Sulfur Dioxide, June 2007.
- 35 Environmental Protection Agency, Air Quality System, Quick Look Report, Nitrogen Dioxide, June 2007.
- 36 Environmental Protection Agency, Air Quality System, Quick Look Report, Particulates, June 2007.

-
- 37 Environmental Protection Agency, Air Quality System, Quick Look Report, Lead, June 2007.
- 38 United States Environmental Protection Agency – Air Trends, Carbon Monoxide, June 2007.
- 39 United States Carbon Monoxide Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].
- 40 United States Ozone Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].
- 41 United States Ozone Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].
- 42 United States Environmental Protection Agency – Air Trends, Sulfur Dioxide, May 4, 2004.
- 43 United States Sulfur Dioxide Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].
- 44 United States Environmental Protection Agency – Air Trends, Nitrogen Dioxide, May 4, 2004.
- 45 United States Nitrogen Dioxide Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].
- 46 United States PM₁₀ Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].
- 47 United States PM_{2.5} Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].
- 48 United States Environmental Protection Agency – Air Trends, Lead, May 4, 2007.
- 49 United States Lead Air Quality Monitors (2006), Monitor Ranking Report, Available: www.epa.gov/air/data/monvals.html. Ranking [June 2007].