

Headline®

fungicide

Supplemental Labeling

EPA Reg. No. 7969-186

EPA SLN No.: CO-050003

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF COLORADO

FOR THE CONTROL OF HEAD MOLD IN BARLEY

This label for Headline® fungicide expires and must not be distributed or used in accordance with this FIFRA 24(c) Special Local Need registration after January 1, 2007.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered product label for Headline® fungicide, EPA Reg. No. 7969-186.

This labeling must be in the possession of the user at the time of application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Read the **Precautionary Statement, Environmental Hazards, Storage and Disposal** statements, and **Conditions of Sale and Warranty** statement appearing on the container label.

GENERAL INFORMATION

Headline fungicide provides optimum disease control when applied in a regularly scheduled protective fungicide program and is used in a spray program that rotates fungicides with different modes of action. Refer to the **Headline** main label for general resistance management

information and to the crop-specific use recommendations and restrictions found in this label.

APPLICATION INFORMATION

Apply **Headline** according to the rate, timing, resistance management and adjuvant use recommendations in the **Crop-Specific Use Directions (Table A)** of this label.

Headline may be applied by ground sprayer, aerial equipment or through sprinkler irrigation systems. Refer to the **Headline** main label for specific instructions on these methods.

RESTRICTIONS AND LIMITATIONS

DO NOT use in greenhouse or transplant production systems.

DO NOT apply more than 18 fluid ounces of **Headline** per acre per season.

Refer to the EPA-approved main label for all other **Restrictions and Limitations**.



The Chemical Company

Table A - Headline® fungicide Crop-Specific Use Directions

Crop	Target Diseases	Use Rate per Application per Acre (fl oz.)	Maximum Number of Applications per Season	Maximum Rate per Acre per Season	Minimum Time from Application to Harvest (PHI)
Barley	Black point (Kernel blight or Head mold) (<i>Alternaria</i> spp., <i>Cochliobolus sativus</i> , <i>Helminthosporium sativum</i>)	6 to 9 fl. oz. per acre	2	18 fl. oz. per acre	21 days (Harvest is defined as the time when the grain is cut)
<p>Application Directions: For optimal disease control and grain quality it is important to apply Headline to protect the heads before they become infected. Headline may be reapplied 10-14 days later if weather conditions favor disease development or if extended residual disease control is desired.</p> <p>Use the higher rate and shorter interval when disease pressure is high. The higher rate can also provide longer residual protection of the heads.</p> <p>Do not apply more than 9 fl. oz. per acre in a single application.</p> <p>Resistance Management: To limit the potential for development of resistance, do not make more than two (2) applications of Headline or other QoI (Group 11) fungicides per season.</p> <p>Do not harvest barley hay within 14 days of last application.</p>					

©2006 BASF Corporation
All rights reserved

Headline is a registered trademark of BASF.

0007969-00186.20060503.NVA 2006-04-088-0161

24(c) Registrant:
BASF Corporation
Agricultural Products
26 Davis Drive
Research Triangle Park, NC 27709

