

Improved Health	Chronic Condition Composite • Use of appropriate medications for asthma • HbA1c Poor Control • Beta blocker therapy for LVSD	Children involved with child-welfare: meeting the periodicity schedule	Annual Well Exam	Use of Appropriate Medications for Asthma	Hospital readmissions 7, 30, 90, and 180 days
	Behavioral Health Composite • Clinical depression screening and follow up • Maternal Depression screening • SUD screening • Developmental screening in the first three years of life *	Transition-aged youth SUD screening, risky behavior screening	Dental	Clinical Depression Screening and Follow Up	Percent of members prescribed redundant or duplicated atypical antipsychotic medications
		Corrections Population: clients have a well-visit within 30 days of being released.		HbA1c Poor Control	Psychotropic utilization in children
		OCL population clients have a well-visit with in the last year		Controlling High Blood Pressure	Mental health engagement
				Weight Assessment and Counseling for Nutrition and physical activity for children and adolescents	Adherence to antipsychotics for individuals with schizophrenia
				BMI screening and follow up plan	Initiation and engagement of alcohol or other drug dependence treatment
				Influenza Immunizations	Penetration rates
				Maternal Depression Screening	Diabetes screening for individuals schizophrenia or bipolar disorder who are using antipsychotic medication
				Colorectal Cancer Screening	Cardiovascular monitoring for people with cardiovascular disease and schizophrenia
				Screening for fall risk	Diabetes monitoring for people with diabetes and schizophrenia
				Tobacco use screening and cessation	Follow up appointments within 7 and 30 days after hospital discharge for a mental health condition - adult and child/adolescent
				Initiation and engagement of alcohol and other drug dependence treatment	Members with physical health well-care visits
				Beta blocker therapy for LVSD	Inpatient utilization
				Diabetes: blood pressure management	ED utilization for a mental health condition
				Developmental Screening in the first three years of life *	Follow up after ED visit for mental health or alcohol or other dependence
			Anxiety screening and follow up	Antidepressant medication management acute and continuation phase	
			SUD screening	Child and Adolescent MDD: Suicide risk assessment	
			Breast Cancer screening	Adult MDD: Suicide risk assessment	
			F/U after dx from ED for MH or SUD	Adult BMI assessment and follow up	
			Cardiovascular screening for people with schizophrenia or bipolar disorder who are prescribed antipsychotics	Child and adolescent BMI assessment with nutrition and physical activity follow up	
			Adherence to mood stabilizers for individuals with bipolar 1 disorder	Controlling high blood pressure	
				Depression remission at 12 months using standard PHQ-9	
				Tobacco use screening and cessation	
				Unhealthy alcohol screening and brief counseling	
				Adolescent health screening and referral	
More Value	Total Cost of Care	Admissions for ambulatory sensitive conditions		Use of Imaging studies for low back pain	
	ED Visits for Ambulatory Sensitive Conditions				
Better Experience	CAMP5 - Coordination of Care Composite		Client Goal		Echo survey
	ECHO - Perceived Improvement Composite		Selecting/Attesting to PCMP		
	NCIAD				