

Colorado Division of Workers' Compensation

General Literature Search Strategy

Studies were identified through the electronic database of PubMed (with specified search topics), and related links from articles identified by searches. For some articles, Web of Science, a literature citation database, was used when it was desirable to find literature that cited a particular article. Relevant evidence statements from Cochrane and British Medical Journal (BMJ) Clinical Evidence were reviewed. Selected guidelines/systematic reviews were also reviewed. The reference lists from other literature and tables of content from related journals were scanned for relevant articles. Suggestions from various volunteer advisory bodies to the Division of Workers' Compensation were solicited.

Literature reviewed was in English. Literature searches were limited according to study type and human adults. Only randomized clinical trials (RCT) or meta-analyses were used for evidence statements regarding treatment. Beginning with the Traumatic Brain Injury Medical Treatment Guidelines Revision of 2012, if meta-analyses were of high enough quality, then previous RCTs that were incorporated into the selected meta-analyses may not have been individually critiqued. Selected RCTs published after Cochrane meta-analyses were evaluated as to whether they would have likely met the Cochrane inclusion criteria. If so, the Cochrane software (RevMan) was used to update the pooled effect measure and compare it with the original Cochrane report. Diagnostic accuracy studies were critiqued for diagnostic testing evidence and cohort, cross-sectional and case-control studies were critiqued for causation evidence statements. Literature which did not meet requirements for evidence statements could be referenced if it furnished useful background information or described interventions which are considered generally accepted by a consensus of health care providers. This information sometimes contributed to consensus decisions by the multi-disciplinary task force drafting the guidelines. Literature that was determined either be unrelated to the clinical issue, or which had such poor quality on initial review that it could not qualify for evidence nor provide meaningful input was not critiqued. All articles sent by the public were formally reviewed.