

## SUPPLEMENTAL LABELLING FOR SPECIAL LOCAL NEED IN COLORADO FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF COLORADO



FOR COMMERCIAL USE ONLY  
EPA REG. No. 10163-251  
SLN No. CO-010006

<b>ACTIVE INGREDIENT:</b> Hexythiazox	<b>% By Wt.</b>
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide .....	50.0%
<b>OTHER INGREDIENTS</b> .....	50.0%
	<b>TOTAL</b> 100.0%

### KEEP OUT OF REACH OF CHILDREN CAUTION

This label must be in the possession of the user at the time of the pesticide application.

Follow all other applicable direction, restriction, Worker Protection Standard requirements, and precautions on the EPA-registered label.

**Application Directions:** Apply lower rates when mite infestation levels are low and higher rates on moderate to high mite infestation levels. Make only one application per crop cycle or once per year. This product is for use in commercial plantings. **It is also intended for use in home plantings and ornamental landscaping by licensed commercial pesticide applicators.**

#### RECOMMENDATIONS

Crop	Pest	Rate	Comments
Ornamental Lawns and Turf (including residential)	Banks Grass	1-2 oz. per 100 gals. applied dilute	Apply to lawns and turf at the first signs of egg deposition. Thorough coverage is required to maximize control of pest mite species. When using high-volume dilute sprays, do not exceed the maximum per acre rate. For concentrate equipment, use the recommended rate per acre.
		<b>OR</b>	
	Mite	3-6 oz. per acre applied concentrate	
		<b>OR</b>	
		0.07-0.14 oz. per 1000 square feet	

- ◆ Do not use Plyac® or household detergents in tank-mix combinations with Hexygon.
- ◆ Do not graze or feed livestock on cover crops growing in treated areas. Do not use on crops used for feed or forage.
- ◆ Do not use water from treated areas for irrigating other crops.