CONFIDENTIAL REPORT OF SEXUALLY TRANSMITTED INFECTIONS

CDPHE DCEED-STI/HIV-REGISTRY A3 4300 Cherry Creek Dr. South Denver CO 80246-1530 (303-692-2697)

PLEASE COMPLETE ALL INFORMATION AND MAIL OR FAX TO CDPHE-STI REGISTRY - FAX 303-782-5393

LAST NAME _____ FIRST NAME _____ MIDDLE _____

STREET _____

CITY _____ ZIP _____ COUNTY _____ STATE _____ PHONE _____

BIRTHDAY _____ AGE ____ SEX ____ AMER. INDIAN ___ ASIAN/PACIFIC ___ BLACK ___ WHITE ___ HISPANIC ___ NON-HISPANIC ____

LAB PERFORMING TEST _____ PHONE (_____) _____-_____

FACILITY / CLINIC: _____ PROVIDER _____

STREET _____ CITY _____ STATE _____ ZIP _____ COUNTY _____ PHONE _____

Positive **SYPHILIS** ____ Positive **GONORRHEA** ____ Positive **CHLAMYDIA** ____ **NON-GONOCOCCAL PID** ____ **OTHER** (SPECIFY) _____
QUANTITATIVE RESULT _____

ACCESSION ID _____ **SPECIMEN TYPE** _____ **TEST TYPE** _____

SPECIMEN DATE _____ **RESULT DATE** _____

SPECIMEN SOURCE **Cervix** **Vagina** **Urine** **Urethra** **Throat** **Rectum** **Eye** **Other** (SPECIFY) _____

PREGNANT ____ **YES** ____ **NO**
Per CDC - Doxy Not Recommended in Pregnancy

PELVIC INFLAMMATORY DISEASE (PID) ____ **YES** ____ **NO**

CDC TREATMENT GUIDELINES

Please Circle ALL Treatment Administered

Partner(s) Treated **Yes** **No** **UNK** **Date Treated** _____

SYPHILIS	**GONORRHEA**	**CHLAMYDIA**	**PELVIC INFLAMMATORY DISEASE**
Primary and Secondary Syphilis Benzathine Penicillin G 2.4 million units IM in a single dose **Early Latent Syphilis** Benzathine penicillin G 2.4 million units IM in a single dose **Late Latent Syphilis or Latent Syphilis of Unknown Duration** Benzathine penicillin g 7.2 million units total, administered as 3 doses of 2.4 million units IM each at 1-week intervals. Not to exceed 10 days between doses. Missed doses are not acceptable for pregnant patients receiving therapy for late latent syphilis.	**Uncomplicated Gonococcal Infections of the Cervix, Urethra, and Rectum** Ceftriaxone 250 mg IM in a single dose OR, IF NOT AN OPTION Cefixime 400 mg orally in a single dose OR Single-dose injectible cephalosporin regimens **PLUS** Azithromycin 1g orally in a single dose OR Doxycycline 100 mg orally twice a day for 7 days **Uncomplicated Gonococcal Infections of the Pharynx** Ceftriaxone 250 mg IM in a single dose PLUS Azithromycin 1g orally in a single dose OR Doxycycline 100 mg orally twice a day for 7 days	**Uncomplicated Chlamydia** Azithromycin 1 g orally in a single dose OR Doxycycline 100 mg orally twice a day for 7 days *Alternative Regimens* Erythromycin base 500 mg orally four times a day for 7 days OR Erythromycin ethylsuccinate 800 mg orally four times a day for 7 days OR Levofloxacin 500 mg orally once daily for 7 days OR Ofloxacin 300 mg orally twice a day for 7 days	Ceftriaxone 250 mg IM in a single dose **PLUS** Doxycycline 100 mg orally twice a day for 14 days WITH or WITHOUT Metronidazole 500 mg orally twice a day for 14 days OR Cefoxitin 2 g IM in a single dose and Probenecid, 1 g orally administered concurrently in a single dose **PLUS** Doxycycline 100 mg orally twice a day for 14 days WITH or WITHOUT Metronidazole 500 mg orally twice a day for 14 days OR Ceftizoxime 2 g IM in a single dose OR / Cefotaxime 2 g IM in a single dose **PLUS** Doxycycline 100 mg orally twice a day for 14 days WITH or WITHOUT Metronidazole 500 mg orally twice a day for 14 days

CDC Treatment Guidelines - www.cdc.gov/mmwr FOR INFORMATION, REPORT FORMS, PARTNER NOTIFICATION SERVICES OR CONSULTATION PLEASE CONTACT CDPHE SEXUALLY TRANSMITTED DISEASE/AIDS CONTROL 303-692-2697 CDPHE MAY NEED TO CONTACT YOUR CLIENT REGARDING NOTIFICATION OR TREATMENT FOR A COMMUNICABLE DISEASE.