

Table 1.13A (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2010 and 2011

Drug	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
ILLCIT DRUGS¹	34,163	33,552	10,163	9,844	3,074	2,888	2,485	2,490	4,603	4,466	24,001	23,707
Marijuana and Hashish	25,887	26,186	7,198	7,127	2,080	2,052	1,811	1,821	3,307	3,253	18,689	19,059
Cocaine	4,294 ^a	3,628	1,512 ^a	1,210	387	434	402 ^a	261	723 ^a	516	2,782	2,418
Crack	866 ^a	597	469 ^a	291	158	108	140	71	170	112	397	306
Heroin	593	566	292	366	137	144	45	84	110	137	301	201
Hallucinogens	3,795	3,421	1,256	1,084	397	378	364	264	495	443	2,540	2,336
LSD	746	721	287	235	88	86	115 ^a	49	84	100	459	486
PCP	61	80	34	37	24	10	3	11	6	16	27	43
Ecstasy	2,188	2,003	751	675	207	233	216	156	328	285	1,436	1,328
Inhalants	1,163	1,041	415	390	141	114	112	119	162	157	748	651
Nonmedical Use of Psychotherapeutics ^{2,3}	14,238 ^a	12,908	5,333	4,764	1,747	1,517	1,308	1,230	2,278	2,016	8,905	8,144
Pain Relievers	10,720 ^a	9,659	4,072	3,586	1,440	1,162	1,002	1,024	1,629	1,400	6,648	6,073
OxyContin [®]	1,645	1,404	783	667	384 ^a	198	195	204	204	265	862	738
Tranquilizers	5,117	4,635	2,275	1,948	691	676	630	491	955	781	2,841	2,686
Stimulants ³	2,577	2,401	1,061	1,043	320	393	255	231	486	419	1,517	1,358
Methamphetamine ³	861	928	367	507	118	197	74	127	176	183	494	420
Sedatives	796 ^b	442	394	252	187	123	77	73	131	57	401 ^a	190
ILLCIT DRUGS OTHER THAN MARIJUANA¹	17,866 ^a	16,395	6,427 ^a	5,724	2,058	1,811	1,619	1,459	2,750	2,454	11,440	10,671

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.13B (REVISED) Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2010 and 2011

Drug	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
ILLCIT DRUGS¹	14.9	14.4	24.5	23.8	32.9	32.1	25.2	24.1	20.7	20.3	12.8	12.4
Marijuana and Hashish	11.3	11.3	17.4	17.2	22.3	22.8	18.4	17.6	14.9	14.8	9.9	10.0
Cocaine	1.9 ^b	1.6	3.7 ^a	2.9	4.1	4.8	4.1 ^a	2.5	3.3	2.3	1.5	1.3
Crack	0.4 ^a	0.3	1.1 ^a	0.7	1.7	1.2	1.4	0.7	0.8	0.5	0.2	0.2
Heroin	0.3	0.2	0.7	0.9	1.5	1.6	0.5	0.8	0.5	0.6	0.2	0.1
Hallucinogens	1.7	1.5	3.0	2.6	4.2	4.2	3.7 ^a	2.6	2.2	2.0	1.4	1.2
LSD	0.3	0.3	0.7	0.6	0.9	1.0	1.2 ^a	0.5	0.4	0.5	0.2	0.3
PCP	0.0	0.0	0.1	0.1	0.3	0.1	0.0	0.1	0.0	0.1	0.0	0.0
Ecstasy	1.0	0.9	1.8	1.6	2.2	2.6	2.2	1.5	1.5	1.3	0.8	0.7
Inhalants	0.5	0.4	1.0	0.9	1.5	1.3	1.1	1.1	0.7	0.7	0.4	0.3
Nonmedical Use of Psychotherapeutics ^{2,3}	6.2 ^b	5.5	12.9	11.5	18.7	16.8	13.3	11.9	10.3	9.1	4.7 ^a	4.3
Pain Relievers	4.7 ^b	4.2	9.8	8.7	15.4	12.9	10.2	9.9	7.3	6.4	3.5	3.2
OxyContin [®]	0.7	0.6	1.9	1.6	4.1 ^a	2.2	2.0	2.0	0.9	1.2	0.5	0.4
Tranquilizers	2.2	2.0	5.5	4.7	7.4	7.5	6.4	4.8	4.3	3.5	1.5	1.4
Stimulants ³	1.1	1.0	2.6	2.5	3.4	4.4	2.6	2.2	2.2	1.9	0.8	0.7
Methamphetamine ³	0.4	0.4	0.9	1.2	1.3	2.2	0.7	1.2	0.8	0.8	0.3	0.2
Sedatives	0.3 ^b	0.2	1.0	0.6	2.0	1.4	0.8	0.7	0.6	0.3	0.2 ^a	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	7.8 ^b	7.0	15.5 ^a	13.8	22.0	20.1	16.4	14.1	12.4	11.1	6.1 ^a	5.6

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14A (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Numbers in Thousands, 2010 and 2011

Substance	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
PAST YEAR												
Tobacco Products ¹	78,965	77,656	18,937	17,909	4,944	4,744	4,662	4,667	9,330	8,498	60,029	59,747
Cigarettes	65,139	63,839	16,722	15,851	4,573	4,369	4,105	4,131	8,044	7,351	48,417	47,988
Smokeless Tobacco	11,169	10,621	2,151 ^a	1,715	477	398	504 ^a	312	1,170	1,005	9,018	8,905
Cigars	24,179	23,040	5,246	4,648	1,376	1,263	1,248	1,152	2,622	2,233	18,933	18,392
Alcohol	161,399	163,482	29,515	28,927	6,591	6,640	7,072	7,335	15,852	14,952	131,884	134,555
PAST MONTH												
Tobacco Products ¹	67,051	65,735	16,359	15,571	4,292	4,259	4,091	3,995	7,976	7,317	50,692	50,164
Cigarettes	56,295	54,880	14,593	13,880	4,015	3,950	3,655	3,579	6,923	6,351	41,702	41,000
Smokeless Tobacco	8,309	7,726	1,492	1,180	316	242	324 ^a	187	852	751	6,817	6,546
Cigars	12,485	12,007	2,864	2,516	721	643	699	579	1,444	1,294	9,621	9,491
Pipe Tobacco	2,000	1,955	646	534	190	140	128	197	328	197	1,354	1,421
Alcohol	128,055	130,059	22,829	22,029	5,000	4,840	5,345	5,392	12,484	11,797	105,226	108,030
Binge Alcohol Use ²	56,623	56,491	11,155	10,569	2,594	2,552	2,657	2,652	5,904	5,365	45,467	45,922
Heavy Alcohol Use ²	16,490	15,486	3,523	3,314	885	867	879	822	1,760	1,625	12,967	12,171

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 1.14B (REVISED) Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Level of Mental Illness: Percentages, 2010 and 2011

Substance	Total (2010)	Total (2011)	Any Mental Illness (2010)	Any Mental Illness (2011)	Serious Mental Illness (2010)	Serious Mental Illness (2011)	Moderate Mental Illness (2010)	Moderate Mental Illness (2011)	Mild Mental Illness (2010)	Mild Mental Illness (2011)	No Mental Illness (2010)	No Mental Illness (2011)
PAST YEAR												
Tobacco Products ¹	34.4 ^a	33.4	45.7 ^a	43.3	53.0	52.6	47.2	45.2	42.0 ^a	38.6	32.0	31.2
Cigarettes	28.4	27.4	40.4	38.3	49.0	48.5	41.6	40.0	36.2	33.4	25.8	25.1
Smokeless Tobacco	4.9	4.6	5.2 ^a	4.1	5.1	4.4	5.1 ^b	3.0	5.3	4.6	4.8	4.7
Cigars	10.5 ^a	9.9	12.7 ^a	11.2	14.7	14.0	12.6	11.2	11.8	10.1	10.1	9.6
Alcohol	70.4	70.3	71.3	69.9	70.6	73.7	71.7	71.0	71.4 ^a	67.8	70.2	70.4
PAST MONTH												
Tobacco Products ¹	29.2 ^a	28.3	39.5	37.6	46.0	47.3	41.5	38.7	35.9	33.2	27.0	26.2
Cigarettes	24.6 ^a	23.6	35.2	33.5	43.0	43.8	37.0	34.6	31.2	28.8	22.2	21.4
Smokeless Tobacco	3.6	3.3	3.6	2.9	3.4	2.7	3.3 ^a	1.8	3.8	3.4	3.6	3.4
Cigars	5.4	5.2	6.9	6.1	7.7	7.1	7.1	5.6	6.5	5.9	5.1	5.0
Pipe Tobacco	0.9	0.8	1.6	1.3	2.0	1.6	1.3	1.9	1.5	0.9	0.7	0.7
Alcohol	55.9	55.9	55.1	53.2	53.6	53.7	54.2	52.2	56.2	53.5	56.0	56.5
Binge Alcohol Use ²	24.7	24.3	26.9	25.5	27.8	28.3	26.9	25.7	26.6	24.3	24.2	24.0
Heavy Alcohol Use ²	7.2 ^a	6.7	8.5	8.0	9.5	9.6	8.9	8.0	7.9	7.4	6.9	6.4

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings*).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-IV)*. Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes persons in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled *Revised Estimates of Mental Illness from the National Survey on Drug Use and Health* at <http://samhsa.gov/data/default.aspx>.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.