



Colorado Department of Health Care Policy and Financing
Preferred Drug List (PDL)
 Effective April 1, 2020

PA Forms: Available online at <https://www.colorado.gov/hcpf/pharmacy-resources>

PA Requests: Colorado Pharmacy Call Center Phone Number: 800-424-5725 | Colorado Pharmacy Call Center Fax Number: 800-424-5881

The PDL applies to Medicaid fee-for-service members. It does not apply to members enrolled in Rocky Mountain Health HMO or Denver Health Medicaid Choice.

Initiation of pharmaceutical product subject to Prior Authorization:

Please note that starting the requested drug, including a non-preferred drug, prior to a PA request being reviewed and approved, through either inpatient use, by using office "samples", or by any other means, does not necessitate Medicaid approval of the PA request.

Health First Colorado, at 25.5-5-501, requires the generic of a brand name drug be prescribed if the generic is therapeutically equivalent to the brand name drug. Exceptions to this rule are: 1) If the brand name drug is more cost effective than the generic as determined by the Department, 2) If the patient has been stabilized on a brand name drug and the prescriber believes that transition to a generic would disrupt care, and 3) If the drug is being used for treatment of mental illness, cancer, epilepsy, or human immunodeficient virus and acquired immune deficiency syndrome.

Brand Name Required = BNR, Prior Authorization = PA, AutoPA = authorization can be automated at the point of sale transaction if criteria is met
Preferred drug list applies only to prescription (RX) products, unless specified

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-preferred products will be approved for one year unless otherwise stated.)
I. Analgesics		
Therapeutic Drug Class: NON-OPIOID ANALGESIA AGENTS - Oral - Effective 7/1/2019		
No PA Required <i>Brand/generic changes effective 10/01/19</i> Duloxetine capsule (generic Cymbalta)	PA Required CYMBALTA (duloxetine) Duloxetine capsule (generic Irenka) GRALISE (gabapentin)	Non-preferred oral non-opioid analgesic agents may be approved if member meets all of the following criteria: <ul style="list-style-type: none"> Member has trialed and failed duloxetine (20mg, 30mg, or 60mg) AND has trialed and failed gabapentin OR Lyrica (Failure is defined as lack of efficacy with 8 week trial, allergy, intolerable side effects, or significant drug-drug interaction) AND Duloxetine (20mg, 30mg, or 60mg) will be approved for members with a diagnosis of fibromyalgia, neuropathic pain, or chronic musculoskeletal pain (e.g. osteoarthritis or chronic lower back pain)

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Gabapentin capsule, tablet, solution</p> <p>Pregabalin capsules</p>	<p>LYRICA (pregabalin) capsule, solution, CR tablet</p> <p>NEURONTIN (gabapentin) capsule, tablet, solution</p> <p>Pregabalin solution</p> <p>SAVELLA (milnacipran) tablet</p>	<p>through automated verification (AutoPA) upon claim submission of the corresponding ICD-10 diagnosis code related to indicated use of the medication</p> <p>Prior authorization will be required for Lyrica dosages > 600mg per day (maximum of 3 capsules daily) and gabapentin dosages > 3600mg per day.</p>
--	--	--

Therapeutic Drug Class: NON-OPIOID ANALGESIA AGENTS - Topical - Effective 7/1/2019

No PA Required	PA Required	
<p>Lidocaine Patch</p>	<p>LIDODERM Patch (lidocaine)</p> <p>ZTLIDO Patch (lidocaine)</p>	<p>Non-preferred topical products require a trial/failure with an adequate 8-week trial of gabapentin AND Lyrica AND duloxetine AND lidocaine patch. Failure is defined as lack of efficacy with an 8 week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Prior authorization will be required for Lidocaine Patch quantities exceeding 90 patches per 30 days (maximum of 3 patches daily).</p>

Therapeutic Drug Class: NON-STEROIDAL ANTI-INFLAMMATORIES (NSAIDS) - Oral - Effective 1/1/2020

No PA Required	PA Required	
<p>Celecoxib capsule</p> <p>Diclofenac potassium tablet</p> <p>Diclofenac sodium EC/DR tablet</p> <p>Ibuprofen suspension, tablet (RX)</p> <p>Indomethacin capsule, ER capsule</p> <p>Ketorolac tablet**</p> <p>Meloxicam tablet</p>	<p>ARTHROTEC (diclofenac sodium/misoprostol) tablet</p> <p>CELEBREX (celecoxib) capsule</p> <p>DAYPRO (oxaprozin) caplet</p> <p>Diclofenac sodium ER tablets</p> <p>Diclofenac sodium/misoprostol tablet</p> <p>Diffunisal tablet</p> <p>DUEXIS (ibuprofen/famotidine) tablet</p> <p>Etodolac capsule, IR and ER tablet</p>	<p>Non-preferred oral agents may be approved for members who have trialed and failed four preferred agents. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.)</p> <p>Duexis (ibuprofen/famotidine) or Vimovo (naproxen/esomeprazole) may be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> ● Trial and failure of all preferred NSAIDs at maximally tolerated doses AND ● Trial and failure of three preferred proton pump inhibitors in combination with NSAID within the last 6 months AND ● Have a documented history of gastrointestinal bleeding <p>(Failure is defined as: lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions)</p> <p>**Ketorolac tablets quantity limit: 5 days of therapy for every 30 days Tablets:20 tablets for 30 days</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Nabumetone tablet</p> <p>Naproxen EC, DR/ER, suspension, tablet (RX)</p> <p>Sulindac tablet</p>	<p>FELDENE (piroxicam) capsule</p> <p>Fenoprofen capsule, tablet</p> <p>Flurbiprofen tablet</p> <p>INDOCIN (indomethacin) susp</p> <p>Ketoprofen IR, ER capsule</p> <p>Meclofenamate capsule</p> <p>Mefenamic acid capsule</p> <p>NALFON (fenoprofen) capsule, tablet</p> <p>NAPRELAN (naproxen CR) tablet</p> <p>Naproxen sodium CR, ER, IR tablet</p> <p>Oxaprozin tablet</p> <p>Piroxicam capsule</p> <p>QMIIZ (meloxicam) ODT</p> <p>TIVORBEX (indomethacin) capsule</p> <p>Tolmetin tablet, capsule</p> <p>VIMOVO (naproxen/esomeprazole) tablet</p> <p>VIVLODEX (meloxicam) capsule</p> <p>ZIPSOR (diclofenac) capsule</p>	
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

	ZORVOLEX (diclofenac) capsule	
--	-------------------------------	--

Therapeutic Drug Class: NON-STEROIDAL ANTI-INFLAMMATORIES (NSAIDS) - Non-Oral - Effective 1/1/2020

No PA Required	PA Required	
Diclofenac 1.5% topical solution	Diclofenac 1.3% topical patch (generic Flector)	<p>Non-preferred topical agents may be approved for members who have trialed and failed one preferred agent. Failure is defined as lack of efficacy with 14 day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Sprix (ketorolac) intranasal will be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> ● Unable to tolerate, swallow or absorb oral NSAIDs OR ● Trial and failure of three preferred oral or topical NSAID agents (failure is defined as: lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) ● Quantity limit: 5-single day nasal spray bottles per 30 days <p>Flector (diclofenac) patch quantity limit: 2 patches per day</p> <p>Solaraze (diclofenac sodium) gel prior authorization criteria can be found on the Appendix P.</p>
VOLTAREN (diclofenac) 1% gel	FLECTOR (diclofenac) 1.3% topical patch	
Diclofenac sodium 1% (generic Voltaren) gel	PENNSAID (diclofenac solution) 2% Pump, 2% Solution Packet	
	SPRIX (ketorolac) nasal spray	

Opioid Utilization Policy (long-acting and short-acting opioids):

It is highly encouraged that the healthcare team utilize the Prescription Drug Monitoring Program (PDMP) to aid in ensuring safe and efficacious therapy for members using controlled substances.

Total Morphine Milligram Equivalent Policy Effective 10/1/17:

- The maximum allowable morphine milligram equivalent (MME) is 200 MME. Prescriptions for short-acting (SA) and long-acting (LA) opioids are cumulatively included in this calculation. The prescription that exceeds the cumulative MME limit of 200 MME for a member will require prior authorization and may require a provider to provider telephone consultation with the pain management physician (free of charge and provided by Health First Colorado).
- Prior authorization will be granted to allow for tapering
- Prior authorization for 1 year will be granted for diagnosis of sickle cell anemia
- Prior authorization for 1 year will be granted for admission to or diagnosis of hospice or end of life care
- Prior authorization for 1 year will be granted for pain associated with cancer

MME calculation is conducted using conversion factors from the following website: <http://agencymeddirectors.wa.gov/Calculator/DoseCalculator.htm>

Only one long-acting opioid agent (including different strengths) and one short-acting opioid agent (including different strengths) will be considered for a prior authorization.

Medicaid provides guidance on the treatment of pain, including tapering, on our webpage under the heading Pain Management Resources and Opioid Use at: <https://www.colorado.gov/pacific/hcpf/pain-management-resources-and-opioid-use>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Opioid Naïve Policy Effective 8/1/17 (Update effective 11/27/19 in Italics):

Members who have not filled a prescription for an opioid within the past 180 days will be identified as “opioid treatment naïve” and have the following limitations placed on the initial prescription(s):

- The prescription is limited to short-acting opioid agents *or Butrans (buprenorphine) 5mcg patch*. Use of other long-acting opioid agents will require prior authorization approval for members identified as opioid treatment naïve.
- The days supply of the first, second, and third prescription for an opioid will be limited to 7 days, the quantity will be limited to 8 dosage forms per day (tablets, capsules), maximum #56 tablets/capsules for a 7 day supply
- The fourth prescription for an opioid will require prior authorization, filling further opioid prescriptions may require a clinical pharmacist review or provider to provider telephone consultation with a pain management physician (free of charge and provided by Health First Colorado).
- If a member has had an opioid prescription filled within the past 180 days, then this policy would not apply to that member and other opioid policies would apply as applicable.

Dental Prescriptions Opioid Policy Effective 11/15/18 (implemented in the claims system 01/07/19):

Members who receive an opioid prescribed by a dental provider will be subject to day supply limits and quantity per day limits for short acting opioids.

- The prescription is limited to short-acting opioid agents only. Use of long-acting opioid agents and short acting fentanyl agents will require prior authorization approval for members’ prescriptions written by a dental provider.
- The days supply of the first, second, and third prescription for an opioid will be limited to 4 days, the quantity will be limited to 6 dosage forms per day (tablets, capsules), maximum #24 tablets/capsules for a 4 day supply
- The fourth prescription for an opioid will require prior authorization. A prior authorization for the fourth fill may be approved for up to a 7 day supply and the quantity will be limited to 8 dosage forms per day (#56 tablets/capsules) for members with any of the following diagnoses/undergoing any of the following procedures:
 - Traumatic oro-facial tissue injury with major mandibular/maxillary surgical procedures
 - Severe cellulitis of facial planes
 - Severely impacted teeth with facial space infection necessitating surgical management
- Other potential exemptions that exceed the first 3 fill limits (day supply and quantity) may be evaluated with a provider to provider telephone consult with a pain management specialist (free of charge and provided by Health First Colorado)

If a member has had an opioid prescription prescribed by a non-dental provider, then this policy would not apply to that member and other opioid policies would apply as applicable. Dental prescriptions do not impact the opioid treatment naïve policy, but the prescriptions will be counted towards the Morphine Milligram Equivalent (MME) daily dose.

Opioid and Benzodiazepine Combination Effective 9/15/19:

Prior authorization will be required for members receiving long-term therapy with an opioid medication who are newly started on a benzodiazepine medication OR for members receiving long-term therapy with a benzodiazepine medication who are newly started on an opioid medication. Prior authorization may be approved if meeting the following:

- The member discontinued or is no longer taking either the opioid or benzodiazepine medication and will not be using these in combination **OR**
- The member will not be taking the prescribed opioid and benzodiazepine medications at the same time based on prescribed dosing interval (such as prn administration) for the regimen **AND** the prescriber attests that the member has received appropriate counseling* regarding the risks associated with combining opioid and

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

benzodiazepine medications including increased risk for sedation, respiratory depression, overdose, and overdose-related death and counseling regarding the FDA Boxed Warning for combining these medications **OR**

- The prescriber has evaluated the regimen and attests that it is appropriate for the member to continue use of the concomitant opioid and benzodiazepine medication regimen as prescribed **AND** the prescriber attests that the member has received appropriate counseling* regarding the risks associated with combining opioid and benzodiazepine medications including increased risk for sedation, respiratory depression, overdose, and overdose-related death and counseling regarding the FDA Boxed Warning for combining these medications **OR**
- Prior authorization may be approved for members receiving palliative or hospice care **OR**
- For benzodiazepine prior authorizations, approval may be granted if the benzodiazepine is being prescribed for seizure disorder or convulsions.

**If counseling has not been provided, the prescriber attests that a reasonable effort will be made to contact the member or the member's pharmacy to ensure that counseling is provided.*

Opioid and Quetiapine Combination Effective 9/15/19:
Pharmacy claims for members receiving opioid and quetiapine medications in combination will require entry of point-of-sale DUR service codes (Reason for Service, Professional Service, Result of Service) for override of drug-drug interaction (DD) related to risk of increased sedation from concomitant use of this drug combination.

Therapeutic Drug Class: OPIOIDS, Short Acting - Effective 7/1/2019		
No PA Required* (if criteria is met)	PA Required	*Tramadol and tramadol-containing products will require prior authorization approval to verify that the following criteria are met:
Hydrocodone/apap tablet	Acetaminophen / codeine elixir, tablets**	<ul style="list-style-type: none"> • Member is \geq 12 years of age AND • If member is less than 18 years of age, tramadol is NOT being prescribed for post-surgical pain following tonsil or adenoid procedure AND • If member is between 12 and 18 years of age, member is not obese (BMI greater than 30kg/m²), does not have obstructive sleep apnea or severe lung disease AND • Non-preferred tramadol products will require trial/failure of generic tramadol tablet AND generic tramadol/APAP tablet. Failure is defined as allergy‡, lack of efficacy, intolerable side effects, or significant drug-drug interaction. <p>Rybix® ODT (tramadol hydrochloride) will be approved without trial/failure of three preferred agents for members who meet the tramadol products criteria above AND who are unable to swallow oral tablets or absorb oral medications.</p> <p>**Codeine and codeine-containing products will receive prior authorization approval for members meeting the following criteria:</p> <ul style="list-style-type: none"> • Member is \geq 12 years of age AND • If member is less than 18 years of age, codeine is NOT being prescribed for post-surgical pain following tonsil or adenoid procedure AND
Hydrocodone/apap solution	Butalbital / caffeine / acetaminophen w/ codeine**	
Hydrocodone/ibuprofen	Butalbital compound w/ codeine**	
Hydromorphone tablet	Butorphanol tartrate (nasal)	
Morphine IR tablet	Carisoprodol compound / codeine**	
Morphine soln	Codeine (all forms)**	
Oxycodone tablet	DILAUDID liquid	
Oxycodone Soln	DVORAH (acetaminophen / caffeine / dihydrocodeine)	
Oxycodone/apap tablet		

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Tramadol 50mg*</p> <p>Tramadol/apap tablet*</p>	<p>Fiorinal/codeine**</p> <p>Fioricet / codeine**</p> <p>Hydromorphone liquid</p> <p>IBUDONE (hydrocodone/ibuprofen)</p> <p>LORTAB (hydrocodone/apap)</p> <p>Levorphanol</p> <p>Meperidine solution, tablet</p> <p>Morphine concentrated solution</p> <p>NORCO (hydrocodone/apap)</p> <p>NUCYNTA*** (tapentadol)</p> <p>OPANA (oxymorphone)</p> <p>OXAYDO (oxycodone)</p> <p>Oxycodone / aspirin</p> <p>Oxycodone / acetaminophen solution</p> <p>Oxycodone / ibuprofen</p> <p>Oxycodone capsule, syringe, conc solution</p> <p>Oxymorphone</p> <p>Pentazocine / naloxone</p> <p>PERCOCET (oxycodone/apap)</p>	<ul style="list-style-type: none"> ● If member is between 12 and 18 years of age, member is not obese (BMI greater than 30kg/m2), does not have obstructive sleep apnea or severe lung disease ● Member is not pregnant or breastfeeding AND ● Renal function is not impaired (GFR > 50 ml/min) AND ● Member is not receiving strong inhibitors of CYP3A4 (e.g. erythromycin, clarithromycin, telithromycin, itraconazole, ketoconazole, posaconazole, fluconazole [≥200mg daily], voriconazole, delavirdine, and milk thistle) AND ● Member meets one of the following: <ul style="list-style-type: none"> ○ Member has trialed codeine or codeine-containing products in the past no history of allergy or adverse drug reaction to codeine ○ Member has not trialed codeine or codeine-containing products in the past and the prescriber acknowledges reading the following statement: “Approximately 1-2% of the population metabolizes codeine in a manner that exposes them to a much higher potential for toxicity. Another notable proportion of the population may not clinically respond to codeine. We ask that you please have close follow-up with members newly starting codeine and codeine-containing products to monitor for safety and efficacy.” <p><u>Maximum Doses:</u> **Tramadol maximum dose is 400mg/day **Codeine maximum dose is 360mg/day</p> <p>***Nucynta® IR (tapentadol) may be approved for members who meet the following criteria:</p> <ul style="list-style-type: none"> ● Member has history of trial/failure of 7-days utilization of preferred product(s) in the last 21 days OR ● If member does not meet the above criteria, prior authorization approval for Nucynta IR will require trial and failure of three preferred agents. Failure is defined as lack of efficacy, intolerable side effects, significant drug-drug interaction, allergy‡, or significant adverse drug reaction. ● Nucynta IR will have a maximum daily quantity of 6 tablets (180 tabs per 30 days). <p>Prior authorization for all other non-preferred short-acting opioid products will be approved if the member has trialed/failed three preferred products. Failure is defined as allergy‡, or significant adverse drug reaction.</p> <p>‡Allergy: hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema</p> <p><u>Quantity Limits:</u> Short-acting opioids will be limited to a total of 120 tablets per 30 days (4/day) per member for members who are not included in the opioid treatment naive policy. Exceptions will be</p>
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	Roxicodone tablet Tramadol 100mg TYLENOL w/codeine ULTRACET* (tramadol/apap) ULTRAM* (tramadol) ZAMICET (hydrocodone/apap)	made for members with a diagnosis of a terminal illness (hospice or palliative care) or sickle cell anemia. For members who are receiving more than 120 tablets currently and who do not have a qualifying exemption diagnosis, a 6-month prior authorization can be granted via the prior authorization process for providers to taper members. Please note that if more than one agent is used, the combined total utilization may not exceed 120 units in 30 days. There may be allowed certain exceptions to this limit for acute situations (for example: post-operative surgery, fractures, shingles, car accident). Butorphanol intranasal maximum quantity is 10ml per 30 days (four 2.5ml 10mg/ml package units per 30 days).
--	---	---

Therapeutic Drug Class: FENTANYL PREPARATIONS (buccal, intranasal, transmucosal, sublingual) - Effective 7/1/2019

	PA Required	
	Abstral (fentanyl citrate) Actiq (fentanyl citrate) Fentanyl citrate Fentora (fentanyl citrate) Lazanda (fentanyl citrate)	Fentanyl buccal, intranasal, transmucosal, and sublingual products: Prior authorization approval will be granted for members experiencing breakthrough cancer pain and those that have already received and are tolerant to opioid drugs for the cancer pain AND are currently being treated with a long-acting opioid drug. The prior authorization may be granted for up to 4 doses per day. For patients in hospice or palliative care, prior authorization will be automatically granted regardless of the number of doses prescribed. Ionsys transdermal system requires administration in the hospital setting and is not covered under the pharmacy benefit

Therapeutic Drug Class: OPIOIDS, Long Acting - Effective 7/1/2019

No PA Required	PA Required	
BUTRANS (buprenorphine) patch ^{BNR} EMBEDA (morphine/naltrexone) Fentanyl patches 12mcg, 25mcg, 50mcg, 75mcg, 100mcg	*NUCYNTA ER (tapentadol ER) BELBUCA (buprenorphine) buccal film Buprenorphine patch CONZIP (tramadol ER) DOLOPHINE (methadone)	* Nucynta ER will be approved for members who have trialed and failed‡ treatment with TWO preferred agents in the last 6 months. <u>Non-Preferred Agents:</u> All non-preferred abuse-deterrent formulations (OxyContin®, Xtampza® ER, Hysingla® ER, etc) will require trial and failure‡ of three preferred agents within the past year. All other non-preferred agents may be approved for members who have trialed and failed‡ three preferred products within the past year.

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Morphine ER (generic MS Contin)</p> <p>Tramadol ER (generic Ultram ER)</p>	<p>DURAGESIC (fentanyl) patch</p> <p>EXALGO (hydromorphone ER)</p> <p>Fentanyl patches 37mcg, 62mcg, 87mcg</p> <p>Hydromorphone ER</p> <p>HYSINGLA (hydrocodone ER)</p> <p>KADIAN (morphine ER capsules) brand and generic</p> <p>Methadone (all forms)</p> <p>MS CONTIN (morphine ER)</p> <p>MORPHABOND (morphine ER)</p> <p>OXYCONTIN (oxycodone ER)</p> <p>Tramadol ER (generic Ryzolt/ Conzip)</p> <p>VANTRELA ER (hydrocodone bitartrate)</p> <p>XARTEMIS XR (oxycodone/acetaminophen)</p> <p>XTAMPZA ER (oxycodone ER)</p> <p>ZOHYDRO ER (hydrocodone ER)</p>	<p>‡Failure is defined as lack of efficacy with 14 day trial due to allergy (hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema), intolerable side effects, or significant drug-drug interaction.</p> <p><u>Methadone Continuation:</u> Members who have been receiving methadone for pain indications do not have to meet non-preferred criteria. All new starts for methadone will require prior authorization under the non-preferred criteria listed above.</p> <p><i>If a prescriber would like to discuss strategies for tapering off methadone or transitioning to other pain management therapies for a Health First Colorado member, consultation with the Health First Colorado pain management physician is available free of charge by contacting the pharmacy call center helpdesk and requesting an opioid prescriber consult.</i></p> <p><u>Reauthorization:</u> Reauthorization for a non-preferred agent may be approved if the following criteria are met:</p> <ul style="list-style-type: none"> ● Provider attests to continued benefit outweighing risk of opioid medication use AND ● Member met original prior authorization criteria for this drug class at time of original authorization <p><u>Quantity/Dosing Limits:</u></p> <ul style="list-style-type: none"> ● OXYCONTIN®, OPANA ER®, NUCYNTA ER®, and ZOHYDRO ER® will only be approved for twice daily dosing. ● HYSINGLA ER® will only be approved for once daily dosing. ● Fentanyl patches will require a PA for doses of more than 15 patches/30 days (taking one strength) or 30 patches for 30 days (taking two strengths). For fentanyl patch strengths of 37mcg/hr, 62mcg/hr, and 87mcg/hr. Member must trial and fail two preferred strengths of separate patches summing desired dose (i.e. 12mcg/hr + 50mcg/hr =62mcg/hr)
---	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

II. Anti-Infectives

Therapeutic Drug Class: ANTI-HERPETIC AGENTS - Oral -Effective 1/1/2020

No PA Required	PA Required													
<p>Acyclovir tablet, capsule</p> <p>Acyclovir suspension <i>(members under 5 years or with a feeding tube)</i></p> <p>Valacyclovir tablet</p>	<p>Famciclovir tablet</p> <p>SITAVIG (acyclovir) buccal tablet</p> <p>VALTREX (valacyclovir) tablet</p> <p>ZOVIRAX (acyclovir) capsule, tablet</p>	<p>Non-preferred products may be approved for members who have failed an adequate trial with oral acyclovir AND valacyclovir. Failure is defined as lack of efficacy with 14 day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Sitavig (acyclovir) buccal tablet may be approved for diagnosis of recurrent herpes labialis (cold sores) if member meets non-preferred criteria listed above AND has failed trial with oral acyclovir suspension. Failure is defined as lack of efficacy with 14 day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>For members with a diagnosis of Bell's palsy, valacyclovir 1000 mg three times daily will be approved for 7 days if member presents with severe facial palsy.</p> <p>Acyclovir suspension may be approved for:</p> <ul style="list-style-type: none"> • Members under 5 years of age OR • Members with a feeding tube OR • Members meeting non-preferred criteria listed above. <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th colspan="3" style="text-align: center;">Maximum Dose Table</th> </tr> <tr> <th></th> <th style="text-align: center;">Adult</th> <th style="text-align: center;">Pediatric</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">Acyclovir</td> <td style="text-align: center;">4000 mg daily</td> <td style="text-align: center;">1200 mg daily</td> </tr> <tr> <td style="text-align: center;">Valacyclovir</td> <td style="text-align: center;">4000 mg daily</td> <td style="text-align: center;">Age 2-11 years: 3000mg daily Age ≥ 12 years: 4000mg daily</td> </tr> </tbody> </table>	Maximum Dose Table				Adult	Pediatric	Acyclovir	4000 mg daily	1200 mg daily	Valacyclovir	4000 mg daily	Age 2-11 years: 3000mg daily Age ≥ 12 years: 4000mg daily
Maximum Dose Table														
	Adult	Pediatric												
Acyclovir	4000 mg daily	1200 mg daily												
Valacyclovir	4000 mg daily	Age 2-11 years: 3000mg daily Age ≥ 12 years: 4000mg daily												

Therapeutic Drug Class: ANTI-HERPETIC AGENTS- Topical - Effective 1/1/2020

No PA Required	PA Required	
<p>DENAVIR (penciclovir) cream</p> <p>ZOVIRAX^{BNR} (acyclovir) cream</p>	<p>Acyclovir cream</p> <p>Acyclovir ointment</p>	<p>Generic Acyclovir ointment/cream will be approved for members who have failed an adequate trial with Zovirax ointment/cream (diagnosis, dose and duration) as deemed by approved compendium. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction)</p> <p>Xerese (acyclovir/hydrocortisone) prior authorization will be approved for members that meet the following criteria:</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

ZOVIRAX ^{BNR} (acyclovir) ointment	XERESE (acyclovir/hydrocortisone) cream	<ul style="list-style-type: none"> • Documented diagnosis of recurrent herpes labialis AND • Member is immunocompetent AND • Member has failed treatment of at least 10 days with acyclovir (Failure will be defined as significant drug-drug interaction, lack of efficacy, contraindication to or intolerable side effects) AND • Member has failed treatment of at least one day with famciclovir 1500 mg OR valacyclovir 2 GM twice daily (Failure is defined as significant drug-drug interaction, lack of efficacy, contraindication to or intolerable side effects)
---	---	--

Therapeutic Drug Class: TETRACYCLINES - Effective 7/1/2019

No PA Required	PA Required	
Doxycycline hyclate capsules	Demeclocycline	<p>Prior authorization for non-preferred tetracycline agents will be approved if member has trialed/failed a preferred doxycycline agent AND preferred minocycline capsule. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction</p> <p>Prior authorization for liquid oral tetracycline formulations will be approved if member has difficulty swallowing and cannot take solid oral dosage forms.</p>
Doxycycline hyclate tablets	DORYX (doxycycline)	
Doxycycline monohydrate 50mg, 100mg, capsule	Doxycycline hyclate tablet DR	
Doxycycline monohydrate tablets	Doxycycline monohydrate 40mg, 75mg, 150mg, capsule	
Minocycline capsules	Doxycycline monohydrate Suspension	
	Minocycline ER	
	Minocycline tablets	
	MINOLIRA (minocycline)	
	MORGIDOX (doxycycline)	
	NUZYRA (Omadacycline)	
	SOLODYN (minocycline)	
	Tetracycline	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	VIBRAMYCIN (doxycycline) XIMINO (minocycline)	
--	--	--

Therapeutic Drug Class: FLUOROQUINOLONES -Oral -Effective 1/1/2020

No PA Required	PA Required	
CIPRO (ciprofloxacin) oral suspension (<5 years old)	AVELOX (moxifloxacin) tablet	<p>Non-preferred products will be approved for members who have failed an adequate trial (7 days) with at least one preferred product. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.)</p> <p>CIPRO/ciprofloxacin suspension approved for members < 5 years of age without PA</p> <p>For members ≥ 5 years of age, CIPRO/ciprofloxacin suspension will only be approved for those members who cannot swallow a whole or crushed tablet</p> <p>Levofloxacin solution will be approved for members who require administration via feeding tube OR who have failed an adequate trial (7 days) of ciprofloxacin suspension. (Failure is defined as: lack of efficacy, presence of feeding tube, allergy, intolerable side effects, or significant drug-drug interaction.)</p>
Ciprofloxacin oral suspension (<5 years old)	BAXDELA (delafloxacin) tablet	
Ciprofloxacin tablet	CIPRO (ciprofloxacin) tablet	
Levofloxacin tablet	CIPRO XR (ciprofloxacin ER) tablet	
	Ciprofloxacin oral suspension (>5 years old), ER tablet	
	LEVAQUIN (levofloxacin) tablet	
	Levofloxacin oral solution	
	Moxifloxacin tablet	
	Ofloxacin tablet	

Therapeutic Drug Class: HEPATITIS C VIRUS TREATMENTS - Effective 1/1/2020

Direct Acting Antivirals (DAAs)

PA Required for all agents in this class		Preferred Hepatitis C Virus Treatment Regimens	
EPCLUSA ^{BNR} (sofosbuvir/velpatasvir)	Sofosbuvir/ledipasvir	Harvoni (ledipasvir/sofosbuvir)	Harvoni will be approved for members 3 years and older with chronic HCV infection; GT 1, 4-6; who are NC, have CC, or in combination with ribavirin in adults with DC; and meet the below applicable criteria
	Sofosbuvir/velpatasvir	Mavyret (glecapravir/pibrentasvir)	Mavyret will be approved for members 12 years and older or weighing at least 45 kg with chronic HCV infection, GT 1-6 who are NC or have CC (Child-Pugh A), and meet the below applicable criteria
HARVONI ^{BNR} (sofosbuvir/ledipasvir)	SOVALDI (sofosbuvir)		

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>MAVYRET (glecaprevir/pibrentasvir)</p>	<p>VOSEVI (sofosbuvir/velpatasvir/voxilaprevir)</p> <p>ZEPATIER (elbasvir/grazoprevir)</p>	<table border="1" data-bbox="934 277 2020 375"> <tr> <td data-bbox="934 277 1213 375"> <p>Epclusa (sofosbuvir/velpatasvir)</p> </td> <td data-bbox="1213 277 2020 375"> <p>Epclusa will be approved for adult members with chronic HCV infection, GT 1-6, who are NC, have CC, or in combination with ribavirin in DC; and meet the below applicable criteria</p> </td> </tr> </table> <p><i>(GT-Genotype, NC-Non-Cirrhotic, CC-Compensated Cirrhosis, DC-Decompensated Cirrhosis)</i></p> <p>All preferred agents will be granted prior authorization if the following criteria are met:</p> <ul style="list-style-type: none"> ● Physician attests to provide one HCV RNA test result from 12-24 weeks post-treatment showing SVR, AND ● Member must have received, or be in the process of receiving, full courses of both Hepatitis A and Hepatitis B vaccinations, or have immunity; AND ● Members must have genotyping results within 1 year before anticipated therapy start date; AND ● If member is abusing/misusing alcohol or controlled substances, member must be receiving or be enrolled in counseling or a substance use treatment program for at least 1 month prior to starting treatment; AND ● Agent must be prescribed by an infectious disease specialist, gastroenterologist, or hepatologist OR prescribed by any primary care provider in consultation with an infectious disease specialist, gastroenterologist or hepatologist; OR for treatment naïve members without cirrhosis, prescribed by any primary care who has completed the hepatitis C (HCV) ECHO series (four, 1-hour trainings); AND ● Physician attests to the member’s readiness for adherence; AND <ul style="list-style-type: none"> ○ Prescribers may utilize assessment tools to evaluate readiness of the patient for treatment, some examples are available at: http://www.integration.samhsa.gov/clinical-practice/screening-tools#drugs or Psychosocial Readiness Evaluation and Preparation for Hepatitis C Treatment (PREP-C) is available at: https://prepc.org/ ● Physician attests to member having Chronic HCV infection (Presence of HCV RNA viral load for ≥ 6 months to confirm infection is not acute or evidence that the infection has spontaneously resolved) AND ● For women of childbearing potential, serum pregnancy testing is conducted within 30 days of expected direct-acting antiviral start date AND ● The provider must provide the following laboratory tests within 6 months of initiating therapy: <ul style="list-style-type: none"> ○ Complete Blood Count (CBC) ○ Hepatic Function Panel (i.e. albumin, total and direct bilirubin, alanine aminotransferase (ALT), aspartate aminotransferase (AST), and alkaline phosphatase levels) ○ Calculated glomerular filtration rate (GFR) ○ If cirrhosis is present, calculation of the Child-Turcotte-Pugh (CTP) Score ○ Transplant status as applicable (pre-, post-, N/A) 	<p>Epclusa (sofosbuvir/velpatasvir)</p>	<p>Epclusa will be approved for adult members with chronic HCV infection, GT 1-6, who are NC, have CC, or in combination with ribavirin in DC; and meet the below applicable criteria</p>
<p>Epclusa (sofosbuvir/velpatasvir)</p>	<p>Epclusa will be approved for adult members with chronic HCV infection, GT 1-6, who are NC, have CC, or in combination with ribavirin in DC; and meet the below applicable criteria</p>			

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<p>For ribavirin-containing regimens only:</p> <ul style="list-style-type: none"> ● Member is not a pregnant female or a male with a pregnant female partner AND ● Women of childbearing potential and their male partners must attest that they will use two forms of effective (non-hormonal) contraception during treatment AND ● Member does not meet any of the following ineligibility criteria for use of ribavirin: <ul style="list-style-type: none"> ● Pregnant women and men whose female partners are pregnant ● Known hypersensitivity to ribavirin ● Autoimmune hepatitis ● Hemoglobinopathies ● Creatinine Clearance < 50mL/min ● Co-administered with didanosine <p>Non-Preferred Agents: All non-preferred agents or treatment regimens will be granted prior authorization if the criteria for preferred agents above is satisfied PLUS documentation is provided indicating an acceptable rationale for not prescribing a preferred treatment regimen. (Acceptable rationale may include: patient-specific medical contraindications to a preferred treatment, member has initiated treatment on a non-preferred drug and needs to complete therapy.)</p> <p>Re-treatment: All requests for HCV re-treatment for members who have failed therapy with a DAA will be reviewed on a case-by-case basis.</p> <p>Additional information will be requested for retreatment requests including, but not limited to:</p> <ul style="list-style-type: none"> ● Previous regimen medications and dates treated ● Genotype of previous HCV infection ● Any information regarding adherence to previously trialed regimen(s) and current chronic medications ● Adverse effects experienced from previous treatment regimen ● Concomitant therapies during previous treatment regimen <p>For regimens ≥ 12 weeks in duration:</p> <ul style="list-style-type: none"> ● Physician attests that if the week 4 HCV RNA is detectable (>25 copies) while on therapy, HCV RNA will be reassessed in 2 weeks. If the repeated HCV RNA level has not decreased (i.e. >1
--	--	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

		<p>log10 IU/ml from nadir) all treatment will be discontinued unless documentation is provided which supports continuation of therapy; AND</p> <ul style="list-style-type: none"> • All approvals will initially be for an 8-week time period, with further approvals dependent on the submission of HCV RNA levels at treatment times of 4 weeks, 12 weeks, and 20 weeks as applicable to justify continuing drug therapy; AND • Refills should be reauthorized in order to continue the appropriate treatment plan. The member MUST receive refills within one week of completing the previous fill. Please allow ample time for reauthorization after HCV RNA levels are submitted. <p>Grandfathering: Members currently receiving treatment with a non-preferred agent will receive approval to finish their treatment regimen, provided required documentation is sent via normal PAR process.</p> <p>Hepatitis C Treatment requests must be submitted via the Hepatitis C specific PAR form which can be accessed on the Pharmacy Resources page at: https://www.colorado.gov/hcpf/pharmacy-resources</p>
--	--	--

Ribavirin Products

No PA Required	PA Required	
Ribavirin capsule	MODERIBA (ribavirin)	<p>Non-preferred ribavirin products require prior authorizations which will be evaluated on a case-by-case basis.</p> <p>Members currently receiving non-preferred ribavirin product will receive approval to continue that product for the duration of their HCV treatment regimen.</p>
Ribavirin tablet	REBETOL (ribavirin) solution	
	RIBASPHERE (ribavirin)	
	Ribavirin solution	

III. Cardiovascular

Therapeutic Drug Class: ANGIOTENSIN MODIFIERS - <i>Effective 7/1/2019</i>
--

Angiotensin-converting enzyme inhibitors (ACE Inh)

No PA Required	PA Required	
Benazepril tablet	Captopril	Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products will be approved for members who have trialed

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Enalapril tablet	EPANED powder/solution* (enalapril)	<p>and failed treatment with three preferred products in the last 12 months (Failure is defined as lack of efficacy with a 4 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>*Epaned® (enalapril) powder and solution will be approved without trial/failure of three preferred agents for members under the age of 5 years who cannot swallow a whole or crushed tablet.</p>
Fosinopril tablet	QBRELIS solution (lisinopril)	
Lisinopril tablet	Moexipril	
Quinapril tablet	Perindopril	
Ramipril tablet	Trandolapril	

ACE Inh Combinations		
-----------------------------	--	--

No PA Required	PA Required	
Enalapril HCTZ	Benazepril HCTZ	<p>Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products will be approved for members who have trialed and failed treatment with three preferred products in the last 12 months (Failure is defined as lack of efficacy with a 4 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p>
Lisinopril HCTZ	Captopril HCTZ	
	Fosinopril HCTZ	
	Moexipril HCTZ	
	Quinapril HCTZ	
	ZESTORETIC (lisinopril HCTZ)	

Angiotensin II receptor blockers (ARBs)		
--	--	--

No PA Required	PA Required	
BENICAR (olmesartan)	ATACAND (candesartan)	<p>Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products will be approved for members who have trialed and failed treatment with three preferred products in the last 12 months (Failure is defined as lack of efficacy with a 4 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p>
Irbesartan	AVAPRO (irbesartan)	
Losartan	Candesartan	
Olmesartan	COZAAR (losartan)	
Telmisartan	DIOVAN (valsartan)	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

Valsartan	EDARBI (azilsartan) Eprosartan MICARDIS (telmisartan) TEVETEN (eprosartan)	
-----------	---	--

ARB Combinations		
-------------------------	--	--

No PA Required	PA Required	
Irbesartan/HCTZ Losartan/HCTZ Olmesartan/HCTZ Valsartan/HCTZ	Amlodipine/olmesartan Amlodipine/olmesartan/HCTZ Amlodipine/valsartan Amlodipine/valsartan/HCTZ ATACAND HCT (candesartan/HCTZ) AVALIDE (irbesartan/HCTZ) AVALIDE (irbesartan/HCTZ) AZOR (amlodipine/olmesartan) BENICAR HCT (olmesartan/HCTZ) BYVALSON (nebivolol/valsartan) Candesartan/HCTZ DIOVAN HCT (valsartan/HCTZ) EDARBYCLOR (azilsartan/chlorthalidone)	Non-preferred ACE inhibitors, ACE inhibitor combinations, ARBs, ARB combinations, renin inhibitors, and renin inhibitor combination products will be approved for members who have trialed and failed treatment with three preferred products in the last 12 months (Failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects, or significant drug-drug interaction).

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

	Eprosartan/HCTZ EXFORGE (amlodipine/valsartan) EXFORGE HCT (amlodipine/valsartan/HCTZ) HYZAAR HCT (losartan/HCTZ) MICARDIS-HCT (telmisartan/HCTZ) Telmisartan/HCTZ Telmisartan/amlodipine Telmisartan/HCTZ TRIBENZOR (amlodipine/olmesartan/HCTZ) TWYNSTA (telmisartan/amlodipine)	
--	---	--

Renin Inhibitors & Renin Inhibitor Combinations		
--	--	--

	<p style="text-align: center;">PA Required</p> Aliskiren TEKTURNA (aliskiren) TEKTURNA HCT (aliskiren/HCTZ)	<p>Non-preferred renin inhibitors and renin inhibitor combination products will be approved for members who have failed treatment with three preferred products in the last 12 months (Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>Renin inhibitors and combinations will not be approved in patients with diabetes. Renin inhibitors are contraindicated when used in combination with an ACE-inhibitor, ACE-inhibitor combination, ARB, or ARB-combination.</p>
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

Therapeutic Drug Class: **PULMONARY ARTERIAL HYPERTENSION THERAPIES** - *Effective 1/1/2020*

Phosphodiesterase Inhibitors

<p>*Must meet eligibility criteria</p> <p>*Sildenafil (generic Revatio) 20 mg tablet</p> <p>*Tadalafil 20mg</p>	<p>PA Required</p> <p>ADCIRCA (tadalafil)</p> <p>ALYQ (tadalafil) 20mg</p> <p>REVATIO (sildenafil) 20mg tablet, suspension</p> <p>Sildenafil (generic Revatio) oral suspension</p>	<p>*Eligibility Criteria for all agents in the class</p> <p>Approval will be granted for a diagnosis of pulmonary hypertension.</p> <p>Non-preferred products may be approved for members who have failed treatment with preferred sildenafil AND preferred tadalafil. Failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Revatio (sildenafil) suspension will approved for members who are unable to take/swallow tablets</p> <p>Grandfathering: Members who have been previously stabilized on a Non-preferred product can receive approval to continue on the medication.</p>
--	---	--

Endothelin Antagonists

<p>*Must meet eligibility criteria</p> <p>*LETAIRIS^{BNR} (ambrisentan) tablet</p> <p>*TRACLEER 62.5mg, 125mg (bosentan) tablet ^{BNR}</p>	<p>PA Required</p> <p>Ambrisentan (generic Letairis) tablet</p> <p>Bosentan (generic Tracleer) 62.5mg, 125mg tablet</p> <p>OPSUMIT (macitentan)</p> <p>TRACLEER (bosentan) 32mg tablet for suspension</p>	<p>*Eligibility Criteria for all agents in the class</p> <p>Approval will be granted for a diagnosis of pulmonary hypertension. Member and prescriber should be enrolled in applicable REMS program for prescribed medication.</p> <p>Non-preferred agents will be approved for members who have trialed and failed two preferred agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Grandfathering: Members who have been previously stabilized on a Non-preferred product can receive approval to continue on the medication.</p>
---	--	--

Prostanoids

<p>*Must meet eligibility criteria</p> <p>*Epoprostenol (generic Flolan) vial</p> <p>*ORENITRAM (treprostini) ER tablet</p>	<p>PA Required</p> <p>FLOLAN (epoprostenol) vial</p> <p>REMOTULIN (treprostini) vial</p> <p>Treprostini (generic Remodulin) vial</p>	<p>*Eligibility Criteria for all agents in the class</p> <p>Approval will be granted for a diagnosis of pulmonary hypertension.</p> <p>Non-preferred products will be approved for members who have failed treatment with a Preferred Product. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, contraindication to IV therapy or significant drug-drug interaction)</p>
--	---	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

*VENTAVIS (iloprost) inhalation solution	TYVASO (treprostinil) inhalation solution UPTRAVI (selexipag) tablet VELETRI (epoprostenol) vial	Grandfathering: Members who have been previously stabilized on a non-preferred product can receive approval to continue on the medication.
--	--	---

Guanylate Cyclase (sGC) Stimulator

	<p style="text-align: center;">PA Required</p> <p>ADEMPAS (riociguat) tablet</p>	<p>Adempas will be approved for patients who meet the following criteria:</p> <ul style="list-style-type: none"> ● Patient is not a pregnant female and is able to receive monthly pregnancy tests while taking Adempas and one month after stopping therapy. AND ● Women of childbearing potential and their male partners must use one of the following contraceptive methods during treatment and one month after stopping treatment (e.g. IUD, contraceptive implants, tubal sterilization, a hormone method with a barrier method, two barrier methods, vasectomy with a hormone method, or vasectomy with a barrier method). AND ● Patient is not receiving dialysis or has severe renal failure (e.g. Crcl < 15 ml/min). AND ● Patient does not have severe liver impairment (e.g. Child Pugh C). AND ● Prescriber must be enrolled with the Adempas REMS Program. AND ● Female patients, regardless of reproductive potential, must be enrolled in the Adempas REMS program prior to starting therapy. AND ● Patient has a diagnosis of persistent/recurrent chronic thromboembolic pulmonary hypertension (CTEPH) (WHO Group 4) after surgical treatment or has inoperable CTEPH OR ● Patient has a diagnosis of pulmonary hypertension and has failed treatment with a preferred product for pulmonary hypertension. (Failure is defined as a lack of efficacy, allergy, intolerable side effects, or significant drug-drug interactions).
--	---	---

Therapeutic Drug Class: LIPOTROPICS - Effective 4/1/2020

<p style="text-align: center;">No PA Required</p> <p>Colesevelam tablet</p> <p>Colestipol tablet</p> <p>Cholestyramine packet, light packet</p>	<p style="text-align: center;">PA Required</p> <p>ANTARA (fenofibrate)</p> <p>Colesevelam packet</p> <p>COLESTID (colestipol) tablet, granules</p> <p>Colestipol granules</p>	<p>Non-preferred bile acid sequestrates may be approved if the member has failed treatment with 2 preferred products in the last 12 months (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Non-preferred fibrates may be approved if the member has failed treatment with generic gemfibrozil or generic fenofibrate and niacin ER in the last 12 months (failure is defined as lack of efficacy with 4 week trial of each drug, allergy, intolerable side effects or significant drug-drug interactions).</p>
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Ezetimibe</p> <p>Fenofibrate capsule, tablet (generic Lofibra/Tricor)</p> <p>Gemfibrozil</p> <p>Niacin ER tablet</p> <p>*Omega-3 ethyl esters cap (generic Lovaza)</p>	<p>Fenofibric acid DR capsule</p> <p>Fenofibric acid tablet</p> <p>LOPID (gemfibrozil)</p> <p>LOVAZA (omega-3 ethyl esters)</p> <p>PREVALITE (cholestyramine/aspartame) packet</p> <p>QUESTRAN (cholestyramine/sugar) packet</p> <p>NIASPAN ER (niacin ER)</p> <p>TRIGLIDE (fenofibrate)</p> <p>TRILIPIX (fenofibric acid)</p> <p>VASCEPA (icosapent ethyl)</p> <p>WELCHOL (colesevalam) tablet, packet</p> <p>ZETIA (ezetimibe)</p>	<p>Non-preferred lipotropic agents with a preferred product with same strength, dosage form, and active ingredient will be approved with adequate trial and/or failure of the preferred product with the same ingredient (such as preferred ezetimibe and Zetia) and 2 additional agents. (Failure is defined as: lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>*Omega-3 ethyl esters (generic Lovaza) may be approved for members who have a baseline triglyceride level ≥ 500 mg/dL</p> <p>Lovaza (brand name) may be approved if meeting the following:</p> <ul style="list-style-type: none"> • Member has a baseline triglyceride level ≥ 500 mg/dl AND • Member has failed an adequate trial of omega-3 Ethyl Esters AND an adequate trial of gemfibrozil or fenofibrate (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions) <p>Vascepa (icosapent ethyl) may be approved if meeting the following:</p> <ul style="list-style-type: none"> • Member has a baseline triglyceride level > 500 mg/dl AND • Member has failed an adequate trial of generic omega-3 ethyl esters AND an adequate trial of gemfibrozil or fenofibrate (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions) <p>OR</p> <ul style="list-style-type: none"> • Vascepa (icosapent ethyl) is being prescribed to reduce CV risk for members on maximally tolerated statin therapy with triglyceride levels ≥ 150mg/dL and LDL-C levels between 41-100 mg/dL AND member meets <u>one</u> of the following: <ul style="list-style-type: none"> ○ Member is ≥ 45 years of age and has established atherosclerotic CV disease (e.g. coronary artery disease, cerebrovascular/carotid disease, peripheral arterial disease) OR ○ Member is ≥ 50 years of age with diabetes mellitus and has <u>one or more</u> of the following additional risk factors for CV disease: <ul style="list-style-type: none"> ▪ Male ≥ 55 years of age or female ≥ 65 years of age ▪ Cigarette smoker ▪ Hypertension ▪ HDL-C ≤ 40 mg/dL for men or ≤ 50 mg/dL for women ▪ hsCRP >3.00 mg/L (0.3 mg/dL) ▪ CrCl 30 to 59 mL/min ▪ Retinopathy ▪ Micro- or macroalbuminuria ▪ ABI <0.9 without symptoms of intermittent claudication
---	--	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		Maximum Dose: Vascepa (icosapent ethyl) 4g daily
--	--	--

Therapeutic Drug Class: STATINS -Effective 4/1/2020		
--	--	--

No PA Required	PA Required	
Atorvastatin tablet	ALTOPREV (lovastatin ER) tablet	Non-preferred Statins may be approved following trial and failure of treatment with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). Age Limitations: Altoprev will not be approved for members < 18 years of age. Fluvastatin and lovastatin will not be approved for members < 10 years of age. Livalo will not be approved for members < 6 years of age.
Lovastatin tablet	CRESTOR (rosuvastatin) tablet	
Pravastatin tablet	EZALLOR (rosuvastatin) sprinkle capsule	
Rosuvastatin tablet	Fluvastatin capsule	
Simvastatin tablet	LESCOL XL (fluvastatin ER) tablet	
	LIPITOR (atorvastatin) tablet	
	LIVALO (pitavastatin) tablet	
	PRAVACHOL (pravastatin) tablet	
	ZOCOR (simvastatin) tablet	

Therapeutic Drug Class: STATIN COMBINATIONS -Effective 4/1/2020		
--	--	--

	PA Required	
	Amlodipine /atorvastatin	Non-preferred Statin combinations may be approved following trial and failure of treatment with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions).
	CADUET (amlodipine/atorvastatin)	
	Ezetimibe/simvastatin	Children: Vytorin will not be approved for members < 18 years of age. Caduet will not be approved for members < 10 years of age.
	VYTORIN (ezetimibe/simvastatin)	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

IV. Central Nervous System

Therapeutic Drug Class: ANTI-CONVULSANTS -Oral-Effective 10/1/2019

<p>No PA Required (age and dosing limits may apply*)</p> <p>Carbamazepine IR tablet, ER tablet, chewable, ER capsule</p> <p>Clobazam tablet</p> <p>Clonazepam tablet, ODT</p> <p>Divalproex capsule, IR tablet, ER tablet</p> <p>DILANTIN^{BNR} (phenytoin) 30 mg capsules</p> <p>Ethosuximide capsule, solution</p> <p>FELBATOL^{BNR} (felbamate) tablet, suspension</p> <p>Lamotrigine tablet, chewable/disperse tabs</p> <p>Levetiracetam IR, ER tablet, solution</p> <p>Oxcarbazepine tablet, suspension</p> <p>Phenobarbital elixir, soln, tab</p> <p>PHENYTEK^{BNR} (phenytoin ER)</p>	<p style="text-align: center;">PA Required</p> <p style="text-align: center;"><i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i></p> <p>APTIOM (eslicarbazepine)</p> <p>BANZEL (rufinamide)</p> <p>BRIVIACT (brivaracetam)</p> <p>CARBATROL ER (carbamazepine)</p> <p>Carbamazepine suspension</p> <p>CELONTIN (methsuximide)</p> <p>DEPAKENE (valproic acid)</p> <p>DEPAKOTE (divalproex)</p> <p>DILANTIN (phenytoin ER) suspension, infatab, 100 mg capsules</p> <p>EPIDIOLEX (cannabidiol)</p> <p>Felbamate tablet, suspension</p> <p>FYCOMPA (perampanel)</p>	<p>Prior Authorization for members currently stabilized (in outpatient or acute care settings) on any non-preferred medication will be approved.</p> <p><u>Non-Preferred Products Newly Started for Treating Seizure Disorder or Convulsions:</u></p> <ul style="list-style-type: none"> ● Non-preferred medications newly started for members with a diagnosis of seizure disorder/convulsions may be approved if meeting the following criteria: <ul style="list-style-type: none"> ○ The medication is being prescribed by a neurologist OR ○ The medication is being prescribed in conjunction with prescriber consultation by a neurologist and meets the following: <ul style="list-style-type: none"> ▪ The prescription meets minimum age and maximum dose limits listed in Table 1 AND ▪ For medications indicated for use as adjunctive therapy, the medication is being used in conjunction with another anticonvulsant medication <p style="text-align: center;">AND</p> <ul style="list-style-type: none"> ○ The prescription meets additional criteria listed for any of the following: <p>Sympazan (clobazam) film:</p> <ul style="list-style-type: none"> ○ Member has history of trial and failure[‡] of clobazam tablet or solution OR ○ Provider attests that member cannot take clobazam tablet or solution <p>Epidiolex (cannabidiol):</p> <ul style="list-style-type: none"> ○ Member has diagnosis of Lennox-Gastaut syndrome (LGS) or Dravet Syndrome <p>Briviact (brivaracetam):</p> <ul style="list-style-type: none"> ○ Member has history of trial and failure[‡] of any levetiracetam-containing product. <p>Aptiom (eslicarbazepine):</p> <ul style="list-style-type: none"> ○ Member has history of trial and failure[‡] of any carbamazepine-containing product. <p>Diacomit (stiripentol):</p> <ul style="list-style-type: none"> ○ Member is concomitantly taking clobazam AND ○ Member has diagnosis of seizures associated with Dravet syndrome <p><u>Non-Preferred Products Newly Started for Non-Seizure Disorder Diagnoses:</u></p>
---	---	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Phenytoin suspension, chewable, ER capsule</p> <p>Primidone tablet</p> <p>TEGRETOL^{BNR} (carbamazepine) suspension</p> <p>Topiramate tablet, sprinkle cap</p> <p>Valproic acid capsule, solution</p> <p>Zonisamide capsule</p>	<p>EQUETRO (carbamazepine)</p> <p>GABITRIL (tiagabine)</p> <p>KEPPRA (levetiracetam) IR tablet, XR tablet, solution</p> <p>KLONOPIN (clonazepam)</p> <p>LAMICTAL (lamotrigine)</p> <p>Lamotrigine ODT, ER tablet</p> <p>MYSOLINE (primidone)</p> <p>ONFI (clobazam)</p> <p>OXTELLAR XR (oxcarbazepine) tablet</p> <p>PEGANONE (ethotoin)</p> <p>QUDEXY XR capsule</p> <p>SPRITAM tablet</p> <p>TEGRETOL (carbamazepine) IR tablet, XR tablet, capsule, chewable</p> <p>Tiagabine tablet</p> <p>TOPAMAX tablet, sprinkle cap</p> <p>Topiramate ER capsule</p> <p>TROKENDI XR capsule</p> <p>TRILEPTAL tablet, suspension</p>	<ul style="list-style-type: none"> Non-preferred medications newly started for non-seizure disorder diagnoses may be approved if meeting the following criteria: <ul style="list-style-type: none"> Member has history of trial and failure[‡] of two preferred agents AND The prescription meets minimum age and maximum dose limits listed in Table 1 <p>[‡]Failure is defined as lack of efficacy, allergy, intolerable side effects, significant drug-drug interaction, or documented contraindication to therapy, or inability to take preferred formulation. Members identified as HLA-B*15:02 positive, carbamazepine and oxcarbazepine should be avoided per Clinical Pharmacogenetics Implementation Consortium Guideline. This may be considered a trial for prior authorization approvals of a non-preferred agent.</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th colspan="3" style="text-align: center;">Table 1: Non-preferred Anticonvulsant Product Table</th> </tr> <tr> <th style="width: 70%;"></th> <th style="width: 15%; text-align: center;">Minimum Age*</th> <th style="width: 15%; text-align: center;">Maximum Dose*</th> </tr> </thead> <tbody> <tr> <td>Mysoline (primidone)</td> <td></td> <td>2000 mg per day</td> </tr> <tr> <td>Dilantin (phenytoin ER)</td> <td></td> <td>1000 mg per loading day 600 mg maintenance dose</td> </tr> <tr> <td>Peganone (ethotoin)</td> <td></td> <td>3000 mg per day</td> </tr> <tr> <td>Celontin (methsuximide)</td> <td></td> <td>Not listed</td> </tr> <tr> <td>Zarontin (ethosuximide)</td> <td></td> <td>Not listed</td> </tr> <tr> <td>Klonopin (clonazepam)</td> <td></td> <td></td> </tr> <tr> <td>Onfi (clobazam) tablet, suspension</td> <td>1 year</td> <td>40 mg per day</td> </tr> <tr> <td>Diacomit (stiripentol)</td> <td>2 years</td> <td>50mg/kg/day</td> </tr> <tr> <td>Aptiom (eslicarbazepine)</td> <td>4 years</td> <td>1600 mg per day</td> </tr> <tr> <td>Carbatrol (carbamazepine ER)</td> <td></td> <td>1600 mg per day</td> </tr> <tr> <td>Epitol (carbamazepine)</td> <td></td> <td>1600 mg per day</td> </tr> <tr> <td>Equetro (carbamazepine ER)</td> <td></td> <td>1600 mg per day</td> </tr> <tr> <td>Oxtellar XR (oxcarbazepine ER)</td> <td></td> <td>Not listed</td> </tr> <tr> <td>Tegretol (carbamazepine) all except suspension</td> <td></td> <td>Not listed</td> </tr> <tr> <td>Tegretol XR (carbamazepine ER)</td> <td></td> <td>Not listed</td> </tr> <tr> <td>Trileptal (oxcarbazepine)</td> <td></td> <td>Not listed</td> </tr> <tr> <td>Depakene (valproic acid)</td> <td>10 years</td> <td></td> </tr> <tr> <td>Depakote (divalproex DR)</td> <td>10 years</td> <td></td> </tr> <tr> <td>Depakote ER (divalproex ER)</td> <td>10 years</td> <td></td> </tr> <tr> <td>Depakote Sprinkle (divalproex DR)</td> <td>10 years</td> <td></td> </tr> <tr> <td>Lamictal (lamotrigine)</td> <td>2 years</td> <td>400 mg per day</td> </tr> <tr> <td>Lamictal ODT (lamotrigine)</td> <td>2 years</td> <td>400 mg per day</td> </tr> <tr> <td>Lamictal XR (lamotrigine ER)</td> <td>13 years</td> <td>600 mg per day</td> </tr> </tbody> </table>	Table 1: Non-preferred Anticonvulsant Product Table				Minimum Age*	Maximum Dose*	Mysoline (primidone)		2000 mg per day	Dilantin (phenytoin ER)		1000 mg per loading day 600 mg maintenance dose	Peganone (ethotoin)		3000 mg per day	Celontin (methsuximide)		Not listed	Zarontin (ethosuximide)		Not listed	Klonopin (clonazepam)			Onfi (clobazam) tablet, suspension	1 year	40 mg per day	Diacomit (stiripentol)	2 years	50mg/kg/day	Aptiom (eslicarbazepine)	4 years	1600 mg per day	Carbatrol (carbamazepine ER)		1600 mg per day	Epitol (carbamazepine)		1600 mg per day	Equetro (carbamazepine ER)		1600 mg per day	Oxtellar XR (oxcarbazepine ER)		Not listed	Tegretol (carbamazepine) all except suspension		Not listed	Tegretol XR (carbamazepine ER)		Not listed	Trileptal (oxcarbazepine)		Not listed	Depakene (valproic acid)	10 years		Depakote (divalproex DR)	10 years		Depakote ER (divalproex ER)	10 years		Depakote Sprinkle (divalproex DR)	10 years		Lamictal (lamotrigine)	2 years	400 mg per day	Lamictal ODT (lamotrigine)	2 years	400 mg per day	Lamictal XR (lamotrigine ER)	13 years	600 mg per day
Table 1: Non-preferred Anticonvulsant Product Table																																																																													
	Minimum Age*	Maximum Dose*																																																																											
Mysoline (primidone)		2000 mg per day																																																																											
Dilantin (phenytoin ER)		1000 mg per loading day 600 mg maintenance dose																																																																											
Peganone (ethotoin)		3000 mg per day																																																																											
Celontin (methsuximide)		Not listed																																																																											
Zarontin (ethosuximide)		Not listed																																																																											
Klonopin (clonazepam)																																																																													
Onfi (clobazam) tablet, suspension	1 year	40 mg per day																																																																											
Diacomit (stiripentol)	2 years	50mg/kg/day																																																																											
Aptiom (eslicarbazepine)	4 years	1600 mg per day																																																																											
Carbatrol (carbamazepine ER)		1600 mg per day																																																																											
Epitol (carbamazepine)		1600 mg per day																																																																											
Equetro (carbamazepine ER)		1600 mg per day																																																																											
Oxtellar XR (oxcarbazepine ER)		Not listed																																																																											
Tegretol (carbamazepine) all except suspension		Not listed																																																																											
Tegretol XR (carbamazepine ER)		Not listed																																																																											
Trileptal (oxcarbazepine)		Not listed																																																																											
Depakene (valproic acid)	10 years																																																																												
Depakote (divalproex DR)	10 years																																																																												
Depakote ER (divalproex ER)	10 years																																																																												
Depakote Sprinkle (divalproex DR)	10 years																																																																												
Lamictal (lamotrigine)	2 years	400 mg per day																																																																											
Lamictal ODT (lamotrigine)	2 years	400 mg per day																																																																											
Lamictal XR (lamotrigine ER)	13 years	600 mg per day																																																																											

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)	
------------------	----------------------	---	--

<p>SABRIL (vigabatrin) powder packet and tablet</p> <p>Vigadrone powder packet</p> <p>Vigabatrin tablet</p> <p>VIMPAT tablet, solution, start kit</p> <p>ZARONTIN capsule, solution</p>	<table border="1"> <tr><td>Qudexy XR (topiramate ER)</td><td>2 years</td><td>400 mg per day</td></tr> <tr><td>Topamax (topiramate)</td><td></td><td>400 mg per day</td></tr> <tr><td>Trokendi XR (topiramate ER)</td><td>6 years</td><td>400 mg per day</td></tr> <tr><td>Briviact (brivaracetam)</td><td>4 years</td><td>200 mg per day</td></tr> <tr><td>Gabitril (tiagabine)</td><td>12 years</td><td>64 mg per day</td></tr> <tr><td>tiagabine</td><td>12 years</td><td>64 mg per day</td></tr> <tr><td>Vimpat (lacosamide)</td><td>4 years</td><td>400 mg per day</td></tr> <tr><td>Banzel (rufinamide)</td><td>1 year</td><td>3200 mg per day</td></tr> <tr><td>Felbamate</td><td>18 years</td><td></td></tr> <tr><td>Fycompa (perampanel)</td><td>4 years</td><td>12 mg per day</td></tr> <tr><td>Sabril (vigabatrin)</td><td>1 month</td><td>3000 mg per day</td></tr> <tr><td>Spritam (levetiracetam)</td><td>4 years</td><td>3000 mg per day</td></tr> <tr><td>Vigabatrin</td><td>1 month</td><td>3000 mg per day</td></tr> <tr><td>Zonegran (zonisamide)</td><td>16 years</td><td>600 mg per day</td></tr> <tr><td>Keppra (levetiracetam)</td><td></td><td>3000 mg per day</td></tr> <tr><td>Keppra XR (levetiracetam ER)</td><td>12 years</td><td>3000 mg per day</td></tr> <tr><td>Epidiolex (cannabidiol)</td><td>2 years</td><td>20 mg/kg/day</td></tr> </table> <p>** Limits based on data from FDA package insert. Approval for age/dosing that falls outside of the indicated range may be evaluated on a case-by-case basis</p>	Qudexy XR (topiramate ER)	2 years	400 mg per day	Topamax (topiramate)		400 mg per day	Trokendi XR (topiramate ER)	6 years	400 mg per day	Briviact (brivaracetam)	4 years	200 mg per day	Gabitril (tiagabine)	12 years	64 mg per day	tiagabine	12 years	64 mg per day	Vimpat (lacosamide)	4 years	400 mg per day	Banzel (rufinamide)	1 year	3200 mg per day	Felbamate	18 years		Fycompa (perampanel)	4 years	12 mg per day	Sabril (vigabatrin)	1 month	3000 mg per day	Spritam (levetiracetam)	4 years	3000 mg per day	Vigabatrin	1 month	3000 mg per day	Zonegran (zonisamide)	16 years	600 mg per day	Keppra (levetiracetam)		3000 mg per day	Keppra XR (levetiracetam ER)	12 years	3000 mg per day	Epidiolex (cannabidiol)	2 years	20 mg/kg/day
	Qudexy XR (topiramate ER)	2 years	400 mg per day																																																	
	Topamax (topiramate)		400 mg per day																																																	
	Trokendi XR (topiramate ER)	6 years	400 mg per day																																																	
	Briviact (brivaracetam)	4 years	200 mg per day																																																	
	Gabitril (tiagabine)	12 years	64 mg per day																																																	
	tiagabine	12 years	64 mg per day																																																	
	Vimpat (lacosamide)	4 years	400 mg per day																																																	
	Banzel (rufinamide)	1 year	3200 mg per day																																																	
	Felbamate	18 years																																																		
	Fycompa (perampanel)	4 years	12 mg per day																																																	
	Sabril (vigabatrin)	1 month	3000 mg per day																																																	
	Spritam (levetiracetam)	4 years	3000 mg per day																																																	
	Vigabatrin	1 month	3000 mg per day																																																	
	Zonegran (zonisamide)	16 years	600 mg per day																																																	
	Keppra (levetiracetam)		3000 mg per day																																																	
	Keppra XR (levetiracetam ER)	12 years	3000 mg per day																																																	
	Epidiolex (cannabidiol)	2 years	20 mg/kg/day																																																	

Therapeutic Drug Class: NEWER GENERATION ANTI-DEPRESSANTS -Effective 1/1/2020

No PA Required	PA Required	
<p>Bupropion IR, SR, XL</p> <p>Citalopram tablet, solution</p> <p>Desvenlafaxine succ ER (generic Pristiq) tablet</p> <p>Duloxetine capsule (generic Cymbalta)</p> <p>Escitalopram tablet</p>	<p style="text-align: center;"><i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i></p> <p>APLENZIN ER (bupropion ER) tablet</p> <p>CELEXA (citalopram) tablet</p> <p>CYMBALTA (duloxetine) capsule</p>	<p>Prior authorization for Fetzima, Trintellix, or Viibryd will be approved for members who have failed an adequate trial with four preferred newer generation anti-depressant products (failure is defined as lack of efficacy with 6 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>All non-preferred products not listed above will be approved for members who have failed adequate trial with three preferred newer generation anti-depressant products. If three preferred newer generation anti-depressant products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all preferred products FDA approved for that indication (failure is defined as lack of efficacy with 6 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Fluoxetine capsules, solution</p> <p>Fluvoxamine tablet (generic Luvox)</p> <p>Mirtazapine tablet, ODT</p> <p>Paroxetine IR tablet</p> <p>Sertraline tablet, solution</p> <p>Trazodone tablet</p> <p>Venlafaxine IR tablet</p> <p>Venlafaxine ER capsules</p>	<p>Desvenlafaxine ER (generic Khedzela)</p> <p>Desvenlafaxine fumarate ER</p> <p>Duloxetine capsule (generic Irenka)</p> <p>EFFEXOR XR (venlafaxine ER) capsule</p> <p>Escitalopram solution</p> <p>FETZIMA (levomilnacipran) capsule</p> <p>Fluoxetine tablets, fluoxetine DR capsules</p> <p>Fluvoxamine ER capsule</p> <p>FORFIVO XL (bupropion ER) tablet</p> <p>LEXAPRO (escitalopram) tablet</p> <p>Nefazodone tablet</p> <p>Paroxetine ER tablet</p> <p>PAXIL (paroxetine) tablet, suspension</p> <p>PAXIL CR (paroxetine ER) tablet</p> <p>PEXEVA (paroxetine) tablet</p> <p>PRISTIQ ER (desvenlafaxine succ ER) tablet</p> <p>PROZAC (fluoxetine) pulvule</p>	<p>Citalopram doses higher than 40mg/day for ≤60 years of age and 20mg for >60 years of age will require prior authorization. Please see the FDA guidance at: https://www.fda.gov/drugs/drugsafety/ucm297391.htm for important safety information.</p> <p>Grandfathering: Members currently stabilized on a Non-preferred newer generation antidepressant can receive approval to continue on that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.</p>
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
	REMERON (mirtazapine) tablet, soltab (ODT) SARAFEM (fluoxetine) tablet TRINTELLIX (vortioxetine) tablet Venlafaxine ER tablets VIIBRYD (vilazodone) tablet WELLBUTRIN SR, XL (bupropion) tablet ZOLOFT (sertraline) tablet, solution	
Therapeutic Drug Class: MONOAMINE OXIDASE INHIBITORS (MAOis) -Effective 1/1/2020		
	<p style="text-align: center;">PA Required</p> EMSAM (selegiline) patch MARPLAN (isocarboxazid) tablet NARDIL (phenelzine) tablet Phenelzine tablet Tranylcypromine tablet	<p>Non-preferred products will be approved for members who have failed adequate trial (8 weeks) with three preferred anti-depressant products. If three preferred anti-depressant products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all preferred anti-depressant products FDA approved for that indication. (Failure is defined as: lack of efficacy after 8 week trial, allergy, intolerable side effects, or significant drug-drug interaction)</p> <p>Grandfathering: Members currently stabilized on a Non-preferred MAOi antidepressant can receive approval to continue on that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.</p>
Therapeutic Drug Class: TRICYCLIC ANTI-DEPRESSANTS (TCAs) -Effective 1/1/2020		
<p style="text-align: center;">No PA Required</p> Amitriptyline tablet Doxepin 10mg, 25mg, 50mg, 75mg, 100mg, 150mg capsule	<p style="text-align: center;">PA Required</p> <p style="text-align: center;"><i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as</i></p>	<p>Non-preferred products will be approved for members who have failed adequate trial (8 weeks) with three preferred tricyclic products. If three preferred products are not available for indication being treated, approval of prior authorization for non-preferred products will require adequate trial of all tricyclic preferred products FDA approved for that indication. (Failure is defined as: lack of efficacy after 8 week trial, allergy, intolerable side effects, or significant drug-drug interaction)</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Doxepin solution</p> <p>Imipramine HCl tablet</p> <p>Nortriptyline capsule, solution</p>	<p><i>written” is indicated on the prescription.</i></p> <p>Amoxapine tablet</p> <p>ANAFRANIL (clomipramine) capsule</p> <p>Clomipramine capsule</p> <p>Desipramine tablet</p> <p>Imipramine pamoate capsule</p> <p>Maprotiline tablet</p> <p>NORPRAMIN (Desipramine) tablet</p> <p>PAMELOR (nortriptyline) capsule</p> <p>Protriptyline tablet</p> <p>SURMONTIL (trimipramine) capsule</p> <p>TOFRANIL (imipramine HCl)</p> <p>Trimipramine capsule</p>	<p>Grandfathering: Members currently stabilized on a Non-preferred TCA antidepressant can receive approval to continue on that agent for one year if medically necessary. Verification may be provided from the prescriber or the pharmacy.</p> <p>Silenor (doxepin 3mg, 6mg) approval criteria can be found on the Appendix P</p>
---	--	--

Therapeutic Drug Class: ANTI-PARKINSON’S AGENTS -Effective 4/1/2020

Dopa decarboxylase inhibitors, dopamine precursors and combinations

No PA Required	PA Required	
<p>Carbidopa/Levodopa IR, ER tablet</p>	<p>Carbidopa tablet</p> <p>Carbidopa/Levodopa ODT</p> <p>DUOPA (carbidopa/levodopa) Suspension</p>	<p>Non-preferred agents may be approved with adequate trial and failure of carbidopa-levodopa IR and ER formulations (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Carbidopa or levodopa single agent products may be approved for members with diagnosis of Parkinson’s Disease as add-on therapy to carbidopa-levodopa.</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	INBRIJA (levodopa) capsule for inhalation RYTARY ER (carbidopa/levodopa) capsule SINEMET (carbidopa/levodopa) IR, ER tablet STALEVO (carbidopa/levodopa/entacapone) tablet	<p>Non-preferred medications that <u>are not</u> prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval without meeting trial and failure step therapy criteria.</p> <p>Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred.</p> <p><u>Grandfathering</u>: Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.</p>
--	---	---

MAO-B inhibitors

No PA Required	PA Required	
Selegiline capsule	AZILECT (Rasagiline) tablet	<p>Non-preferred agents may be approved with adequate trial and failure of selegiline capsule or tablet (failure is defined as lack of efficacy with a 4-week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Non-preferred medications that <u>are not</u> prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval without meeting trial and failure step therapy criteria.</p> <p>Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred.</p> <p><u>Grandfathering</u>: Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.</p>
Selegiline tablet	Rasagiline tablet XADAGO (safinamide) tablet ZELAPAR (selegiline) ODT	

Dopamine Agonists

No PA Required	PA Required	
Pramipexole IR tablet	Bromocriptine capsule, tablet	<p>Non-preferred agents may be approved with adequate trial and failure of ropinirole IR AND pramipexole IR (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Non-preferred medications that <u>are not</u> prescribed for Parkinson’s Disease (or an indication related to Parkinson’s Disease) may receive approval without meeting trial and failure step therapy criteria.</p> <p>Members with history of trial and failure of a non-preferred Parkinson’s Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred.</p>
Ropinirole IR tablet	CYCLOSET (bromocriptine) tablet MIRAPEX (pramipexole) IR, ER tablet NEUPRO (rotigotine) patch	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	PARLODEL (bromocriptine) Pramipexole ER tablet REQUIP (ropinirole) tablet, XR tablet Ropinirole ER tablet	<u>Grandfathering:</u> Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.
--	--	--

Other Parkinson's agents

No PA Required	PA Required	
Amantadine cap, tab, syrup Benztropine tablet Trihexyphenidyl tab, elixir	COMTAN (entacapone) tablet Entacapone tablet GOCOVRI (amantadine) capsule NOURIANZ (istradefylline) tablet OSMOLEX ER (amantadine) tab TASMAR (tolcapone) tablet Tolcapone tablet	Non-preferred agents may be approved with adequate trial and failure of two preferred agents (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions). Non-preferred medications that <u>are not</u> prescribed for Parkinson's Disease (or an indication related to Parkinson's Disease) may receive approval without meeting trial and failure step therapy criteria. Members with history of trial and failure of a non-preferred Parkinson's Disease agent that is the brand/generic equivalent of a preferred product (same strength, dosage form and active ingredient) may be considered as having met a trial and failure of the equivalent preferred. <u>Grandfathering:</u> Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.

Therapeutic Drug Class: ATYPICAL ANTI-PSYCHOTICS - Oral - Effective 4/1/2020

No PA Required*	PA Required	
<p style="text-align: center;">For injectable Atypical Antipsychotics please see Appendix P for criteria</p> Aripiprazole tablet Clozapine tablet LATUDA (lurasidone) 2nd line**	<p style="text-align: center;"><i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and "dispense as written" is indicated on the prescription.</i></p> ABILIFY (aripiprazole) tablet, oral soln, ODT, MyCite	Non-preferred products may be approved for members meeting all of the following: <ul style="list-style-type: none"> • Medication is being prescribed for an FDA-Approved indication (Table 1) AND • Prescription meets dose and age limitations (Table 3) AND • Member has history of trial and failure of three preferred products (failure defined as lack of efficacy with 6 week trial, allergy, intolerable side effects, significant drug-drug interactions, or known interacting genetic polymorphism that prevents safe preferred product dosing) <p>*<u>Age Limits:</u> All products including preferred products will require a PA for members younger than the FDA approved age for the agent (Table 3). Members younger than the FDA approved age for the agent who are currently stabilized on an atypical antipsychotic will be eligible for grandfathering. Atypical Antipsychotic prescriptions for members under 5 years of age may require a</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Olanzapine tablet, ODT</p> <p>Quetiapine IR tablet***</p> <p>Quetiapine ER tablet</p> <p>Risperidone tablet, oral soln, ODT</p> <p>Ziprasidone</p>	<p>Aripiprazole oral solution****, ODT</p> <p>CAPLYTA (lumateperone)</p> <p>CLOZARIL (clozapine)</p> <p>Clozapine ODT</p> <p>GEODON (ziprasidone)</p> <p>FANAPT (iloperidone)</p> <p>FAZACLO (clozapine ODT)</p> <p>Iloperidone</p> <p>INVEGA (paliperidone)</p> <p>olanzapine/fluoxetine</p> <p>NUPLAZID (pimavanserin)</p> <p>Paliperidone</p> <p>REXULTI (brexpiprazole)</p> <p>RISPERDAL (risperidone) tablet, M-tab (ODT), oral solution</p> <p>SAPHRIS (asenapine)</p> <p>SEROQUEL IR (quetiapine IR)***</p> <p>SEROQUEL XR (quetiapine ER)***</p> <p>SYMBYAX (olanzapine/fluoxetine)</p> <p>VERSACLOZ (clozapine suspension)</p>	<p>provider-provider telephone consult with a child and adolescent psychiatrist (provided at no cost to provider or member).</p> <p>**Latuda (lurasidone) may be approved for the treatment of schizophrenia or bipolar depression if the member has tried and failed treatment with one preferred product (qualifying diagnosis verified by AutoPA).</p> <p>***Quetiapine IR when given at sub-therapeutic doses may be restricted for therapy. Low-dose quetiapine (<150mg/day) is only FDA approved as part of a drug titration schedule to aid patients in getting to the target quetiapine dose. PA will be required for quetiapine < 150mg per day except for utilization (when appropriate) in members 65 years or older. PA will be approved for members 10-17 years of age with approved diagnosis (Table 3) stabilized on <150mg quetiapine IR per day.</p> <p>****Aripiprazole solution: Aripiprazole <u>tablet</u> quantity limit is 2 tablets/day for pediatric members to allow for incremental dose titration, and use of the preferred tablet formulation should be considered for dose titrations when possible and clinically appropriate. If incremental dose cannot be achieved with titration of the aripiprazole tablet for members < 18 years of age OR for members unable to swallow solid tablet dosage form, aripiprazole solution may be approved. For all other cases, aripiprazole solution is subject to meeting non-preferred product approval criteria listed above.</p> <p>Nuplazid (pimavanserin tartrate) may be approved for the treatment of hallucinations and delusions associated with Parkinson’s Disease psychosis AND following trial and failure of therapy with quetiapine or clozapine (failure will be defined as intolerable side effects, drug-drug interaction, or lack of efficacy).</p> <p>Abilify MyCite may be approved if meeting all of the following:</p> <ul style="list-style-type: none"> • Member has history of adequate trial and failure of 5 preferred agents (one trial must include aripiprazole tablet). Failure is defined as lack of efficacy with 6 week trial on maximally tolerated dose, allergy, intolerable side effects, significant drug-drug interactions AND • Information is provided regarding adherence measures being recommended by provider and followed by member (such as medication organizer or digital medication reminders) AND • Member has history of adequate trial and failure of 3 long-acting injectable formulations of atypical antipsychotics, one of which must contain aripiprazole (failure is defined as lack of efficacy with 8 week trial, allergy, intolerable side effects, significant drug-drug interactions) AND • Abilify MyCite is being used with a MyCite patch and member is using a compatible mobile application. AND
---	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	VRAYLAR (cariprazine) ZYPREXA (olanzapine) ZYPREXA ZYDIS (olanzapine ODT)	<ul style="list-style-type: none"> Medication adherence information is being shared with their provider via a web portal or dashboard. <p><u>Quantity Limits:</u> Quantity limits will be applied to all products (Table 2). In order to receive approval for off-label dosing, the member must have an FDA approved indication and must have tried and failed on the FDA approved dosing regimen.</p> <p><u>Grandfathering:</u> Members currently stabilized on a non-preferred atypical antipsychotic or Latuda can receive approval to continue therapy with that agent for one year.</p>
--	---	---

Table 1: Approved Indications

Drug	Indication
Abilify (aripiprazole)	<ul style="list-style-type: none"> Schizophrenia Acute treatment of manic or mixed episodes associated with bipolar I disorder Adjunctive treatment of major depressive disorder Irritability associated with autistic disorder Treatment of Tourette's Disorder
Caplyta (lumateperone)	<ul style="list-style-type: none"> Schizophrenia
Fanapt (iloperidone)	<ul style="list-style-type: none"> Acute treatment of schizophrenia in adults
Fazaclo, Versacloz (clozapine)	<ul style="list-style-type: none"> Treatment-resistant schizophrenia Reducing the risk of recurrent suicidal behavior in patients with schizophrenia or schizoaffective disorder
Geodon (ziprasidone)	<ul style="list-style-type: none"> Schizophrenia Bipolar I disorder (acute mixed or manic episodes and maintenance treatment as adjunct to lithium or valproate) Acute treatment of agitation in schizophrenia
Latuda (lurasidone)	<ul style="list-style-type: none"> Schizophrenia Bipolar I disorder
Nuplazid (pimavanserin)	<ul style="list-style-type: none"> hallucinations and delusions associated with Parkinson's disease psychosis
Invega (paliperidone)	<ul style="list-style-type: none"> Schizophrenia Schizoaffective disorder
Risperdal (risperidone)	<ul style="list-style-type: none"> Schizophrenia Bipolar mania Irritability associated with autistic disorder
Rexulti (brexpiprazole)	<ul style="list-style-type: none"> Adjunctive therapy to antidepressants for the treatment of major depressive disorder Schizophrenia
Saphris (asenapine)	<ul style="list-style-type: none"> Acute and maintenance of schizophrenia Bipolar mania, monotherapy Maintenance treatment of bipolar I disorder as an adjunct to lithium or divalproex

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Seroquel (quetiapine) Seroquel XR (quetiapine ER)	<ul style="list-style-type: none"> • Treatment of schizophrenia • Acute treatment of manic or mixed episodes associated with bipolar I disorder, as monotherapy or as an adjunct to lithium or divalproex • Maintenance treatment of bipolar I disorder as an adjunct to lithium or divalproex • Adjunctive treatment of major depressive disorder (Seroquel XR only)
Symbyax (olanzapine/fluoxetine)	<ul style="list-style-type: none"> • Treatment resistant depression • Bipolar I disorder
Vraylar (cariprazine)	<ul style="list-style-type: none"> • Schizophrenia • Bipolar (acute treatment)
Zyprexa (olanzapine)	<ul style="list-style-type: none"> • Schizophrenia • Bipolar I disorder

Table 2: Quantity Limits

Brand Name	Generic Name	Quantity Limits
Abilify	Aripiprazole	Maximum one tablet per day (maximum of two tablets per day allowable for members < 18 years of age to accommodate for incremental dose changes)
Caplyta	Lumateperone	Maximum dosage of 42mg per day
Clozaril	Clozapine	Maximum dosage of 900mg per day
Fazaclio	Clozapine	Maximum dosage of 900mg per day
Fanapt	Iloperidone	Maximum two tablets per day
Geodon	Ziprasidone	Maximum two capsules per day
Invega	Paliperidone	Maximum one capsule per day
Latuda	Lurasidone	Maximum one tablet per day (If dosing 160mg for schizophrenia, then max of two tablets per day)
Nuplazid	Pimavanserin	Maximum dosage of 34mg per day
Risperdal	Risperidone	Maximum dosage of 12mg/day
Rexulti	Brexpiprazole	Maximum of 3mg/day for MDD adjunctive therapy, Maximum of 4mg/day for schizophrenia
Saphris	Asenapine	Maximum two tablets per day
Secuado	Asenapine	Maximum 1 patch per day
Seroquel	Quetiapine	Maximum three tablets per day
Seroquel XR	Quetiapine ER	Maximum one tablet per day (for 300mg & 400mg tablets max 2 tablets per day)
Symbyax	Olanzapine/ fluoxetine	Maximum three capsules per day (18mg olanzapine/75mg fluoxetine)

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Vraylar	Cariprazine	Maximum dosage of 6mg/day
Zyprexa	Olanzapine	Maximum one tablet per day
Zyprexa Zydis	Olanzapine ODT	Maximum one tablet per day

Table 3: FDA Approved Pediatric Dosing by Age

Drug	FDA Approved Indication	FDA-Approved Age	Max FDA-Approved Dose
Asenapine (Saphris, Secuado)	APPROVED FOR ADULTS ONLY		
Brexpiprazole (Rexulti)			
Cariprazine (Vraylar)			
Clozapine (Fazaclo, Clozaril)			
Iloperidone (Fanapt)			
Lumateperone (Caplyta)			
Pimavanserin (Nuplazid)			
Quetiapine ER (Seroquel XR)			
Ziprasidone (Geodon)			
Aripiprazole (Abilify)	Autism/Psychomotor Agitation	6-17 years	15mg/day
	Bipolar Disorder/Mixed Mania	10-17 years	30mg/day
	Schizophrenia	13-17 years	30mg/day
	Gilles de la Tourette's Syndrome	6-17 years	20mg/day
Lurasidone (Latuda)	Schizophrenia	13-17 years	80mg/day
	Bipolar Depression	10-17 years	80mg/day
Olanzapine (Zyprexa)	Schizophrenia	13-17 years	10mg/day
Olanzapine (Zyprexa Zydis)	Bipolar Disorder/Mixed Mania	13-17 years	10mg/day
Paliperidone (Invega ER)	Schizophrenia	12-17 years	12mg/day
Risperidone (Risperdal)	Autism/Psychomotor Agitation	5-16 years	3mg/day
	Bipolar Disorder/Mixed Mania	10-17 years	6mg/day
	Schizophrenia	13-17 years	6mg/day
Quetiapine Fumarate (Seroquel)	Schizophrenia	13-17 years	800 mg/day
	Bipolar Disorder/Mixed Mania	10-17 years	600 mg/day

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)	
------------------	----------------------	---	--

Olanzapine/fluoxetine (Symbyax)	Bipolar I disorder	10-17 years	12mg/50mg/day
---------------------------------	--------------------	-------------	---------------

Therapeutic Drug Class: LITHIUM AGENTS -Effective 4/1/2020

No PA Required	PA Required	
<p>Lithium Carbonate capsule</p> <p>Lithium Carbonate tablet</p> <p>Lithium ER tablet</p>	<p style="text-align: center;"><i>Non-preferred brand name medications do not require a prior authorization when the equivalent generic is preferred and “dispense as written” is indicated on the prescription.</i></p> <p>LithoBID ER (lithium ER) tablet</p> <p>Lithium Citrate soln</p>	<p>Non-preferred products may be approved with trial and failure of one preferred agent (failure is defined as lack of efficacy with 6 week trial, allergy, intolerable side effects, significant drug-drug interactions, intolerance to dosage form).</p> <p>Grandfathering: Members currently stabilized on a non-preferred product may receive approval to continue therapy with that product.</p>

Therapeutic Drug Class: CALCITONIN GENE – RELATED PEPTIDE INHIBITORS (CGRPs) -Effective 4/1/2020

PA Required for all agents		
<p>*AIMOVIG (erenumab) autoinjector</p> <p>*EMGALITY 120mg (galcanezumab) pen, syringe</p>	<p>AJOVY (fremanezumab) syringe</p> <p>EMGALITY 100mg (galcanezumab) syringe</p> <p>UBRELVY (ubrogepant) tablet</p>	<p>*Emgality 120mg (galcanezumab) or Aimovig (erenumab) may be approved for members meeting Migraine Prevention Prior Authorization Criteria below.</p> <p><u>Migraine Prevention Prior Authorization Criteria (must meet all of the following):</u></p> <ul style="list-style-type: none"> • Member is 18 years of age or older AND • Member is in need of prevention of episodic or chronic migraine AND • Member has diagnosis of migraine with or without aura AND • Member has tried and failed 2 oral preventative pharmacological agents listed as Level A per American Headache Society/American Academy of Neurology (i.e. divalproex, topiramate, metoprolol, propranolol). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND • Headache count: If prescribed for episodic migraine member has history of 4-14 migraine days per month OR if prescribed for chronic migraine member has history of 15 or more headache days per month where 8 or more were migraine days for three or more months AND • Member does not have history of MI, stroke, TIA, unstable angina, coronary artery bypass surgery, or other revascularization procedures within previous 12 months AND • Prescription meets one of the following:

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<ul style="list-style-type: none"> ○ Medication <u>is not</u> prescribed for chronic migraine with medication overuse headache OR ○ Member is prescribed Aimovig for chronic migraine with medication overuse headache resulting from taking triptans ≥ 10 days/month, non-narcotic analgesics ≥ 15 days/month (such as acetaminophen, NSAID), or a combination of analgesics ≥ 10 days/month (including non-narcotic, ergot, opioid, butalbital) AND member has not been using a migraine prevention medication for 2 months prior to Aimovig prescription <p>AND</p> <ul style="list-style-type: none"> ● Initial authorization will be limited to the following: <ul style="list-style-type: none"> ○ For episodic migraine: Initial authorization will be for 6 months. Continuation (12 month authorization) will require documentation of clinically significant improvement after 4 months use (and documentation of number of migraine days per month) ○ For chronic migraine: Initial authorization will be for 4 months. Continuation (12 month authorization) will require documentation of clinically significant improvement after 3 months use (and documentation of number of migraine days per month) <p><u>Non-Preferred Medications for Migraine Prevention:</u></p> <p>Non-preferred medications for migraine prevention may be approved if the member meets the Migraine Prevention Prior Authorization Criteria above AND the member has history of adequate trial and failure of Emgality 120mg AND Aimovig therapy (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>Members taking a non-preferred agent for migraine prevention that have not shown clinically significant improvement for 4 months for acute episodic migraine treatment or 3 months for chronic migraine treatment will be allowed to transition to a preferred CGRP agent without meeting the “headache count” criteria listed above.</p> <p><u>Non-Preferred Medications for Acute Migraine Treatment or Cluster Headache Treatment:</u></p> <p>Non-preferred medications for acute migraine treatment (Ubrelyvy) may be approved for members meeting all of the following:</p> <ul style="list-style-type: none"> ● Member is 18 years of age or older AND ● Medication is being prescribed to treat migraine headache with moderate to severe pain AND ● Member is not receiving an injectable form of CGRP medication for any indication AND
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<ul style="list-style-type: none"> • Member has history of trial and failure of all of the following (failure is defined as lack of efficacy with 4 week trial, contraindication, allergy, intolerable side effects, or significant drug-drug interaction): <ul style="list-style-type: none"> ○ Three triptans (including at least two different routes of administration) AND ○ Two NSAID agents AND ○ Dihydroergotamine vial or an ergotamine combination product <p>Non-preferred medications for treatment of cluster headache (Emgality 100mg) may be approved for members meeting all of the following:</p> <ul style="list-style-type: none"> • Member is 19-65 years of age AND • Member meets diagnostic criteria for episodic cluster headache (has had no more than 8 attacks per day, a minimum of one attack every other day, and at least 4 attacks during the week prior to this medication being prescribed) AND • Member is not taking other preventative medications to reduce the frequency of cluster headache attacks AND • Member has history of trial and failure of all of the following (failure is defined as lack of efficacy with 4 week trial, contraindication, allergy, intolerable side effects, or significant drug-drug interaction): <ul style="list-style-type: none"> ○ Oxygen therapy AND ○ Sumatriptan subcutaneous or intranasal AND ○ Zolmitriptan intranasal AND • Member is not prescribed this medication for medication overuse headache AND • Member does not have ECG abnormalities compatible with acute cardiovascular event or conduction delay AND • Member does not have a history within the last 6 months of myocardial infarction, unstable angina, percutaneous coronary intervention, coronary artery bypass grafting, deep vein thrombosis, or pulmonary embolism AND • Member does not have a history of stroke, intracranial or carotid aneurysm, intracranial hemorrhage, or vasospastic angina, clinical evidence of peripheral vascular disease, or diagnosis of Raynaud's AND • Initial authorization will be limited to 8 weeks. Continuation (12 month authorization) will require documentation of clinically relevant improvement with no less than 30% reduction in headache frequency in a 4 week period. <p><u>Maximum Dosing:</u> Aimovig (erenumab): 140mg per 30 days Emgality 120mg (galcanezumab): 240mg once as first loading dose then 120mg monthly</p>
--	--	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		Emgality 100mg (galcanezumab): 300mg per 30 days Ajovy (fremanezumab): 225mg monthly or 675mg every three months Ubrelvy 50mg (ubrogepant): 16 tablets/30 days (800mg per 30 days) Ubrelvy 100mg (ubrogepant): 16 tablets/30 days (1600mg per 30 days)
--	--	---

Therapeutic Drug Class: NEUROCOGNITIVE DISORDER AGENTS -Effective 4/1/2020		
---	--	--

<p>*Must meet eligibility criteria</p> <p>*Donepezil 5mg, 10mg tablet</p> <p>*Donepezil ODT</p> <p>*Memantine tablets</p> <p>*Rivastigmine capsule, patch</p>	<p>PA Required</p> <p>ARICEPT (donepezil) tablets (all strengths), ODT</p> <p>Donepezil 23mg tablet</p> <p>EXELON (rivastigmine) cap, patch, soln.</p> <p>Galantamine IR tablet, soln</p> <p>Galantamine ER capsule</p> <p>Memantine ER capsule, IR solution</p> <p>MESTINON (pyridostigmine) tab, syrup</p> <p>NAMENDA IR, XR (memantine)</p> <p>NAMZARIC (memantine/donepezil)</p> <p>RAZADYNE (galantamine) tab, oral soln</p> <p>RAZADYNE ER (galantamine) cap</p>	<p>*Eligibility criteria for Preferred Agents – All preferred products may be approved without PA if the member has a diagnosis of neurocognitive disorder which can be verified by SMART PA.</p> <p>Non-preferred products may be approved if the member has failed treatment with one of the preferred products in the last 12 months. (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions)</p> <p>Members currently stabilized on a non-preferred product may receive approval to continue on that agent for one year if medically necessary and if there is a diagnosis of neurocognitive disorder.</p>
--	---	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

Therapeutic Drug Class: SEDATIVE HYPNOTICS -Effective 4/1/2020		
Non-Benzodiazepines		
<p>No PA Required* (unless age, dose, or duplication criteria apply)</p> <p>Eszopiclone tablet</p> <p>Zaleplon capsule</p> <p>Zolpidem IR tablet</p> <p>Zolpidem ER tablet</p>	<p style="text-align: center;">PA Required</p> <p>AMBIEN (zolpidem) tablet</p> <p>AMBIEN CR (zolpidem) tablet</p> <p>BELSOMRA (suvorexant) tablet</p> <p>EDLUAR (zolpidem) SL tablet</p> <p>INTERMEZZO (zolpidem) SL tablet</p> <p>LUNESTA (eszopiclone) tablet</p> <p>Ramelteon tablet</p> <p>ROZEREM (ramelteon) tablet</p> <p>SONATA (zaleplon) capsule</p> <p>Zolpidem SL tablet</p>	<p>Non-preferred non-benzodiazepine sedative hypnotics may be approved for members who have failed treatment with two preferred non-benzodiazepine agents (failure is defined as lack of efficacy with a 2 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p><u>Children:</u> Prior authorization will be required for all agents for children < 18 years of age.</p> <p><u>Duplications:</u> Only one agent in the sedative hypnotic drug class will be approved at a time (concomitant use of agents in the same sedative hypnotic class or differing classes will not be approved).</p> <p>All sedative hypnotics will require prior authorization for members ≥ 65 years of age when exceeding 90 days of therapy.</p> <p>Belsomra (suvorexant) may be approved for adult members that meet the following:</p> <ul style="list-style-type: none"> • Members has trialed and failed therapy with two preferred agents (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND • Member is not receiving strong inhibitors (such as erythromycin, clarithromycin, telithromycin, itraconazole, ketoconazole, posaconazole, fluconazole, voriconazole, delavirdine, and milk thistle) or inducers (such as carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifampin, rifabutin, rifapentine, dexamethasone, efavirenz, etravirine, nevirapine, darunavir/ritonavir, ritonavir, and St John’s Wort) of CYP3A4 AND • Member does not have a diagnosis of narcolepsy <p>Dayvigo (lemborexant) may be approved for adult member that meet the following:</p> <ul style="list-style-type: none"> • Member has trialed and failed therapy with two preferred agents AND Belsomra (surovexant). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND • Member is not receiving strong inhibitors (such as erythromycin, clarithromycin, telithromycin, itraconazole, ketoconazole, posaconazole, fluconazole, voriconazole, delavirdine, and milk thistle) or inducers (such as carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifampin, rifabutin, rifapentine, dexamethasone, efavirenz, etravirine, nevirapine, darunavir/ritonavir, ritonavir, and St John’s Wort) of CYP3A4 AND • Member does not have a diagnosis of narcolepsy

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

		<p>Rozerem (ramelteon) may be approved for adult members with a history/concern of substance abuse or for documented concern of diversion within the household without failed treatment on a preferred agent</p> <p>Prior authorization will be required for prescribed doses exceeding maximum (Table 1).</p>
--	--	---

Benzodiazepines		
<p>No PA Required* (unless age, dose, or duplication criteria apply)</p> <p>Temazepam 15mg, 30mg capsule</p> <p>Triazolam tablet</p>	<p style="text-align: center;">PA Required</p> <p>Estazolam tablet</p> <p>Flurazepam capsule</p> <p>HALCION (triazolam) tablet</p> <p>RESTORIL (all strengths) capsule</p> <p>Temazepam 7.5mg, 22.5mg capsule</p>	<p>Non-preferred benzodiazepine sedative hypnotics may be approved for members who have trialed and failed therapy with two preferred benzodiazepine agents (failure is defined as lack of efficacy with a 2 week trial, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>Temazepam 7.5mg and 22.5 mg may be approved if the member has trialed and failed temazepam 15mg or 30mg AND one other preferred product (failure is defined as lack of efficacy with a 2 week trail, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p><u>Children:</u> Prior authorization will be required for all sedative hypnotic agents when prescribed for children < 18 years of age.</p> <p><u>Duplications:</u> Only one agent in the sedative hypnotic drug class will be approved at a time (concomitant use of agents in the same sedative hypnotic class or differing classes will not be approved).</p> <p>All sedative hypnotics will require prior authorization for member's ≥ 65 years of age when exceeding 90 days of therapy.</p> <p><u>Grandfathering:</u> Members currently stabilized on a non-preferred benzodiazepine medication may receive authorization to continue that medication.</p> <p>Prior authorization will be required for prescribed doses exceeding maximum (Table 1).</p>

Brand	Generic	Maximum Dose
Non-Benzodiazepine		
Ambien CR	Zolpidem CR	12.5 mg/day
Ambien IR	Zolpidem IR	10 mg/day
Belsomra	Suvorexant	20 mg/day

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	Dayvigo	Lemborexant	10mg/day
	Edluar	Zolpidem sublingual	Men: 10 mg/day Women: 5 mg/day
	Intermezzo	Zolpidem sublingual	Men: 3.5mg/day Women: 1.75 mg/day
	Lunesta	Eszopiclone	3 mg/day
	Sonata	Zaleplon	20 mg/day
	Rozerem	Ramelteon	8 mg/day
	Zolpimist	Zolpidem spray	Men: 10 mg (2 sprays)/day Women: 5 mg (1 spray)/day
	Benzodiazepine		
	Halcion	Triazolam	0.5 mg/day
	Restoril	Temazepam	30 mg/day
	-	Estazolam	2 mg/day
	-	Flurazepam	30 mg/day
	-	Quazepam	15 mg/day

Therapeutic Drug Class: SKELETAL MUSCLE RELAXANTS -Effective 7/1/2019

No PA Required (if under 65 years of age)*	PA Required	
Baclofen (generic Lioresal)	AMRIX ER (cyclobenzaprine ER)	<p>All agents in this class will require a PA for members 65 years of age and older. The maximum allowable approval will be for a 7-day supply.</p> <p>Non-preferred skeletal muscle relaxants will be approved for members who have trialed and failed‡ three preferred agents. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions.)</p> <p>Authorization for any CARISOPRODOL product will be given for a maximum 3-week one-time authorization for members with acute, painful musculoskeletal conditions who have failed treatment with three preferred products within the last 6 months.</p> <p>*Dantrolene will be approved for members 5-17 years of age who have trialed and failed‡ one preferred agent and meet the following criteria:</p> <ul style="list-style-type: none"> ● Documentation of age-appropriate liver function tests AND ● One of following diagnoses: Multiple Sclerosis, Cerebral Palsy, stroke, upper motor neuron disorder, or spinal cord injury ● Dantrolene will be approved for the period of one year
Cyclobenzaprine (generic Flexeril) 5mg and 10mg tablet	Carisoprodol	
Methocarbamol	Chlorzoxazone	
Tizanidine (generic Zanaflex) 2mg and 4mg tablet	Cyclobenzaprine 7.5mg tabs	
	DANTRIUM (dantrolene)	
	*Dantrolene	
	FEXMID (cyclobenzaprine)	
	LORZONE (chlorzoxazone)	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	METAXALL (metaxolone) Metaxolone Orphenadrine PARAFON FORTE (chlorzoxazone) ROBAXIN (methocarbamol) SKELAXIN (metaxalone) SOMA (carisoprodol) Tizanidine 2, 4, 6mg caps ZANAFLEX (tizanidine)	<ul style="list-style-type: none"> • If a member is stabilized on dantrolene at <18 years of age, they may continue to receive approval after turning 18 years of age • (Failure is defined as: lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions.) <p>‡Failure is defined as: lack of efficacy with 14 day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions.</p>
--	---	--

Therapeutic Drug Class: STIMULANTS AND RELATED AGENTS -Effective 10/1/2019

<p>*No PA Required (if age, max daily dose, and diagnosis met)</p> <p><i>Brand/generic changes effective 11/01/19</i></p> <p>Armodafinil (generic Nuvigil)</p> <p>Atomoxetine (generic Strattera)</p> <p>Mixed-amphetamine salts (generic Adderall IR)</p> <p>Mixed-Amphetamine salts ER (generic Adderall XR)</p>	<p align="center">PA Required</p> <p>ADDERALL IR (mixed-amphetamine salts)</p> <p>ADDERALL XR (mixed amphetamine salts ER)</p> <p>ADHANSIA XR (methylphenidate ER) capsule</p> <p>ADZENYS ER, XR ODT (amphetamine)</p> <p>APTENSIO XR (methylphenidate XR)</p> <p>Clonidine ER tablet</p>	<p>*Preferred medications may be approved through AutoPA for indications listed in Table 1 (preferred medications may also receive approval for off-label use for fatigue associated with multiple sclerosis).</p> <p>Prior authorization for non-preferred medications used for indications listed in Table 1 may be approved for members meeting the following criteria (For Sunosi (solriamfetol), refer to criteria listed below):</p> <ul style="list-style-type: none"> • Member has documented failure with three preferred products in the last 24 months if age ≥6 years or documented failure with one preferred product in the last 24 months if age 3 –5 years (Failure is defined as: lack of efficacy with a four week trial, allergy, intolerable side effects, or significant drug-drug interaction). Trial and failure of preferred agents will not be required for members meeting the following: <ul style="list-style-type: none"> • For Daytrana, Methylin solution, Quillichew, Quillivant XR and Dyanavel XR, one preferred trial must include Vyvanse chewable tablet, Focalin XR, Vyvanse capsules or mixed amphetamine salts ER (generic Adderall XR) and member must have a documented difficulty swallowing that are unable to utilize alternative dosing with preferred tablet and capsule formulations.
---	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>CONCERTA (Methylphenidate ER) tablet^{BNR}</p> <p>Dexmethylphenidate IR (generic Focalin)</p> <p>FOCALIN XR ^{BNR} (dexmethylphenidate ER)</p> <p>Guanfacine ER</p> <p>Methylphenidate IR (generic Ritalin IR)</p> <p>Modafinil (generic Provigil)</p> <p>VYVANSE (lisdexamfetamine) capsules, chewables</p>	<p>COTEMPLA XR ODT (methylphenidate ER)</p> <p>D-amphetamine spansule</p> <p>DAYTRANA (methylphenidate transdermal)</p> <p>DESOXYN (methamphetamine)</p> <p>DEXEDRINE (dextroamphetamine)</p> <p>DEXTROSTAT (dextroamphetamine)</p> <p>Dexmethylphenidate (generic Focalin XR)</p> <p>DYANAVEL XR solution (amphetamine)</p> <p>EVEKEO (amphetamine)</p> <p>FOCALIN IR (dexmethylphenidate)</p> <p>INTUNIV (guanfacine ER)</p> <p>JORNAY PM (methylphenidate)</p> <p>KAPVAY (clonidine ER)</p> <p>METADATE ER (methylphenidate ER)</p> <p>Methylphenidate ER (generic Concerta)</p> <p>Methylphenidate ER 72mg (generic Relexxii)</p>	<p>**Max Dose: Prior authorization may be approved for doses that are higher than the listed maximum dose (Table 2) if member meets all of the following criteria:</p> <ul style="list-style-type: none"> ● Member is taking medication for indicated use listed in table 1 AND ● Member has 30 day trial or failure of three different preferred or non-preferred agents at maximum doses listed in table 2 AND ● Documentation of member’s symptom response to maximum doses of three other agents is provided AND ● Member is not taking a sedative hypnotic medication (from sedative hypnotic PDL class, i.e. temazepam, triazolam, zolpidem) <p>Sunosi (solriamfetol) prior authorization will be approved if member meets the following criteria:</p> <ul style="list-style-type: none"> ● Member is 18 years of age or older AND ● Member has diagnosis of either narcolepsy or obstructive sleep apnea (OSA) and is experiencing excessive daytime sleepiness AND ● Member does not have end stage renal disease AND ● If Sunosi is being prescribed for OSA, member has 1 month trial of CPAP AND ● Member has trial and failure of modafinil AND armodafinil AND one other agent in stimulant PDL class (Failure is defined as: lack of efficacy with 4 week trial, allergy, intolerable side effects, or significant drug-drug interaction.)
---	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	<p>Methylphenidate ER (generic Metadate CD, ER, Ritalin LA)</p> <p>METHYLIN SUSPENSION (methylphenidate)</p> <p>MYDAYIS ER (dextroamphetamine/amphetamine)</p> <p>NUVIGIL (armodafinil)</p> <p>PROCENTRA (dextroamphetamine liquid)</p> <p>PROVIGIL (modafinil)</p> <p>QUILLICHEW (methylphenidate)</p> <p>QUILLIVANT XR suspension (methylphenidate)</p> <p>RELEXXII (methylphenidate ER)</p> <p>RITALIN IR (methylphenidate)</p> <p>RITALIN LA (methylphenidate ER (LA))</p> <p>STRATTERA (atomoxetine)</p> <p>SUNOSI (solriamfetol)</p> <p>ZENZEDI (dextroamphetamine)</p>	
--	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Table 1: Indication and Age

- Approval for medically accepted indications not listed in Table 1 may be given with prior authorization review and may require submission of peer-reviewed literature or medical compendia showing safety and efficacy of the medication used for the prescribed indication. Medications may also receive approval for off-label use for fatigue associated with multiple sclerosis if meeting all other criteria for approval.
- Prior authorization will be required for doses that are higher than the FDA approved maximum doses.**
- **Bolded Drug names are Preferred**

Drug	Indications
Stimulants – Immediate Release	
amphetamine sulfate (Evekeo™)	ADHD (Age ≥ 3 years), Narcolepsy (Age ≥ 6 years)
armodafinil (Nuvigil®)	Excessive sleepiness associated with narcolepsy, OSA, and SWD for age ≥ 18 years
dexmethylphenidate IR (Focalin®)	ADHD (Age ≥ 6 years)
dextroamphetamine IR (Zenedi™)	ADHD (Age 3 to ≤ 16 years), Narcolepsy (Age ≥ 6 years)
dextroamphetamine solution (ProCentra™)	ADHD (Age 3 to ≤ 16 years), Narcolepsy (Age ≥ 6 years)
methamphetamine (Desoxyn®)	ADHD (Age ≥ 6 years)
methylphenidate IR (Ritalin®)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years), OSA
methylphenidate IR (Methylin®)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)
methylphenidate XR ODT (Contempla® XR ODT)	ADHD (Age ≥ 6 years)
mixed amphetamine salts IR (Adderall®)	ADHD (Age ≥ 3 years), Narcolepsy (Age ≥ 6 years)
modafinil (Provigil®)	Excessive sleepiness associated with narcolepsy, OSA, and SWD (Age ≥ 18 years)
Solriamfetol (Sunosi®)	Excessive sleepiness associated with narcolepsy, OSA (Age ≥ 18)
Stimulants – Extended-Release	
amphetamine ER (Adzenys® XR-ODT and Adzenys® ER suspension)	ADHD (Age ≥ 6 years)
amphetamine ER (Dyanavel™ XR)	ADHD (Age ≥ 6 years)
Mixed-Amphetamine salts ER (generic Adderall XR)	ADHD (Age ≥ 6 years)
dexmethylphenidate ER (Focalin XR®)	ADHD (Age ≥ 6 years)
dextroamphetamine ER (Dexedrine®)	ADHD (Age 3 to ≤ 16 years), Narcolepsy (Age ≥ 6 years)
dextroamphetamine ER/amphetamine ER (Mydayis ER®)	ADHD (Age ≥ 13 years)
lisdexamfetamine dimesylate (Vyvanse® capsule and Vyvanse® chewable)	ADHD (Age ≥ 6 years), Moderate to severe binge eating disorder in adults (Age ≥ 18 years)
methylphenidate ER OROS (Concerta®)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years), OSA
methylphenidate SR (Metadate ER®)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

methylphenidate ER† (Metadate CD®)	ADHD (Age ≥ 6 years)
methylphenidate ER (QuilliChew™ ER)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)
methylphenidate ER (Quillivant XR®)	ADHD (Age ≥ 6 years), Narcolepsy (Age ≥ 6 years)
methylphenidate ER (Ritalin LA®)	ADHD (Age ≥ 6 years)
methylphenidate ER (Aptensio XR®)	ADHD (Age ≥ 6 years)
methylphenidate XR ODT (Contempla® XR ODT)	ADHD (Age ≥ 6 years)
Methylphenidate ER (Jornay PM ®)	ADHD (Age ≥ 6 years)
Non-Stimulants	
atomoxetine (Strattera®)	ADHD (Age ≥ 6 years)
clonidine ER (Kapvay™)	ADHD (Age ≥ 6 years), Treatment of ADHD as adjunct to stimulants
guanfacine ER (Intuniv™)	ADHD (Age ≥ 6 years), Treatment of ADHD as adjunct to stimulants

Table 2: Max Daily Dose

Drug	Maximum Daily Dose
ADDERALL ®	60 mg/day
ADDERALL XR®	60mg/day
ADZENYS XR-ODT® ADZENYS ER-SUSPENSION®	18.8 mg/day (age 6-12) 12.5 mg/day (age >13)
AMPHETAMINE SALTS	40 mg/day
CONCERTA®	54 mg/day or 72 mg/day >age 13
COTEMPLA XR-ODT®	51.8mg/day
DESOXYN ®	25mg/day
DEXEDRINE ®	40mg/day
DEXTROSTAT ®	40mg/day
DYANAVEL XR ®	20mg/day
FOCALIN ®	20 mg/day
FOCALIN XR ®	40 mg/day
JORNAY PM ®	100mg/day
METHYLPHNIDATE ER	60 mg/day
MYDAYIS ER®	25 mg/day (age 13-17) 50 mg/day (age ≥ 18)
INTUNIV ER®	4 mg/day
RITALIN® IR	60 mg/day
RITALIN SR®	60 mg/day

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	RITALIN LA ®	60 mg/day	
	STRATTERA®	100 mg/day	
	VYVANSE CAPS AND CHEWABLE ®	70 mg/day	
	D-AMPHETAMINE ER	40 mg/day	
	DAYTRANA ®	30 mg/day	
	EVEKEO ®	40 mg/day	
	KAPVAY ER®	0.1 mg/day	
	METHYLIN ER ®	60 mg/day	
	METHYLIN	60 mg/day	
	METHYLIN SUSPENSION®	60 mg/day	
	METADATE CD ®	60mg/day	
	METADATE ER ®	60mg/day	
	METHYLPHENIDATE	60 mg/day	
	PROVIGIL ®	400 mg/day	
	NUVIGIL ®	250 mg/day	
	QUILLIVANT ®	60 mg/day	
	SUNOSI ®	150 mg/day	
	ZENZEDI ®	40 mg/day	

Therapeutic Drug Class: TRIPTANS AND OTHER MIGRAINE TREATMENTS - Oral -Effective 1/1/2020

No PA Required (monthly quantity limits may apply)	PA Required	
	Almotriptan tablet	<p>Non-preferred oral triptan products may be approved for members who have trialed and failed three preferred oral products. Failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interaction.</p> <p>Quantity Limits:</p> <p>Amerge (naratriptan), Frova (frovatriptan), Imitrex (sumatriptan), Zomig (zolmitriptan): Max 9 tabs/30 days</p> <p>Treximet (sumatriptan/naproxen): Max 9 tabs/30 days</p> <p>Axert (almotriptan) and Relpax (eletriptan): Max 6 tabs/30 days</p> <p>Maxalt (rizatriptan): Max 12 tabs/30 days</p>
Eletriptan tablet (generic Relpax)	AMERGE (naratriptan) tablet	
Naratriptan tablet (generic Amerge)	FROVA (frovatriptan) tablet	
Rizatriptan tablet, ODT (generic Maxalt)	IMITREX (sumatriptan) tablet	
Sumatriptan tablet (generic Imitrex)	MAXALT (rizatriptan) tablet, MLT	
	RELPAX (eletriptan) tablet	
	REYVOW (lasmiditan) tablet	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	Sumatriptan/Naproxen tablet TREXIMET (sumatriptan/ naproxen) tablet Zolmitriptan tablet, ODT ZOMIG (zolmitriptan) tablet, ZMT	
--	--	--

Therapeutic Drug Class: TRIPTANS AND OTHER MIGRAINE TREATMENTS - Non-Oral -Effective 1/1/2020

No PA Required (monthly quantity limits may apply)	PA Required	
Sumatriptan vial	IMITREX (sumatriptan) nasal spray, cartridge, injection, pen injector	Non-preferred non-oral products will be approved for members who have trialed and failed two preferred non-oral products. Failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions, documented inability to tolerate dosage form. Zembrace Symtouch injection, Tosymra nasal spray, or Onzetra Xsail nasal powder may be approved for members who have trialed and failed two preferred non-oral triptan products AND have trialed and failed two oral triptan agents. Failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interaction, documented inability to tolerate dosage form. Quantity Limits: Imitrex (sumatriptan) injection: Max 4 injectors / 30 days Imitrex (sumatriptan) nasal spray: Max 6 inhalers / 30 days Zomig (zolmitriptan) nasal spray: Max 6 inhalers / 30 days Zembrace Symtouch (sumatriptan) injection: Max 36mg / 30 days Onzetra Xsail (sumatriptan) nasal powder: Max 16 nosepieces / 30 days Tosymra (sumatriptan) nasal spray: 12 nasal spray devices / 30 days
ZOMIG (zolmitriptan) nasal spray	ONZETRA XSAIL (sumatriptan) nasal powder	
	SUMAVEL DOSEPRO (sumatriptan) injection	
	Sumatriptan cartridge, injection, syringe, nasal spray	
	TOSYMRA (sumatriptan) nasal spray	
	ZEMBRACE SYMTOUCH (sumatriptan) injection	

V. Dermatological

Therapeutic Drug Class: ACNE – Topical -Effective 7/1/2019

No PA Required (if age and diagnosis criteria is met*)	PA Required	
*Adapalene gel	ACANYA gel, pump	Authorization for all acne agents prescribed solely for cosmetic purposes will not be approved. Preferred topical acne agents prescribed for members > 25 years of age will require prior authorization and will be approved following prescriber verification that the medication is not being

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

<p>*Adapalene/benzoyl peroxide (generic Epiduo)</p> <p>*Clindamycin phosphate med swab</p> <p>*Clindamycin phosphate solution</p> <p>*Clindamycin/benzoyl peroxide gel jar (generic Benzaclin)</p> <p>*DIFFERIN gel pump (adapalene)^{BNR}</p> <p>*Erythromycin soln</p> <p>*RETIN-A cream^{BNR}</p> <p>*Sodium sulfacetamide/sulfur cleanser, wash</p> <p>*Sulfacetamide suspension</p> <p>Tretinoin gel</p>	<p>ACZONE gel, pump</p> <p>Adapalene cream, gel pump, soln</p> <p>AKTIPAK (erythromycin/benzoyl peroxide)</p> <p>ALTRENO (tretinoin)</p> <p>ATRALIN (tretinoin) gel</p> <p>AVAR (all products)</p> <p>AVITA (tretinoin) cream, gel</p> <p>AZELEX (azelaic acid)</p> <p>BENZAC (benzoyl peroxide)</p> <p>BENZACLIN (all products)</p> <p>Benzoyl peroxide gel, kit, lotion, med pad, microspheres, towelette</p> <p>Benzoyl peroxide / sulfur</p> <p>CLINDACIN PAC Kit</p> <p>Clindamycin phosphate gel, lotion, foam</p> <p>Clindamycin/benzoyl peroxide (generic Duac)</p> <p>Clindamycin/benzoyl peroxide w/ pump (generic Benzaclin)</p> <p>Clindamycin/tretinoin</p>	<p>utilized for cosmetic purposes AND prescriber verification that the indicated use is for acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne. These medications are only eligible for prior authorization approval for the aforementioned diagnoses.</p> <p>Preferred topical acne agents prescribed for members ≤ 25 years of age will only be approved for members with a diagnosis of acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne. Diagnosis will be verified through automated verification (AutoPA) of the appropriate corresponding ICD-10 diagnosis code related to the indicated use of the medication.</p> <p>Preferred topical clindamycin and erythromycin products prescribed for members ≤ 25 may also be approved for a diagnosis of folliculitis, hidradenitis suppurativa, or perioral dermatitis (erythromycin only). Approval of preferred topical clindamycin and erythromycin products for other medically accepted indications for members ≤ 25 may be considered following clinical prior authorization review by a call center pharmacist.</p> <p>Non-preferred topical products will be approved for members meeting all of the following criteria:</p> <ul style="list-style-type: none"> ● Member has trialed/failed three preferred topical products with different mechanisms (i.e. tretinoin, antibiotic). Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction AND ● Prescriber verification that the medication is being prescribed for one of the following diagnoses: acne vulgaris, psoriasis, cystic acne, disorders of keratinization, neoplasms, or comedonal acne.
---	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

	<p>Dapsone gel</p> <p>DIFFERIN (adapalene) cream, gel, lotion</p> <p>EPIDUO (all products)</p> <p>Erythromycin gel, med swab</p> <p>Erythromycin / Benzoyl peroxide</p> <p>ONEXTON (clindamycin/benzyoyl peroxide)</p> <p>OVACE (all products)</p> <p>RETIN-A gel</p> <p>RETIN-A Micro (all products)</p> <p>Sulfacetamide cleanser</p> <p>Sulfacetamide sodium/ sulfur cream, lotion, cleanser kit</p> <p>TAZORAC cream, gel</p> <p>Tazarotene cream</p> <p>Tretinoin cream (generic Retin-A, Avita)</p> <p>Tretinoin gel (generic Atralin)</p> <p>Tretinoin microspheres (all products)</p>	
--	---	--

Therapeutic Drug Class: **ACNE – ISOTRETINOIN** -Effective 7/1/2019

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

PA Required for all agents		
AMNESTEEM capsule	ABSORICA capsule	All preferred and non-preferred oral isotretinoin agents will require prior authorization and will be approved for severe, recalcitrant nodulocystic acne for adults and children ≥ 12 years of age and has been unresponsive to conventional therapy AND
CLARAVIS capsule	Isotretinoin capsule	
	MYORISAN capsule	Non-preferred oral isotretinoin agents will be approved if member has trialed/failed two preferred agents. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.
	ZENATANE capsule	

Therapeutic Drug Class: ANTI-PSORIATICS - Oral -Effective 1/1/2020

No PA Required	PA Required	
SORIATANE ^{BNR} (acitretin) capsule	Acitretin capsule	Prior authorization for non-preferred oral agents will be approved with failure of two preferred anti-psoriatic agents, one of which must be a preferred oral agent. Failure is defined as lack of efficacy of a 4 week trial, allergy, intolerable side effects or significant drug-drug interaction.
	Methoxsalen capsule, softgel	
	Methoxsalen Rapid	
	OXSORALEN-ULTRA (methoxsalen) capsule	

Therapeutic Drug Class: ANTI-PSORIATICS -Topical -Effective 1/1/2020

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

No PA Required	PA Required	
Calcipotriene solution DOVONEX ^{BNR} (calcipotriene) cream TACLONEX SCALP ^{BNR} (calcipotriene/betamethasone) susp TACLONEX OINTMENT ^{BNR} (calcipotriene/betamethasone)	Calcipotriene cream, ointment Calcipotriene/betamethasone dp ointment Calcitriol ointment DUOBRII (halobetasol/tazarotene) lotion ENSTILAR (calcipotriene/betamethasone) foam SORILUX (calcipotriene) foam VECTICAL (calcitriol) ointment	<p>Prior authorization for non-preferred topical agents will be approved with failure of two preferred topical agents. If non-preferred topical agent being requesting is a combination product, trial of two preferred agents must include a preferred combination agent. Failure is defined as lack of efficacy of a 4 week trial, allergy, intolerable side effects or significant drug-drug interaction.</p> <p>Preferred and non-preferred products that contain a corticosteroid ingredient (such as betamethasone) will be limited to 4 weeks of therapy. Continued use will require one week of steroid-free time in between treatment periods.</p> <p>Members with >30% of their body surface area affected may not use Enstilar (calcipotriene/betamethasone DP) foam or Taclonex (calcipotriene/betamethasone DP) ointment products as safety and efficacy have not been established.</p>

Therapeutic Drug Class: ROSACEA AGENTS -Effective 7/1/2019

No PA Required	PA Required	
<p><i>Brand/generic changes effective 10/15/19</i></p> Azelaic acid gel Metronidazole cream, gel, lotion	FINACEA (azelaic acid) foam, gel METROCREAM (metronidazole) METROGEL (metronidazole) METROLOTION (metronidazole) MIRVASO (brimonidine) ORACEA (doxycycline)* NORITATE (metronidazole) RHOFADE (oxymetazoline)	<p>Prior authorization for non-preferred products in this class may be approved if member meets the following criteria:</p> <ul style="list-style-type: none"> ● Member has a diagnosis of persistent (non-transient) facial erythema with inflammatory papules and pustules due to rosacea AND ● Prescriber attests that medication is not being used solely for cosmetic purposes AND ● Member has tried and failed two preferred agents of different mechanisms of action (Failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects) <p>*Oracea® (doxycycline monohydrate DR) may be approved if the member meets all of the following criteria:</p> <ul style="list-style-type: none"> ● Member has taken generic doxycycline for a minimum of three months and failed therapy in the last 6 months. Failure is defined as: lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions AND ● Member has history of an adequate trial/failure (8 weeks) of 2 other preferred agents (oral or topical). Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions AND

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

	ROSADAN Kit (metronidazole) SOOLANTRA (ivermectin)	<ul style="list-style-type: none"> Member is ≥ 18 years of age and has been diagnosed with rosacea with inflammatory lesions
--	---	---

Therapeutic Drug Class: TOPICAL STEROIDS – Effective 4/1/2020

Low potency

No PA Required	PA Required	
Hydrocortisone (Rx) cream, ointment, lotion DERMA-SMOOTHIE-FS ^{BNR} (fluocinolone acetonide) oil Desonide 0.05% cream, ointment Fluocinolone acetonide 0.01% cream	ALA-CORT (hydrocortisone) cream ALA-SCALP (hydrocortisone) lotion Alclometasone cream, ointment CAPEX (fluocinolone) shampoo DESONATE (desonide) gel Desonide lotion DESOWEN (desonide) cream Fluocinolone acetonide 0.01% body oil, 0.01% scalp oil, 0.01% solution MICORT-HC (hydrocortisone) cream SYNALAR (fluocinolone) 0.01% solution TEXACORT (hydrocortisone) solution	Non-preferred Low Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the Low Potency class (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions).

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Medium potency		
No PA Required	PA Required	
Betamethasone dipropionate 0.05% lotion	BESER (fluticasone) lotion	Non-preferred Medium Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the Medium Potency class (failure is defined as: lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions).
Betamethasone valerate 0.1% ointment	Betamethasone dipropionate 0.05% cream	
Fluticasone propionate 0.05% cream, 0.05% ointment	Betamethasone valerate 0.1% cream, 0.1% lotion, 0.12% foam	
Mometasone furoate 0.1% cream, 0.1% ointment, 0.1% solution	Clocortolone cream, cream pump	
Triamcinolone acetonide 0.025% cream, 0.1% cream, 0.025% ointment, 0.1% ointment, 0.025% lotion, 0.1% lotion	CLODERM (clocortolone) cream, cream pump	
	CORDRAN (flurandrenolide) tape	
	CUTIVATE (fluticasone) cream, lotion	
	DERMATOP (prednicarbate) ointment	
	DERMATOP EMOLLIENT (prednicarbate) cream	
	Diflorasone cream	
	ELOCON (mometasone) cream	
	Fluocinolone acetonide 0.025% cream, ointment	
	Fluocinonide-E cream 0.05%	
	Flurandrenolide cream, ointment, lotion	
	Fluticasone propionate 0.05% lotion	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	<p>Hydrocortisone butyrate 0.1% cream, 0.1% lotion, 0.1% solution, 0.1% ointment, 0.1% lipid/lipocream</p> <p>Hydrocortisone valerate 0.2% cream, 0.2% ointment</p> <p>KENALOG (triamcinolone) spray</p> <p>LOCOID (hydrocortisone butyrate) cream, ointment, lotion, solution</p> <p>LOCOID LIPOCREAM 0.1% (hydrocortisone butyrate)</p> <p>LUXIQ (betamethasone valerate) foam</p> <p>ORALONE (triamcinolone) paste</p> <p>PANDEL (hydrocortisone probutate) cream</p> <p>Prednicarbate cream, ointment</p> <p>PSORCON (diflorasone) cream</p> <p>SERNIVO (betamethasone dipropionate) spray</p> <p>SYNALAR (fluocinolone) 0.025% cream/kit, ointment/kit</p> <p>SYNALAR TS (fluocinolone) 0.01%</p> <p>Triamcinolone 0.1% paste, 0.147 mg/gm spray</p>	
--	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

High potency		
<p>No PA Required (unless exceeds duration of therapy*)</p> <p>*Betamethasone dipropionate propylene glycol (aug) 0.05% cream</p> <p>*Fluocinonide 0.05% gel, 0.05% solution, 0.05% ointment</p> <p>*Triamcinolone acetonide 0.5% cream, 0.5% ointment</p>	<p>PA Required</p> <p>Amcinonide cream, lotion</p> <p>APEXICON-E (diflorasone) cream</p> <p>Betamethasone dipropionate 0.05% ointment</p> <p>Desoximetasone cream, gel, ointment</p> <p>Diflorasone ointment</p> <p>Fluocinonide 0.05% cream</p> <p>Halcinonide cream</p> <p>HALOG (halcinonide) cream, ointment</p> <p>TOPICORT (desoximetasone) cream, gel, ointment</p> <p>TRIANEX (triamcinolone) Ointment</p>	<p>Non-preferred High Potency topical corticosteroids may be approved following adequate trial and failure of two preferred agents in the High Potency class (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>*All High Potency topical corticosteroids will require prior authorization beyond 4 weeks of therapy. The provider will be encouraged to transition to a moderate or low potency topical steroid after this time has elapsed.</p>
Very high potency		
<p>No PA Required (unless exceeds duration of therapy*)</p> <p>*Betamethasone dipropionate propylene glycol (aug) 0.05% ointment</p> <p>*Clobetasol 0.05% cream, 0.05% gel, 0.05% ointment, 0.05% solution</p>	<p>PA Required</p> <p>Betamethasone dipropionate propylene glycol (aug) 0.05% gel, 0.05% lotion</p> <p>BRYHALI (halobetasol) lotion</p> <p>Clobetasol emollient/emulsion cream, foam</p>	<p>Non-preferred Very High Potency topical corticosteroids may be approved following adequate trial and failure of clobetasol propionate in the same formulation as the product being requested (if the formulation of the requested non-preferred product is not available in preferred clobetasol product options, then trial and failure of any preferred clobetasol product formulation will be required). Failure is defined as lack of efficacy with 2 week trial, allergy, intolerable side effects or significant drug-drug interactions.</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

	<p>Clobetasol lotion, foam, spray, shampoo</p> <p>CLOBEX (clobetasol) 0.05% lotion, 0.05% spray, 0.05% shampoo</p> <p>CLODAN (clobetasol) 0.05% shampoo, kit</p> <p>Desoximetasone spray</p> <p>DIPROLENE (betamethasone dipropionate/glycol) ointment</p> <p>Fluocinonide 0.1% cream</p> <p>Halobetasol cream, ointment, foam</p> <p>LEXETTE (halobetasol) foam</p> <p>OLUX (clobetasol) foam</p> <p>OLUX-E (clobetasol) foam</p> <p>TEMOVATE (clobetasol) cream, ointment</p> <p>TOPICORT (desoximetasone) spray</p> <p>TOVET EMOLLIENT (clobetasol) foam</p> <p>ULTRAVATE (halobetasol) lotion, cream, ointment</p> <p>ULTRAVATE-X (halobetasol/lactic acid) cream, ointment</p>	<p>*All Very High Potency topical corticosteroids will require prior authorization beyond 2 weeks of therapy. If clobetasol propionate shampoo is being used to treat plaque psoriasis, then prior authorization will be required beyond 4 weeks of therapy. The provider will be encouraged to transition to a moderate or low potency topical steroid after this time has elapsed.</p>
--	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	VANOS (fluocinonide) cream	
--	----------------------------	--

VI. Endocrine

Therapeutic Drug Class: ANDROGENIC AGENTS -Effective 7/1/2019

*Must meet criteria	PA Required	
Testosterone 1.62% packet (generic Androgel)	ANDROGEL 1.62% (testosterone gel) pump	<p><u>Hypogonadotropic or Primary Hypogonadism (may be secondary to Klinefelter Syndrome):</u> Preferred androgenic drugs will be approved for members meeting the following:</p> <ol style="list-style-type: none"> 1. Male patient > 16 years of age AND 2. Has a documented diagnosis of hypogonadotropic or primary hypogonadism (Patients with other diagnoses will require a manual review) AND 3. Has two documented low serum testosterone levels below the lower limit of normal range for testing laboratory prior to initiation of therapy AND 4. Does not have a diagnosis of breast or prostate cancer AND 5. Does not have a palpable prostate nodule or prostate-specific antigen (PSA) > 4ng/mL AND 6. Has normal liver function tests prior to initiation of therapy <p><u>Gender Transition/Affirming Hormone Therapy:</u> Preferred androgenic drugs will be approved for members meeting the following:</p> <ol style="list-style-type: none"> 1. Female sex assigned at birth > 16 years of age AND 2. Is undergoing female to male transition AND 3. Has a negative pregnancy test prior to initiation AND 4. Has normal liver function tests prior to initiation of therapy <p>*Testosterone 1.62% packet (generic Androgel®) is a preferred agent for gender transition/affirmation and is non-preferred for all other indications.</p> <p>Non-preferred <u>topical</u> androgenic agents may be approved for patients meeting the above criteria with trial and failed‡ therapy with two preferred topical androgen formulations.</p> <p>Non-preferred <u>injectable</u> androgenic agents may be approved for patients meeting the above criteria with trial and failed‡ therapy with a preferred injectable androgenic drug.</p> <p>Prior authorization for <u>oral</u> androgen agents (tablet, capsule, buccal) may be approved if member has trialed and failed‡ therapy with a preferred topical agent AND testosterone cypionate injection.</p>
*ANDRODERM (testosterone) patch	ANDROGEL 1% (testosterone gel)	
*Testosterone gel pump (generic Axiron)	ANDROID (methyltestosterone) capsule	
*Testosterone gel (generic Fortesta)	DELATESTRYL (testosterone enanthate) IM injection	
*Testosterone gel (generic Testim)	DEPO TESTOSTERONE (testosterone cypionate) IM injection	
*Testosterone gel, packet, pump (generic Vogelxo)	FORTESTA (testosterone) gel	
*Testosterone gel, packet, pump (generic Vogelxo)	Jatenzo (testosterone undecanoate) capsules	
*Testosterone cypionate IM injection	METHITEST (methyltestosterone) tablet	
<p><i>Injectable testosterone cypionate is a pharmacy benefit when self-administered. Administration in an office setting is a medical benefit.</i></p>	Methyltestosterone capsule	
	STRIANT (testosterone) buccal	
	TESTIM (testosterone gel)	
	Testone CIK (testosterone cypionate) IM injection	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	<p>TESTRED (methyltestosterone) capsule</p> <p>Testosterone enanthate IM injection</p> <p>Testosterone gel 1.62% 1.25 gram/actuation pump</p> <p>VOGELXO (testosterone) gel</p> <p>XYOSTED (testosterone enanthate) SC injection</p>	<p>‡Failure is defined as lack of efficacy with 8 week trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction.</p> <p>For all agents and diagnoses, members < 16 years of age will require a manual prior authorization review by a pharmacist.</p> <p><u>Reauthorization Criteria (for Hypogonadism diagnoses):</u></p> <p>Members may continue to receive preferred agents without requirement of updated low serum testosterone laboratory testing that meet the following criteria:</p> <ul style="list-style-type: none"> ● Male patient > 16 years of age AND ● Has at least one past documented low serum testosterone levels below the lower limit of normal range for testing laboratory prior to initiation of therapy AND ● Has documented diagnosis of hypogonadotropic or primary hypogonadism AND ● Does not have a diagnosis of breast or prostate cancer AND ● Does not have a palpable prostate nodule or prostate-specific antigen (PSA) > 4ng/mL AND ● Has normal liver function tests prior to initiation of therapy
--	--	--

Therapeutic Drug Class: BONE RESORPTION SUPPRESSION AND RELATED AGENTS -Effective 10/1/2019

Bisphosphonates

No PA Required	PA Required	
<p>Alendronate (generic) 5mg, 10mg, 35mg, 70mg tablets</p> <p>Ibandronate tablet</p>	<p>ACTONEL (risedronate)</p> <p>ACTONEL w/Calcium (risedronate w/calcium)</p> <p>Alendronate 40mg tab</p> <p>Alendronate oral solution</p> <p>AELVIA (risedronate)</p> <p>BINOSTO (alendronate)</p> <p>BONIVA (ibandronate)</p>	<p>Non-preferred bisphosphonates may be approved for members who have failed treatment with one preferred product at treatment dose. (Failure is defined as: lack of efficacy with a 12 month trial, allergy, intolerable side effects, or significant drug-drug interaction.)</p> <p>Prior authorization for alendronate 70mg/75ml solution will be approved if member cannot swallow solid oral dosage forms or has a feeding tube.</p> <p>Prior authorization may be approved for etidronate in members with heterotopic ossification without treatment failure of a preferred agent.</p> <ul style="list-style-type: none"> ● For members who have a low risk of fracture, prior authorization will be required for members exceeding 5 years of either a preferred or non-preferred bisphosphonate. Low risk will be defined as having an osteopenic bone mineral density (most recent T-score between -1 and -2.5) AND no history of vertebral fracture.

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	DIDRONEL (etidronate) FOSAMAX (alendronate) FOSAMAX plus D (alendronate w/D) Etidronate Risedronate	
--	---	--

Non-Bisphosphonates		
	<p>PA Required</p> Calcitonin salmon (nasal) EVISTA (raloxifene) FORTEO (teriparatide) Raloxifene TYMLOS (abaloparatide)	<p>Calcitonin salmon (nasal) will be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> • Member has a diagnosis of post-menopausal osteoporosis (BMD T-scores of -2.5 or less) AND • Has trial and failure of preferred bisphosphonate for one year (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) OR • Member cannot swallow solid oral dosage forms or has a feeding tube. Quantity limit of one spray per day <p>Raloxifene will be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> • Diagnosis of postmenopausal osteoporosis (BMD T-scores of -2.5 or less) AND • Has trial and failure of preferred bisphosphonate for one year (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) <p>Maximum Dose of raloxifene is 60mg oral daily</p> <p>Forteo (teriparatide) will be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> • Member has one of the following diagnoses: • Osteoporosis, (BMD T-scores of -2.5 or less) primary or hypogonadal in men • Osteoporosis due to corticosteroid use • Postmenopausal osteoporosis AND • Has trial and failure of preferred bisphosphonate for one year (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) <ul style="list-style-type: none"> • Prior authorization will be given for one year and total exposure of parathyroid hormone analogs (Forteo and Tymlos) shall not exceed two years

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

		<p>Maximum dose of Forteo is 20mcg subcutaneous daily</p> <p>Tymlos (abaloparatide) will be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> • Member has a diagnosis of postmenopausal osteoporosis (BMD T-scores of -2.5 or less) AND • Has trial and failure of preferred bisphosphonate for one year (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND • Prior authorization will be given for one year and total exposure of parathyroid hormone analogs (Forteo and Tymlos) shall not exceed two years. <p>Maximum dose of Tymlos is 80 mcg injection daily</p> <p>Prolia (denosumab) is a physician administered drug and prior authorization criteria may be found on the Appendix P.</p>
--	--	--

Therapeutic Drug Class: CONTRACEPTIVES - Oral Effective 10/1/2019

No PA Required	PA Required	
<p>Monophasic 28:</p> <p>Altavera 28 0.15-30 Alyacen 28 1-35 Apri 28 0.15-30 Aubra EQ-28 0.1-20 Aviane 28 0.1-20 Balziva 28 0.4-35 Chateal 28 0.15-30 Chateal EQ 28 0.15-30 Cryselle 28 0.3-30</p> <p style="text-align: center;">No PA Required</p> <p>Cyclafem 28 1-35 Dasetta 28 1-35 Drosperinone-Eth Estradiol 28 3-30 Elinest 28 0.3-30 Enskyce 28 0.15-30 Estarylla 28 0.25-35 Ethinodiol-Eth Estra 28 1-35 Ethinodiol-Eth Estra 28 1-50 Falmina 28 0.1-20 Femynor 28 0.25-35</p>	<p>Levonor-Eth Estrad 28 0.15-30 Levora 28 0.15-30 Lillow 28 0.15-30 Low-Ogestrel 28 0.3-30 Lutera 28 0.1-20 Marlissa 28 0.15-30 Mili 28 0.25-35 Mono-Linyah 28 0.25-35 Mononessa 28 0.25-35 Norg-Ethin Estra 28 0.25-35</p> <p style="text-align: center;">No PA Required</p> <p>Nortrel 28 0.5-35 Nortrel 28 1-35 Ocella 28 3-30 Philith 28 0.4-35 Pirmella 28 1-35 Portia 28 0.15-30 Previfem 28 0.25-35 Reclipsen 28 0.15-30 Sprintec 28 0.25-35 Sronyx 28 0.1-20 Syeda 28 3-30</p>	<p>All other rebateable products are non-preferred</p> <p>Non-preferred oral contraceptive products will be approved if member fails one-month trial with four preferred agents OR if preferred products with medically necessary ingredients and/or doses are unavailable. (Failure is defined as: allergy, intolerable side effects, or significant drug-drug interaction)</p> <p>Initial fills may be dispensed for three-month supply to establish tolerance (i.e. lack of adverse effects). After established tolerance on the same agent for 3 months, a 12 month supply (365 days) may be dispensed (as one fill).</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)	
------------------	----------------------	---	--

No PA Required	No PA Required		
<p>Isibloom 28 0.15-30 Juleber 28 0.15-30 Kelnor 28 1-35 Kurvelo 28 0.15-30 Larissia 28 0.1-20 Lessina 28 0.1-20 Levonor-Eth Estrad 28 0.1-20</p> <p><u>Biphasic:</u> Azurette 28 Bekyree 28 Desogest-Eth Estra 28 Kariva 28 Lo Loestrin FE 28 1-10 Mircette 28 Viorele 28</p> <p><u>Triphasic:</u> Alyacen 7-7-7 28 Cyclafem 7-7-7 28 Dasetta 7-7-7 28 Enpresse 28 Levonest 28 Levonor-Eth Estrad Triphasic 28 Pirmella 7-7-7 Tri-Estarylla 28 Tri-Femynor 28 Tri-Linyah 28 Tri-Lo Estarylla 28 Tri-Lo Marzia 28 Tri-Lo Sprintec 28 Trinessa 28 Tri-Sprintec 28 Tri-Vylibra Lo 28</p>	<p>Vienva 28 0.1-20 Vyfemla 28 0.4-35</p> <p><u>Monophasic 21:</u> Larin 21 1-20 Larin 21 1.5-30 Norethind-Eth Estrad 21 1-20 Nortrel 21 1-35</p> <p><u>Extended Cycle:</u> Amethia 91 0.03 – 0.15 – 0.01 Ashlyna 91 0.15-10-30 Introvale 91 0.15-30 Jolessa 91 0.15-30 Levonorgest-Eth Estrad 0.09-20 Levonorgest-Eth Estrad 91 0.1-10-20 Levonorgest-Eth Estrad 91 0.15-0.03 Levonorgest-Eth Estrad 91 0.15-0.03-0.01 Levonorgest-Eth Estrad 91 0.15-20-25-30 Quasense 91 0.15-30 Setlakin 91 0.15-30</p>		

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

No PA Required	No PA Required		
<p><u>Norethindrone Only:</u> Camila 28 0.35 Deblitane 28 0.35 Errin 28 0.35 Heather 28 0.35 Jencycla 28 0.35 Jolivette 28 0.35 Norethindrone 28 0.35 Norlyda 28 0.35 Sharobel 28 0.35</p>	<p><u>Continuous Cycle:</u> Aurovela FE 1-20 Blisovi FE 1-20 Blisovi FE 1.5-30 Jasmiel 3-20 Junel FE 1-20 Junel FE 24 1-20 Junel FE 1.5-30 Larin FE 1-20 Larin FE 24 1-20 Larin FE 1.5-30 Loryna 3-20 Minastrin FE 24 1-20 Nikki 3-20 Noreth-Eth Estrad-FE 24 1-20 Noreth-Eth Estrad-FE 1-20 Tarina FE 24 1-20 Tarina FE 1-20 Tarina FE 1-20 EQ</p>		

Therapeutic Drug Class: **DIABETES MANAGEMENT CLASSES, INSULINS-** *Effective 4/1/2020*

Rapid-Acting		
No PA Required	PA Required	
NOVOLOG (insulin aspart) cartridge, vial, FlexTouch HUMALOG (insulin lispro) cartridge, vial, KwikPen, pen HUMALOG Jr. (insulin lispro) KwikPen	ADMELOG (insulin lispro) vial, Solostar AFREZZA (regular insulin) cartridge, unit APIDRA (insulin glulisine) vial, Solostar	<p>Non-preferred products may be approved following trial and failure of treatment with two preferred products (failure is defined as allergy [hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema] or intolerable side effects).</p> <p>Afrezza (human insulin) may be approved if meeting the following criteria:</p> <ul style="list-style-type: none"> • Member is 18 years or older AND • Member has trialed and failed treatment with two preferred products (failure is defined as allergy [hives, maculopapular rash, erythema multiforme, pustular rash, severe hypotension, bronchospasm, and angioedema] or intolerable side effects) AND • Member must not have chronic lung disease such as COPD or asthma AND • If member is a type 1 diabetic, must use in conjunction with long-acting insulin AND • Member must not be a smoker

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
	FIASP (insulin aspart) vial, FlexTouich, PenFill Insulin lispro pen, vial	
Short-Acting		
HUMULIN R (insulin regular) vial (OTC) HUMULIN R (insulin regular) concentrated vial, Kwikpen (U-500)	NOVOLIN R (insulin regular) vial (OTC) HUMULIN R (insulin regular) KwikPen (OTC)	Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).
Intermediate-Acting		
HUMULIN N (insulin NPH) vial, Kwikpen (OTC)	NOVOLIN N (insulin NPH) vial (OTC)	Non-preferred products may be approved following trial and failure of treatment with one preferred product (failure is defined as allergy or intolerable side effects).
Long-Acting		
LEVEMIR (insulin detemir) vial, FlexTouch LANTUS (insulin glargine) vial, Solostar	BASAGLAR (insulin glargine) KwikPen TOUJEO (insulin glargine) Solostar TOUJEO MAX (insulin glargine) Solostar TRESIBA (insulin degludec) vial, FlexTouch	Non-preferred products may be approved if the member has failed treatment with Levemir AND Lantus (failure is defined as allergy or intolerable side effects).
Mixtures		
HUMULIN 70/30 vial, Kwikpen (OTC) HUMALOG MIX 50/50 vial, Kwikpen	NOVOLIN 70/30 vial, FlexPen (OTC)	Non-preferred products may be approved if the member has failed treatment with two of the preferred products (failure is defined as: allergy or intolerable side effects).

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>HUMALOG MIX 75/25 vial, Kwikpen</p> <p>NOVOLOG MIX 70/30 vial, FlexPen</p>		
---	--	--

Therapeutic Drug Class: DIABETES MANAGEMENT CLASSES, NON- INSULINS- 10/1/2019

Amylin

	<p align="center">PA Required</p> <p>SYMLIN (pramlintide)</p>	<p>Symlin® will only be approved after a member has failed a three month trial of metformin and a DPP4-inhibitor or a GLP-1 analogue. Failure is defined as: lack of efficacy (e.g., hemoglobin A1C \geq 7%) OR the member cannot tolerate metformin, DPP4-inhibitor and GLP-1 analogue due to allergy, intolerable side effects, or a significant drug-drug interaction. PA will be approved for Symlin products for members with Diabetes Mellitus Type 1 without failed treatment</p> <p>For all products, dosing will be limited to FDA approved dosing. PA will be required for doses in excess of FDA approved dosing.</p>
--	--	---

Biguanides

<p align="center">No PA Required</p> <p>Metformin 500mg, 850mg, 1000mg tablets</p> <p>Metformin ER 500mg tablets (generic Glucophage XR)</p>	<p align="center">PA Required</p> <p>FORTAMET (metformin)</p> <p>GLUCOPHAGE (brand) (metformin)</p> <p>GLUCOPHAGE XR (brand) (metformin XR)</p> <p>GLUMETZA ER (metformin)</p> <p>Metformin ER 750mg</p> <p>Metformin ER 500 and 1000mg (generic Fortamet, generic Glumetza)</p> <p>RIOMET 500mg/5ml (metformin)</p>	<p>Non-preferred products will be approved for members who have failed treatment with two Preferred Products. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.)</p> <p>Liquid metformin will be approved for members who meet one of the following: under the age of 12 with a feeding tube who have difficulty swallowing</p>
---	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

Dipeptidyl Peptidase-4 Enzyme inhibitors (DPP-4is)														
<p>*Must meet eligibility criteria</p> <p>*Januvia (sitagliptin)</p> <p>*Tradjenta (linagliptin)</p>	<p style="text-align: center;">PA Required</p> <p>Alogliptin</p> <p>Nesina (alogliptin)</p> <p>Onglyza (saxagliptin)</p>	<p>*Approval for preferred products require a three month trial of (or documented contraindication to) metformin therapy prior to initiation of therapy.</p> <p>Non-preferred DPP-4 inhibitors will be approved after a member has failed a three month trial of metformin AND a three month trial of two preferred products. Failure is defined as lack of efficacy (e.g., hemoglobin A1C \geq 7%), allergy, intolerable side effects, or a significant drug-drug interaction.</p> <p>For all products, prior authorization will be required for dosing above the FDA approved maximum dosing listed in the following table:</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="text-align: center;">DPP4</th> <th style="text-align: center;">FDA Approved Max Dose (mg/day)</th> </tr> </thead> <tbody> <tr> <td>Alogliptin (generic Nesina)</td> <td style="text-align: center;">25 mg/day</td> </tr> <tr> <td>Januvia (sitagliptan)</td> <td style="text-align: center;">100 mg/day</td> </tr> <tr> <td>Nesina (alogliptan)</td> <td style="text-align: center;">25 mg/day</td> </tr> <tr> <td>Onglyza (saxagliptan)</td> <td style="text-align: center;">5 mg/day</td> </tr> <tr> <td>Tradjenta (linagliptan)</td> <td style="text-align: center;">5 mg/day</td> </tr> </tbody> </table>	DPP4	FDA Approved Max Dose (mg/day)	Alogliptin (generic Nesina)	25 mg/day	Januvia (sitagliptan)	100 mg/day	Nesina (alogliptan)	25 mg/day	Onglyza (saxagliptan)	5 mg/day	Tradjenta (linagliptan)	5 mg/day
DPP4	FDA Approved Max Dose (mg/day)													
Alogliptin (generic Nesina)	25 mg/day													
Januvia (sitagliptan)	100 mg/day													
Nesina (alogliptan)	25 mg/day													
Onglyza (saxagliptan)	5 mg/day													
Tradjenta (linagliptan)	5 mg/day													

DPP-4 Inhibitors – Combination with Metformin		
<p>*Must Meet eligibility criteria</p> <p>*JANUMET (sitagliptin/metformin)</p> <p>*JANUMET XR (sitagliptin/metformin)</p>	<p style="text-align: center;">PA Required</p> <p>Alogliptin/metformin</p> <p>JENTADUETO (linagliptin/metformin)</p> <p>JENTADUETO XR (linagliptin/metformin)</p> <p>KAZANO (alogliptin/metformin)</p>	<p>*Approval for preferred combination agent products require a three month trial of (or documented contraindication to) metformin therapy prior to initiation of therapy.</p> <p>Non-preferred combination products will be approved for members who have been stable on the two individual ingredients of the requested combination for three months AND have had adequate three-month trial and failure of a preferred combination agent. Failure is defined as lack of efficacy (e.g., hemoglobin A1C \geq 7%), allergy, intolerable side effects, or a significant drug-drug interaction.</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

	KOMBIGLYZE (saxagliptin/metformin)	
--	---------------------------------------	--

Glucagon-like Peptide-1 Receptor Agonists (GLP-1 Analogues)

<p>*Must meet eligibility criteria</p> <p>*BYETTA (exenatide)</p> <p>*BYDUREON (exenatide ER)</p> <p>*VICTOZA (liraglutide)</p>	<p>PA Required</p> <p>ADLYXIN (lixisenatide)</p> <p>BYDUREON BCISE (exenatide ER)</p> <p>OZEMPIC (semaglutide)</p> <p>TRULICITY (dulaglutide)</p>	<p>*Approval for preferred products requires a three month trial of (or documented contraindication to) metformin therapy prior to initiation of therapy.</p> <p>Non-preferred products may be approved following trial and failure of a three month trial of metformin AND a three month trial of two preferred products. Failure is defined as lack of efficacy (e.g., hemoglobin A1C \geq 7%), allergy, intolerable side effects, or a significant drug-drug interaction.</p> <p><u>Maximum Dose:</u> Prior authorization is required for all products exceeding maximum dose listed in product package labeling.</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th colspan="2" style="text-align: center;">Maximum Dose</th> </tr> </thead> <tbody> <tr> <td>Adlyxin (lixisenatide)</td> <td>20mcg per day</td> </tr> <tr> <td>Bydureon (exenatide)</td> <td>2mg weekly</td> </tr> <tr> <td>Bydureon BCISE (exenatide)</td> <td>2mg weekly</td> </tr> <tr> <td>Byetta (exenatide)</td> <td>20mcg per day</td> </tr> <tr> <td>Ozempic (semaglutide)</td> <td>1mg weekly</td> </tr> <tr> <td>Trulicity (dulaglutide)</td> <td>1.5mg weekly</td> </tr> <tr> <td>Victoza (liaglutide)</td> <td>1.8mg per day</td> </tr> </tbody> </table>	Maximum Dose		Adlyxin (lixisenatide)	20mcg per day	Bydureon (exenatide)	2mg weekly	Bydureon BCISE (exenatide)	2mg weekly	Byetta (exenatide)	20mcg per day	Ozempic (semaglutide)	1mg weekly	Trulicity (dulaglutide)	1.5mg weekly	Victoza (liaglutide)	1.8mg per day
Maximum Dose																		
Adlyxin (lixisenatide)	20mcg per day																	
Bydureon (exenatide)	2mg weekly																	
Bydureon BCISE (exenatide)	2mg weekly																	
Byetta (exenatide)	20mcg per day																	
Ozempic (semaglutide)	1mg weekly																	
Trulicity (dulaglutide)	1.5mg weekly																	
Victoza (liaglutide)	1.8mg per day																	

Other Hypoglycemic Combinations

	<p>PA Required</p> <p>Alogliptin/pioglitazone</p> <p>AVANDARYL (rosiglitazone/glimepiride)</p> <p>DUETACT (pioglitazone/glimepiride)</p> <p>Pioglitazone/glimepiride</p>	<p>Non-preferred products may be approved for members who have been stable on each of the individual ingredients in the requested combination for 3 months (including cases where the ingredients are taken as two separate 3 month trials or when taken in combination for at least 3 months).</p>
--	---	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	<p>Glipizide/metformin</p> <p>GLUCOVANCE (glyburide/metformin)</p> <p>Glyburide/metformin</p> <p>GLYXAMBI (empagliflozin/linagliptin)</p> <p>METAGLIP (glipizide/metformin)</p> <p>OSENI (alogliptin/pioglitazone)</p> <p>Qtern (dapagliflozin/saxagliptin)</p> <p>Soliqua (glargine 100 U and lixisenatide 33 mcg)</p> <p>Steglujan (ertugliflozin/sitagliptin)</p> <p>Xultophy (degludec 100 U and liraglutide 3.6 mg)</p>	
--	--	--

Meglitinides		
---------------------	--	--

	<p>PA Required</p> <p>Nateglinide</p> <p>PRANDIN (repaglinide)</p> <p>Repaglinide</p> <p>STARLIX (nateglinide)</p>	<p>Non-preferred products will be approved for members who have failed treatment with one Sulfonylurea (Failure is defined as: lack of efficacy (e.g., hemoglobin A1C \geq 7%), allergy, intolerable side effects, or significant drug-drug interaction.)</p>
--	---	--

Meglitinides Combination with Metformin		
--	--	--

	<p>PA Required</p>	
--	---------------------------	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	PRANDIMET (repaglinide/metformin) Repaglinide/metformin	Non-preferred products will be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months.
--	--	---

Sodium-Glucose Cotransporter 2 inhibitors (SGLT-2is)		
--	--	--

<p>*Must meet eligibility criteria</p> <p>*FARXIGA (dapagliflozin)</p> <p>*INVOKANA (canagliflozin)</p> <p>*JARDIANCE (empagliflozin)</p>	<p>PA Required</p> <p>STEGLATRO (ertugliflozin)</p>	<p>*Approval for preferred products requires a three month trial of (or documented contraindication to) metformin therapy prior to initiation of therapy.</p> <p>Non-preferred products may receive approval following trial and failure with a three month trial of metformin AND a three month trial of two preferred products. Failure is defined as lack of efficacy with three month trial (e.g., hemoglobin A1C \geq 7%) allergy, intolerable side effects, or a significant drug-drug interaction</p> <p><u>Maximum Dose:</u> Prior authorization is required for all products exceeding maximum dose listed in product package labeling.</p>
--	--	---

SGLT-2 Inhibitors Combination with Metformin		
--	--	--

	<p>PA Required</p> <p>INVOKAMET (canagliflozin/metformin)</p> <p>SEGLUROMET (ertugliflozin/metformin)</p> <p>SYNJARDY (empagliflozin/metformin)</p> <p>XIGDUO XR (dapagliflozin/metformin)</p>	<p>Non-preferred products will be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months.</p>
--	---	--

Thiazolidinediones (TZDs)		
---------------------------	--	--

<p>No PA Required</p> <p>Pioglitazone</p>	<p>PA Required</p> <p>ACTOS (pioglitazone)</p>	<p>Non-preferred TZDs will be approved after a member has failed a three month trial of metformin and failed a three month trial of a preferred product. Failure is defined as lack of efficacy (e.g., hemoglobin A1C \geq 7%), OR the member cannot tolerate pioglitazone and metformin due to allergy, intolerable side effects, or a significant drug-drug interaction.</p>
--	---	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	AVANDIA (rosiglitazone)	
--	-------------------------	--

Thiazolidinediones Combination with Metformin		
--	--	--

	<p style="text-align: center;">PA Required</p> <p>ACTOPLUS MET (pioglitazone/metformin)</p> <p>ACTOPLUS MET XR (pioglitazone/metformin)</p> <p>AVANDAMET (rosiglitazone/metformin)</p> <p>Pioglitazone/metformin</p>	<p>Non-preferred products will be approved for members who have been stable on the two individual ingredients of the requested combination for 3 months.</p>
--	---	--

Therapeutic Drug Class: GLUCAGON, SELF-ADMINISTERED -Effective 4/1/2020		
--	--	--

<p style="text-align: center;">No PA Required (*Must meet eligibility criteria)</p> <p>GLUCAGEN HYPOKIT (glucagon)</p> <p>Glucagon Emergency Kit</p> <p>GVOKE (glucagon)*</p>	<p style="text-align: center;">PA Required</p> <p>BAQSIMI (glucagon) Nasal Spray</p>	<p>*Gvoke (glucagon) may be approved following trial and failure of GlucaGen (glucagon) OR glucagon emergency kit (failure is defined as allergy to ingredients in product, intolerable side effects, or inability to administer dosage form).</p> <p>Non-preferred products may be approved if the member has failed treatment with Gvoke (glucagon) AND one other preferred product (failure is defined as allergy to ingredients in product, intolerable side effects, or contraindication to dosing form).</p> <p>Quantity limit: 2 doses per year unless used / damaged / lost</p>
--	---	--

Therapeutic Drug Class: GROWTH HORMONES -Effective 4/1/2020		
--	--	--

<p style="text-align: center;">No PA Required (if diagnosis and dose met)</p> <p>GENOTROPIN</p> <p>NORDITROPIN</p>	<p style="text-align: center;">PA Required</p> <p>HUMATROPE</p> <p>NUTROPIN AQ</p> <p>OMNITROPE</p>	<p>All preferred products may be approved if the member has one of the qualifying diagnoses listed below (diagnosis may be verified through AutoPA) AND if prescription does not exceed limitations for maximum dosing (Table 1).</p> <p>Non-preferred Growth Hormones may be approved if the following criteria are met:</p> <ul style="list-style-type: none"> • Member failed treatment with one preferred growth hormone product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). • Member has a qualifying diagnosis:
---	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>SAIZEN</p> <p>SEROSTIM</p> <p>ZOMACTON</p> <p>ZORBATIVE</p>	<ul style="list-style-type: none"> ○ Prader-Willi ○ Chronic renal insufficiency/failure requiring transplantation (defined as Creatinine Clearance < 30mL/min) ○ Turner’s Syndrome ○ Hypopituitarism: as a result of pituitary disease, hypothalamic disease, surgery, radiation therapy or trauma verified by one of the following: <ul style="list-style-type: none"> ▪ Has failed at least one GH stimulation test (peak GH level < 10 ng/mL) ▪ Has at least one documented low IGF-1 level (below normal range for patient’s age – refer to range on submitted lab document) ▪ Has deficiencies in ≥ 3 pituitary axes (i.e. TSH, LH, FSH, ACTH, ADH) ○ Cachexia associated with AIDS ○ Noonan Syndrome ○ Short bowel syndrome ○ Neonatal symptomatic growth hormone deficiency (limited to three month PA approval) <ul style="list-style-type: none"> ● Prescription does not exceed limitations for maximum dosing (Table 1) based on prescriber submission/verification of patient weight from most recent clinical documentation
--	---

Table 1: Growth Hormone Product Maximum Dosing*		
Medication	Pediatric Max Dosing (age < 18 years)	Adult Max Dosing (age ≥ 18 years)
Genotropin	0.33 mg/kg/week	0.08 mg/kg/week
Humatrope	0.375 mg/kg/week	0.0875 mg/kg/week
Norditropin Flexpro	0.47 mg/kg/week	0.112 mg/kg/week
Nutropin AQ Nuspin	0.357 mg/kg/week	0.175 mg/kg/week for ≤35 years of age 0.0875 mg/kg/week for >35 years of age
Omnitrope	0.33 mg/kg/week	0.08 mg/kg/week
Saizen	0.18 mg/kg/week	0.07 mg/kg/week
Serostim	Not Indicated	42 mg/week for cachexia with HIV only (in combination with antiretroviral therapy)
Zomacton	0.375 mg/kg/week	0.0875 mg/kg/week
Zorbative	Not Indicated	8 mg/28 days for short bowel syndrome only

*Based on FDA labeled indications and dosing

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

VII. Gastrointestinal

Therapeutic Drug Class: **ANTI-EMETICS** -Effective 1/1/2020

No PA Required	PA Required	
Ondansetron ODT, tablet	AKYNZEO (netupitant/palonosetron) capsule	<p>Non-preferred products may be approved for members who have trialed and failed treatment with one preferred product AND one other anti-emetic (for example: prochlorperazine, metoclopramide, promethazine). Failure is defined as lack of efficacy with 14 day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>*Ondansetron solution may be approved for members < 5 years and those members ≥ 5 years of age with a feeding tube.</p> <p>Pyridoxine tablet AND doxylamine tablet may be approved for members who have a diagnosis of nausea and vomiting of pregnancy (NVP). Approval will be given for 9 months.</p> <p>Emend (aprepitant) TriPack or Emend (aprepitant) powder kit prior authorization may be approved for members who have trialed and failed one preferred product AND one other anti-emetic (for example: prochlorperazine, metoclopramide, promethazine) AND Emend (aprepitant) <u>capsule</u>. Failure is defined as lack of efficacy with 14 day trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Diclegis (doxylamine/pyridoxine) DR tablet or Bonjesta (doxylamine/pyridoxine) ER tablet may be approved for 9 months for members who meet the following criteria:</p> <ul style="list-style-type: none"> ● Has nausea and vomiting associated with pregnancy AND ● Has failed* 7-day trial of OTC formulation of pyridoxine (Vitamin B6) at maximally tolerated dose of up to 200mg daily AND ● Has failed* 7-day combination trial of OTC formulations of doxylamine and pyridoxine (Vitamin B6) at maximum daily doses of doxylamine 40mg and pyridoxine 40mg AND ● Has failed* 7 day trial of alternate antihistamine (diphenhydramine, dimenhydrinate, meclizine) OR ● Has failed* 7 day trial of dopamine antagonist (metoclopramide, prochlorperazine, promethazine) OR ● Has failed 7-day trial of serotonin antagonist (ondansetron, granisetron). *Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. <p>Dronabinol prior authorization may be approved for members meeting above non-preferred criteria. OR via AutoPA for members with documented HIV diagnosis.</p>
Ondansetron oral solution* (members under 5 years)	ANZEMET (dolasetron) tablet	
Transderm Scop (scopolamine) <small>BNR</small>	Aprepitant capsule	
	BONJESTA ER (doxylamine/pyridoxine) tablet	
	DICLEGIS DR (doxylamine/pyridoxine) tablet	
	Doxylamine 25mg (OTC)	
	Doxylamine/pyridoxine tablet (generic Diclegis)	
	Dronabinol capsule	
	EMEND (aprepitant) capsule, powder for suspension, dose/tri pack	
	Granisetron tablet	
	MARINOL (dronabinol) capsule	
	Pyridoxine 50mg or 100mg (OTC)	
	SANCUSO (granisetron) patch	
	Scopolamine patch	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	VARUBI (rolapitant) tablet ZOFRAN (ondansetron) tabs ZUPLLENZ (ondansetron)	
--	---	--

Therapeutic Drug Class: BILE SALTS -Effective 4/1/2020

No PA Required	PA Required	
Ursodiol capsule Ursodiol tablet	ACTIGALL (ursodiol) capsule CHENODAL (chenodiol) tablet CHOLBAM (cholic acid) capsule OCALIVA (obeticholic acid) tablet URSO (ursodiol) tablet URSO FORTE (ursodiol) tablet	<p>Chenodal (chenodiol) and Actigall (ursodiol) may be approved for members who meet the following criteria:</p> <ul style="list-style-type: none"> • Member \geq 18 years of age AND • Member has tried and failed therapy with a 12 month trial of a preferred ursodiol (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions). <p>Cholbam (cholic acid) may be approved for members who meet the following criteria:</p> <ul style="list-style-type: none"> • Bile acid synthesis disorders: <ul style="list-style-type: none"> ○ Member must be greater than 3 weeks old in age AND ○ Member has a diagnosis for bile acid synthesis disorder due to single enzyme defect (Single Enzyme-Defect Disorders: Defective sterol nucleus synthesis, 3β-hydroxy-Δ-c27-steroid oxidoreductase deficiency, AKR1D1 deficiency, CYP7A1 deficiency, Defective side-chain synthesis, CYP27A1 deficiency (cerebrotendinous xanthomatosis), 2-methylacyl-CoA racemase deficiency (AMACR), 25-hydroxylation pathway (Smith-Lemli-Opitz). • Peroxisomal disorder including Zellweger spectrum disorders: <ul style="list-style-type: none"> ○ Member must be greater than 3 weeks old in age AND ○ Member has diagnosis of peroxisomal disorders (PDs) including Zellweger spectrum disorders AND ○ Member has manifestations of liver disease, steatorrhea or complications from decreased fat-soluble vitamin absorption. <p>Ocaliva (obeticholic acid), Urso (ursodiol), and Urso Forte (ursodiol) may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none"> • Member is \geq 18 years of age AND • Medication is prescribed by or in consultation with a gastroenterologist, hepatologist, or liver transplant provider AND

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<ul style="list-style-type: none"> • Member has the diagnosis of Primary Biliary Cholangitis as evidenced by two of the following at the time of diagnosis: <ul style="list-style-type: none"> ○ Evidence of cholestasis with an alkaline phosphatase elevation of at least 1.5 times the upper limit of normal ○ Presence of antimitochondrial antibody: a titer of 1:40 or higher ○ Histologic evidence of nonsuppurative destruction cholangitis and destruction of interlobular bile ducts AND • Member has failed treatment with a preferred ursodiol product for at least 1 year with an inadequate response OR • Member has intolerable side effects, drug-drug interaction, or allergy to preferred ursodiol formulation.
--	--	--

Therapeutic Drug Class: GI MOTILITY, CHRONIC -Effective 10/1/2019

PA Required for all agents in this class		All GI Motility Agents will only be approved for FDA labeled indications and up to FDA approved maximum doses (listed below):
AMITIZA (lubiprostone) LINZESS (linaclotide) MOVANTIK (naloxegol)	Alosetron LOTRONEX (Alosetron) MOTEGRITY (prucalopride) RELISTOR (Methylnaltrexone bromide) tablet and syringe SYMPROIC (Naldemedine) TRULANCE (plecanatide) VIBERZI (eluxadoline)	<p>Preferred agents will be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> • Has diagnosis of Irritable Bowel Syndrome – Constipation (IBS-C), Chronic Idiopathic Constipation (CIC), or Opioid Induced Constipation (OIC) in patients with opioids prescribed for noncancer pain AND • Member does not have a diagnosis of GI obstruction AND • For indication of OIC, member opioid use must exceed 4 weeks of treatment • For indications of CIC, OIC, IBS-C; member must have documentation of adequate trial of two or more over-the-counter motility agents (for example; polyethylene glycol, docusate, bisocodyl) (Failure is defined as a lack of efficacy for a 7 day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions) <ul style="list-style-type: none"> ○ If the member cannot take oral medications, then the member must fail a 7-day trial with a nonphosphate enema (docusate or bisocodyl enema) • For indication of IBS-D; must have documentation of adequate trial with loperamide AND dicyclomine OR hyoscamine (Failure is defined as a lack of efficacy for a 7 day trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions) <p>Non-preferred agents may be approved if the member meets the following criteria:</p> <ul style="list-style-type: none"> • Member meets all listed criteria for preferred agents AND • Member has trialed and failed two preferred agents <ul style="list-style-type: none"> ○ If indication OIC caused by methadone, then non-preferred agent may be approved after trial of Movantik (Failure is defined as a lack of efficacy for a 7 day trial,

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<p>allergy, intolerable side effects, contraindication to, or significant drug-drug interactions) AND</p> <ul style="list-style-type: none"> ○ Member meets additional criteria for the agents listed below <p>Viberzi® (eluxadoline) will be approved for members who meet the following criteria:</p> <ul style="list-style-type: none"> ● Has diagnosis of Irritable Bowel Syndrome – Diarrhea (IBS-D) AND ● Member has a gallbladder AND ● Member does not have severe hepatic impairment (Child-Pugh C), history of severe constipation, known mechanical gastrointestinal obstruction, biliary duct obstruction, history of pancreatitis or structural disease of the pancreas AND ● Member does not drink more than 3 alcoholic drinks per day AND <p>Lotronex® (alesotron) and Alesotron will be approved for members who meet the following criteria:</p> <ul style="list-style-type: none"> ● Member is a female with Irritable Bowel Syndrome – Diarrhea (IBS-D) with symptoms lasting 6 months or longer AND ● Member does not have severe hepatic impairment (Child-Pugh C), history of severe constipation or ischemic colitis, hypercoagulable state, Crohn’s disease or ulcerative colitis, or known mechanical gastrointestinal obstruction
--	--	---

Medication	FDA approved indication	FDA Max Dose
Amitiza (lubiprostone)	IBS-C (females only), CIC, OIC (not caused by methadone)	48mcg/day
Linzess (linaclotide)	IBS-C, CIC	290mcg/day
Movantik (naloxegol)	OIC	25mg/day
Viberzi (eluxadoline)	IBS-D	200mg/day
Alosetron	OIC	2mg/day (females only)
Relistor syringe (methylnaltrexone)	OIC	12mg SQ/day
Relistor oral (methylnaltrexone)	OIC	450mg/day
Lotronex (alosetron)	IBS-D (females only)	2mg/day (females only)
Symproic (Naldemedine)	OIC	0.2mg/day
Trulance (plecanatide)	CIC, IBS-C	3mg/day

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Motegrity (prucalopride)	CIC	2mg/day
--------------------------	-----	---------

CIC – chronic idiopathic constipation, OIC – opioid induced constipation, IBS – irritable bowel syndrome, D – diarrhea predominant, C – constipation predominant

Therapeutic Drug Class: HEMORRHOIDAL AND RELATED ANORECTAL AGENTS - Effective 4/1/2020

No PA Required	PA Required	
CORTIFOAM (hydrocortisone) aerosol	ANA-LEX (hydrocortisone-lidocaine)	<p>Non-preferred products may be approved following trial and failure of therapy with 3 preferred products (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Rectiv (nitroglycerin) ointment may be approved if meeting the following:</p> <ul style="list-style-type: none"> • Member has a diagnosis of anal fissure AND • Prescriber attests that member has trialed and maximized use of appropriate supportive therapies including sitz bath, fiber, topical analgesics (such as lidocaine), and stool softeners/laxatives.
Hydrocortisone enema	ANALPRAM HC (hydrocortisone-pramoxine) cream	
Hydrocortisone 25 mg suppository	ANUCORT-HC (hydrocortisone) suppository	
Hydrocortisone 2.5% cream with applicator	ANUSOL-HC (hydrocortisone) suppository, cream	
Hydrocortisone-Pramoxine 1%-1%, 2.5%-1% cream	COLOCORT (hydrocortisone) enema	
Lidocaine-Hydrocortisone 3-0.5% cream	CORTENEMA (hydrocortisone) enema	
PROCTOFOAM (hydrocortisone-pramoxine)	Hydrocortisone 30 mg suppository, 1% cream with applicator	
PROCTO-MED HC (hydrocortisone) 2.5% cream	Lidocaine-Hydrocortisone 3-0.5% cream kit	
PROCTO-PAK (hydrocortisone) 1% cream	Lidocaine-Hydrocortisone 3-2.5% gel	
PROCTOSOL-HC 2.5% (hydrocortisone) cream	MICORT-HC (hydrocortisone) cream	
	PROCORT (hydrocortisone-pramoxine) cream	
	PROCTOCORT (hydrocortisone) suppository	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

PROCTOZONE-HC 2.5% (hydrocortisone) cream	RECTIV (nitroglycerin) ointment	
--	---------------------------------	--

Therapeutic Drug Class: PANCREATIC ENZYMES -Effective 1/1/2020

No PA Required	PA Required	
CREON (pancrelipase) capsule ZENPEP (pancrelipase) capsule	PANCREAZE (pancrelipase) capsule PERTZYE (pancrelipase) capsule VIOKACE (pancrelipase) tablet	Non-preferred products will be approved for members who have failed an adequate trial (4 weeks) with at least two preferred products. (Failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interaction.) Grandfathering: Members currently stabilized on a Non-preferred pancreatic enzyme can receive approval to continue on that agent for one year if medically necessary.

Therapeutic Drug Class: PROTON PUMP INHIBITORS -Effective 1/1/2020

No PA Required	PA Required	
Esomeprazole capsule (generic Nexium) RX Lansoprazole capsules (generic Prevacid) RX NEXIUM (esomeprazole) packets Omeprazole capsule Pantoprazole tablet PREVACID Solutab ^{BNR} (lansoprazole) (members < 2)	ACIPHEX (rabeprazole) tablet, sprinkle capsule DEXILANT (dexlansoprazole) capsule Esomeprazole strontium DR capsule Esomeprazole mag capsule OTC Lansoprazole capsule OTC, ODT RX NEXIUM (esomeprazole) capsule (RX) Omeprazole/Na bicarbonate capsule, packet Omeprazole 20mg tablet, ODT (OTC) PREVACID (lansoprazole) capsule PRILOSEC (omeprazole) suspension	For members treating GERD symptoms that are controlled on PPI therapy, it is recommended that the dose of the PPI be re-evaluated or step-down with an H2 blocker (such as famotidine or ranitidine) be trialed in order to reduce long-term PPI use. Prior authorization for non-preferred proton pump inhibitors may be approved if all of the following criteria are met: <ul style="list-style-type: none"> ● Member has a qualifying diagnosis (below) AND ● Member has trailed and failed therapy with three preferred agents within the last 24 months. (Failure is defined as: lack of efficacy following 4 week trial, allergy, intolerable side effects, or significant drug-drug interaction) AND ● Member has been diagnosed using one of the following diagnostic methods: <ul style="list-style-type: none"> ○ Diagnosis made by GI specialist ○ Endoscopy ○ X-ray ○ Biopsy ○ Blood test ○ Breath Test <p>Qualifying Diagnoses: Barrett's esophagus, duodenal ulcer, erosive esophagitis, gastric ulcer, GERD, GI Bleed, H. pylori infection, hypersecretory conditions (Zollinger-Ellison), NSAID-induced ulcer, pediatric esophagitis, requiring mechanical ventilation, requiring a feeding tube</p> <p>Quantity Limits:</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

	<p>PROTONIX (pantoprazole) tablet, suspension</p> <p>Rabeprazole (generic Aciphex) tablet</p> <p>ZEGERID (omeprazole/Na bicarbonate) capsule, packet</p>	<p>All agents will be limited to once daily dosing except when used for the following diagnoses: Barrett’s esophagus, GI Bleed, H. pylori, hypersecretory conditions (Zollinger-Ellison), or Spinal Cord Injury patients with associated acid reflux.</p> <p>Adult members with GERD on once daily, high-dose PPI therapy who continue to experience symptoms may receive initial prior authorization approval for a 4-week trial of twice daily, high-dose PPI therapy. Continuation of the twice daily dosing regimen for GERD beyond 4 weeks will require additional prior authorization approval verifying adequate member response to the dosing regimen and approval may be placed for one year. If a member with symptomatic GERD does not respond to twice daily, high-dose PPI therapy, this should be considered a treatment failure.</p> <p>Pediatric members (< 18 years of age) on once daily dosing of a PPI who continue to experience symptoms may receive one-year prior authorization approval for twice daily PPI therapy.</p> <p>Age Limits: Nexium 24H and Zegerid will not be approved for members less than 18 years of age.</p> <p>Prevacid Solutab will be approved for members < 2 years of age OR for members ≥ 2 years of age with a feeding tube.</p>
--	--	---

Therapeutic Drug Class: H. PYLORI TREATMENTS -Effective 1/1/2020

	<p style="text-align: center;">PA Required</p> <p>OMECLAMOX-PAK (amoxicillin/omeprazole/clarithromycin)</p> <p>PREVPAC (amoxicillin/lansoprazole/clarithromycin)</p> <p>Amoxicillin/lansoprazole/clarithromycin</p> <p>PYLERA (bismuth subcitrate/metronidazole/tetracycline)</p> <p>TALICIA (omeprazole/amoxicillin/rifabutin)</p>	<p>H. Pylori treatments should be used as individual products unless one of the individual products is not commercially available then a PA for the combination product will be given.</p>
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

Therapeutic Drug Class: ULCERATIVE COLITIS AGENTS- Oral -Effective 1/1/2020

No PA Required	PA Required	
<p>APRISO ER ^{BNR} (mesalamine) capsule</p> <p>LIALDA (mesalamine DR) ^{BNR} tablet</p> <p>PENTASA (mesalamine) capsule</p> <p>Sulfasalazine IR and DR tablet</p>	<p>Asacol HD (mesalamine) tablet</p> <p>AZULFIDINE (sulfasalazine) Entab, tablet</p> <p>Balsalazide disodium capsule</p> <p>Budesonide DR tablet</p> <p>COLAZAL (balsalazide) capsule</p> <p>DELZICOL DR (mesalamine) capsule</p> <p>DIPENTUM (olsalazine) capsule</p> <p>GIAZO (balsalazide) tablet</p> <p>Mesalamine DR (generic Asacol HD, Lialda) tablet</p> <p>Mesalamine capsule (generic Apriso ER)</p> <p>UCERIS (budesonide) tablet</p>	<p>Prior authorization for non-preferred oral formulations will require trial and failure of two preferred oral products with different active ingredients AND a preferred rectal product. If inflammation is not within reach of topical therapy, trial of preferred rectal product is not required. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Uceris (budesonide) tablet: If the above criteria is met, Uceris (budesonide) tablet prior authorization will be approved for 8 weeks. Further prior authorization may be approved if 7 days of steroid-free time has elapsed and member continues to meet the above criteria.</p>

Therapeutic Drug Class: ULCERATIVE COLITIS AGENTS- Rectal -Effective 1/1/2020

No PA Required	PA Required	
<p>Mesalamine suppository (generic Canasa)</p>	<p>CANASA (mesalamine) suppository</p> <p>Mesalamine enema, kit</p> <p>SF ROWASA (mesalamine)</p>	<p>Prior authorization for non-preferred rectal formulations will require trial and failure of one preferred rectal formulation and one preferred oral formulation (Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction).</p> <p>Uceris (budesonide) foam: If the above criteria is met, Uceris (budesonide) foam prior authorization will be approved for 6 weeks. Further prior authorization may be approved if 7 days of steroid-free time has elapsed and member continues to meet the above criteria.</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

	ROWASA (mesalamine w/cleansing wipes) UCERIS (budesonide) foam	
--	---	--

VIII. Hematological

Therapeutic Drug Class: ANTI-COAGULANTS- Oral -Effective 10/1/2019

No PA Required	PA Required	
Warfarin	BEVYXXA (betrixaban)	<p>Bevyxxa (betrixaban) may be approved if all the following criteria have been met:</p> <ul style="list-style-type: none"> ● The member has trialed and failed therapy with two preferred agents. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND ● Member is not on dialysis AND ● The member is need of prophylaxis for DVT following hospitalization for an acute medical illness who are at risk for thromboembolic events due to limited mobility AND ● The member does not have a mechanical prosthetic heart valve <p>Eliquis (apixaban) may be approved if the following criteria have been met:</p> <ul style="list-style-type: none"> ● The member is on dialysis OR ● The member has failed therapy with two preferred agents. Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. If the member is on dialysis, trial and failure of preferred agents is not required AND ● The member has a diagnosis of deep vein thrombosis (DVT), pulmonary embolism (PE) OR ● The member is in need of prophylaxis for DVT following knee or hip replacement surgery OR ● The member has a diagnosis of non-valvular atrial fibrillation AND ● The member does not have a mechanical prosthetic heart valve <p>Savaysa (edoxaban) may be approved if all the following criteria have been met:</p> <ul style="list-style-type: none"> ● The member has failed therapy with two preferred agents. (Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction) AND ● Member is not on dialysis AND ● Member does not have CrCl > 95 mL/min AND ● The member has a diagnosis of deep vein thrombosis (DVT), pulmonary embolism (PE) OR ● The member has a diagnosis of non-valvular atrial fibrillation AND ● The member does not have a mechanical prosthetic heart valve <p>Xarelto 2.5mg (rivaroxaban) may be approved for members meeting all of the following criteria:</p>
PRADAXA (dabigatran)	COUMADIN (warfarin)	
XARELTO (rivaroxaban) 10 mg, 15 mg, 20 mg tablet	ELIQUIS (apixaban)	
XARELTO (rivaroxaban) dose pack	SAVAYSA (edoxaban)	
	XARELTO (rivaroxaban) 2.5 mg tablet	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

		<ul style="list-style-type: none"> • Xarelto 2.5mg is being prescribed to reduce major CV events in members diagnosis of chronic coronary artery disease (CAD) or peripheral artery disease AND • Xarelto 2.5mg is being taken twice daily and in combination with aspirin 75-100mg daily AND • Member must not be receiving dual antiplatelet therapy, other non-aspirin antiplatelet, or other oral anticoagulant AND • Member must not have had an ischemic, non-lacunar stroke within the past month AND • Member must not have had a hemorrhagic or lacunar stroke at any time <p>Continuation of Care: Members with current prior authorization approval on file for a non-preferred <u>oral</u> anticoagulant medication may continue to receive approval for that medication.</p>
--	--	--

Therapeutic Drug Class: ANTI-COAGULANTS- Parenteral -Effective 10/1/2019

No PA Required	PA Required	
Enoxaparin syringe Lovenox 300mg/3ml vial ^{BNR}	Arixtra (fondaparinux) syringe Enoxaparin 300mg/3ml vial (generic Lovenox) Fondaparinux (generic Arixtra) Fragmin (dalteparin) vial and syringe Lovenox syringe	Non-preferred parenteral anticoagulants will be approved if member has trial and failure of one preferred agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction ARIXTRA® (fondaparinux) will be approved if the following criteria have been met: <ul style="list-style-type: none"> • Member is 18 years of age or older AND • Member has a CrCl > 30 ml/min AND • Member weighs > 50 kg AND • Member has a documented history of heparin induced-thrombocytopenia OR • Member has a contraindication to enoxaparin <p>Grandfathering: Members currently stabilized on fondaparinux (Arixtra) and dalteparin (Fragmin) may receive prior authorization approval to continue on that medication.</p>

Therapeutic Drug Class: ANTI-PLATELETS -Effective 1/1/2020

No PA Required	PA Required	
<p align="center"><i>Brand/generic changes effective 05/29/20</i></p> AGGRENOX (ASA/dipyridamole) capsule ASA/dipyridamole ER capsule	EFFIENT (prasugrel) tablet PLAVIX (clopidogrel) tablet PLETAL (cilostazol) Ticlopidine tablet	Patients taking Brilinta (ticagrelor) must also be taking a maintenance dose of aspirin not exceeding 100 mg/day. Ticlopidine should only be considered for patients who can be monitored for neutropenia and thrombocytopenia during the first three months of therapy. Zontivity (vorapaxar) will be approved for patients with a diagnosis of myocardial infarction or peripheral artery disease without a history of stroke, transient ischemic attack, intracranial bleeding,

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

BRILINTA (tigacrelor) tablet Cilostazol tablet Clopidogrel tablet Dipyridamole tablet Pentoxifylline ER tablet Prasugrel tablet	ZONTIVITY (vorapaxar) tablet	or active pathological bleeding. Patients must also be taking aspirin and/or clopidogrel concomitantly. Non-preferred products without criteria will be reviewed on a case by case basis.
--	------------------------------	--

Therapeutic Drug Class: COLONY STIMULATING FACTORS -Effective 10/1/2019

PA Required for all agents in this class		Prior authorization may be approved if meeting the following criteria:
NEUPOGEN (filgrastim) vial, syringe	FULPHILA (pegfilgrastim-jmdb) GRANIX (tbo-filgrastim) LEUKINE (sargramostim) NEULASTA (pegfilgrastim) syringe NIVESYM (filgrastim-aafi) UDENYCA (pegfilgrastim-cbqv) ZARXIO (filgrastim-sndz)	<ul style="list-style-type: none"> ● Medication is being used for one of the following indications: <ul style="list-style-type: none"> ○ Cancer patient receiving myelosuppressive chemotherapy –to reduce incidence of infection (febrile neutropenia) (Either the post nadir ANC is less than 10,000 cells/mm³ or the risk of neutropenia for the member is calculated to be greater than 20%) ○ Acute Myeloid Leukemia (AML) patients receiving chemotherapy ○ Bone Marrow Transplant (BMT) ○ Peripheral Blood Progenitor Cell Collection and Therapy ○ Hematopoietic Syndrome of Acute Radiation Syndrome ○ Severe Chronic Neutropenia (Evidence of neutropenia Infection exists or ANC is below 750 cells/mm³) <p>AND</p> <ul style="list-style-type: none"> ● All non-preferred agents will require a documented failure of Neupogen vial or syringe for approval (Failure is defined as a lack of efficacy with a 3 month trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions) AND ● For long-acting formulations (such as Fulphila and Neulasta), the member has trialed and failed a three month trial of Udenyca. Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions)

Therapeutic Drug Class: ERYTHROPOIESIS STIMULATING AGENTS Effective 10/1/2019

PA Required for all agents in this class*		*Prior Authorization is required for all products and may be approved if meeting the following:
		<ul style="list-style-type: none"> ● Medication is being administered in the member’s home or in a long-term care facility AND

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>RETACRIT (epoetin alfa-epbx)</p>	<p>ARANESP (darbepoetin alfa)</p> <p>EPOGEN (epoetin alfa)</p> <p>MIRCERA (methoxy peg-epoetin beta)</p> <p>PROCRIT (epoetin alfa)</p>	<ul style="list-style-type: none"> • Members meets one of the following: <ul style="list-style-type: none"> ○ A diagnosis of cancer, currently receiving chemotherapy, with chemotherapy-induced anemia, and hemoglobin[†] of 10g/dL or lower OR ○ A diagnosis of chronic renal failure, and hemoglobin[†] below 10g/dL OR ○ A diagnosis of hepatitis C, currently taking Ribavirin and failed response to a reduction of Ribavirin dose, and hemoglobin[†] less than 10g/dL (or less than 11g/dL if symptomatic). OR ○ A diagnosis of HIV, currently taking Zidovudine, hemoglobin[†] less than 10g/dL, and serum erythropoietin level of 500mUnits/mL or less OR ○ Member is undergoing elective, noncardiac, nonvascular surgery and medication is given to reduce receipt of allogenic red blood cell transfusions, hemoglobin[†] is greater than 10g/dL, but less than or equal to 13g/dL and high risk for perioperative blood loss. Member is not willing or unable to donate autologous blood pre-operatively. <p>AND</p> <ul style="list-style-type: none"> • For any non-preferred product, member has trialed and failed treatment with one preferred product. Failure is defined as lack of efficacy with a 6 week trial, allergy, intolerable side effects, or significant drug-drug interaction. <p>[†]Hemoglobin results must be from the last 30 days.</p>
-------------------------------------	--	---

IX. Immunological

Therapeutic Drug Class: **IMMUNE GLOBULINS** -Effective 4/1/2020

PA Required for all agents in this class*		
<p>CUVITRU 20% SQ liquid</p> <p>GAMMAGARD 10% IV/SQ liquid</p> <p>GAMMAKED 10% IV/SQ liquid</p> <p>GAMMAPLEX 5%, 10% IV liquid</p>	<p>BIVIGAM 10% IV liquid</p> <p>CUTAQUIG 16.5% SQ liquid</p> <p>FLEBOGAMMA DIF 5%, 10% IV liquid</p> <p>GAMMAGARD S-D solution</p> <p>HYQVIA 10% SQ liquid</p>	<p>Preferred agents may be approved for members meeting at least one of the approved conditions listed below for prescribed doses not exceeding maximum (Table 1).</p> <p>Non-preferred agents may be approved for members meeting the following:</p> <ul style="list-style-type: none"> • Member meets at least one of the approved conditions listed below AND • Member has history of trial and failure of two preferred agents (failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions) AND • Prescribed dose does not exceed listed maximum (Table 1) <p><u>Approved Conditions for Immune Globulin Use:</u></p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>GAMUNEX-C 10% IV/SQ liquid</p> <p>HIZENTRA 20% SQ liquid</p> <p>PRIVIGEN 10% IV liquid</p> <p><i>If immune globulin is being administered in a long-term care facility or in a member's home by a home healthcare provider, it should be billed as a pharmacy claim. All other claims must be submitted through the medical benefit.</i></p>	<p>OCTAGAM 5%, 10% IV liquid</p> <p>PANZYGA 10% IV liquid</p> <p>XEMBIFY 20% IV liquid</p>	<ul style="list-style-type: none"> • Primary Humoral Immunodeficiency disorders including: <ul style="list-style-type: none"> ○ Common Variable Immunodeficiency (CVID) ○ Severe Combined Immunodeficiency (SCID) ○ X-Linked Agammaglobulinemia ○ X-Linked with Hyperimmunoglobulin M (IgM) Immunodeficiency ○ Wiskott-Aldrich Syndrome ○ Members < 13 years of age with pediatric Human Immunodeficiency Virus (HIV) and CD-4 count > 200/mm³ • Neurological disorders including: <ul style="list-style-type: none"> ○ Guillain-Barré Syndrome ○ Relapsing-Remitting Multiple Sclerosis ○ Chronic Inflammatory Demyelinating Polyneuropathy ○ Myasthenia Gravis ○ Polymyositis and Dermatomyositis ○ Multifocal Motor Neuropathy • Chronic Lymphocytic Leukemia (CLL) • Autoimmune Neutropenia (AN) with absolute neutrophil count < 800 mm and history of recurrent bacterial infections • Autoimmune Hemolytic Anemia (AHA) • Liver or Intestinal Transplant • Immune Thrombocytopenia Purpura (ITP) including: <ul style="list-style-type: none"> ○ Requiring preoperative therapy for undergoing elective splenectomy with platelet count < 20,000 ○ Members with active bleeding & platelet count <30,000 ○ Pregnant members with platelet counts <10,000 in the third trimester ○ Pregnant members with platelet count 10,000 to 30,000 who are bleeding <table border="1" data-bbox="934 1110 1745 1365"> <thead> <tr> <th colspan="2">Table 1: FDA-Approved Maximum Immune Globulin Dosing</th> </tr> </thead> <tbody> <tr> <td>Gammaplex 5% - IV Infusion</td> <td>800mg/kg every 3 weeks</td> </tr> <tr> <td>Privigen IV Infusion</td> <td>800mg/kg every 3 weeks</td> </tr> <tr> <td>Gammagard liquid SQ or IV admin</td> <td>2.4 grams/kg/month</td> </tr> <tr> <td>Gammaked SQ or IV admin</td> <td>600 mg/kg every 3 weeks</td> </tr> <tr> <td>Gamunex-C SQ or IV admin</td> <td>600 mg/kg every 3 weeks</td> </tr> <tr> <td>Hizentra SQ admin</td> <td>12 grams every 2 weeks</td> </tr> <tr> <td>Cuvitru SQ admin</td> <td>12.6 grams every 2 weeks</td> </tr> </tbody> </table> <p>Grandfathering: Members currently receiving a preferred or non-preferred immunoglobulin product</p>	Table 1: FDA-Approved Maximum Immune Globulin Dosing		Gammaplex 5% - IV Infusion	800mg/kg every 3 weeks	Privigen IV Infusion	800mg/kg every 3 weeks	Gammagard liquid SQ or IV admin	2.4 grams/kg/month	Gammaked SQ or IV admin	600 mg/kg every 3 weeks	Gamunex-C SQ or IV admin	600 mg/kg every 3 weeks	Hizentra SQ admin	12 grams every 2 weeks	Cuvitru SQ admin	12.6 grams every 2 weeks
Table 1: FDA-Approved Maximum Immune Globulin Dosing																		
Gammaplex 5% - IV Infusion	800mg/kg every 3 weeks																	
Privigen IV Infusion	800mg/kg every 3 weeks																	
Gammagard liquid SQ or IV admin	2.4 grams/kg/month																	
Gammaked SQ or IV admin	600 mg/kg every 3 weeks																	
Gamunex-C SQ or IV admin	600 mg/kg every 3 weeks																	
Hizentra SQ admin	12 grams every 2 weeks																	
Cuvitru SQ admin	12.6 grams every 2 weeks																	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		may receive approval to continue therapy with that product at prescribed doses not exceeding maximum (Table 1).
--	--	---

Therapeutic Drug Class: NEWER GENERATION ANTIHISTAMINES -Effective 7/1/2019		
--	--	--

No PA Required	PA Required	
Cetirizine (generic OTC Zyrtec) 5mg and 10mg tab Cetirizine (RX) syrup Levocetirizine (OTC) tablet Loratadine (generic OTC Claritin) 10mg tab and syrup	ALAVERT (loratadine) ALLEGRA (fexofenadine) Cetirizine (OTC) chewable tablet, syrup CLARINEX (desloratadine) CLARITIN (loratadine) Desloratadine Fexofenadine Levocetirizine (RX) tablets, solution Loratadine chewable, ODT XYZAL (levocetirizine) ZYRTEC (cetirizine)	Non-preferred single agent antihistamine products may be approved for members who have failed treatment with two preferred products in the last 6 months. For members with respiratory allergies, an additional trial of an intranasal corticosteroid will be required in the last 6 months. Failure is defined as lack of efficacy with a 14 day trial, allergy, intolerable side effects, or significant drug-drug interaction.

Therapeutic Drug Class: ANTIHISTAMINE/DECONGESTANT COMBINATIONS - Effective 7/1/2019		
---	--	--

	PA Required	
	ALLEGRA-D (fexofenadine/PSE) Cetirizine-D (OTC) CLARINEX-D (desloratadine-D)	Non-preferred antihistamines and antihistamine/decongestant combinations will be approved for members who have failed treatment with two preferred products in the last 6 months. For members with respiratory allergies, an additional trial of an intranasal corticosteroid will be required in the last 6 months. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	CLARITIN-D (loratadine-D) (OTC) Fexofenadine/PSE (OTC) Loratadine-D (OTC) SEMPREX-D (acrivastine-D) ZYRTEC-D (cetirizine-D) (OTC)	
--	---	--

Therapeutic Drug Class: INTRANASAL RHINITIS AGENTS -Effective 4/1/2020

No PA Required	PA Required	
Azelastine 0.15%, 137 mcg Budesonide 32 mcg (OTC) Fluticasone 50 mcg (generic FLONASE) RX only Ipratropium Triamcinolone acetonide (generic Nasacort) (OTC)	ASTEPRO (azelastine) 0.15% BECONASE AQ (beclomethasone dipropionate) CHILD NASACORT (triamcinolone) DYMISTA (azelastine/ fluticasone propionate) FLONASE (fluticasone) 50 mcg (OTC) FLONASE SENSIMIST (fluticasone) 27.5 mcg (OTC) Flunisolide 0.025% Mometasone 50 mcg NASACORT AQ (triamcinolone) NASONEX (mometasone)	<p>Non-preferred products may be approved following trial and failure of treatment with three preferred products (failure is defined as lack of efficacy with a 2 week trial, allergy, intolerable side effects or significant drug-drug interactions).</p> <p>Non-preferred combination agents may be approved following trial of individual products with same active ingredients AND trial and failure of one additional preferred agent (failure is defined as lack of efficacy with 2 week trial, allergy, intolerable side effects or significant drug-drug interactions).</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	Olopatadine 665 mcg OMNARIS (ciclesonide) PATANASE (olopatadine) QNASL (beclomethasone dipropionate) XHANCE (fluticasone propionate) ZETONNA (ciclesonide)	
--	---	--

Therapeutic Drug Class: LEUKOTRIENE MODIFIERS -Effective 4/1/2020
--

No PA Required	PA Required	
Montelukast tab, chewable	ACCOLATE (zafirlukast) tablet SINGULAIR (montelukast) tablet, chewable tab, granules Montelukast granules Zafirlukast tablet ZYFLO (zileuton ER) tablet	Non-preferred products may be approved if meeting the following criteria: <ul style="list-style-type: none"> Member has trialed and failed treatment with one preferred product (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions) AND Member has a diagnosis of asthma. Montelukast granules may be approved if a member has tried and failed montelukast chewable tablets AND has difficulty swallowing.

Therapeutic Drug Class: MULTIPLE SCLEROSIS AGENTS -Effective 4/1/2020
--

Disease Modifying Therapies

No PA Required (unless indicated*)	PA Required	
AVONEX (interferon beta 1a) injection	COPAXONE (glatiramer) 40MG injection	*Second-line preferred agents (Gilenya, Tecfidera, Aubagio) may be approved if meeting the following: <ul style="list-style-type: none"> Member has documented diagnosis of multiple sclerosis made by neurologist in the last 3 years OR member has history of diagnosis made by a neurologist > 3 years ago but is naïve to all medications indicated for the treatment of relapsing forms of multiple sclerosis AND Documentation is provided by prescribing neurologist (or name of neurologist consulted may

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>BETASERON (interferon beta 1b) injection</p> <p>COPAXONE^{BNR} (glatiramer) 20MG injection</p> <p>*AUBAGIO (teriflunomide) tablet **2nd Line**</p> <p>*GILENYA^{BNR} (fingolimod) 0.5 mg tablet (30-ct bottle) **2nd Line**</p> <p>*TECFIDERA (dimethyl fumarate) tablet **2nd Line**</p>	<p>EXTAVIA (interferon beta 1b) vial</p> <p>GLATOPA (glatiramer) injection</p> <p>Glatiramer 20mg, 40mg injection</p> <p>GILENYA (fingolimod) 0.25 mg, 0.5 mg tablet (7-ct box)</p> <p>MAVENCLAD (cladribine) tablet</p> <p>MAYZENT (siponimod) tablet, pack</p> <p>PLEGRIDY (peg-interferon beta 1a)</p> <p>REBIF (interferon beta 1a) injection</p> <p>VUMERITY (diroximel) capsules</p>	<p>be indicated) supporting marked functional decline as demonstrated by <u>two</u> of the following: MRI, EDSS scale, or medical chart notes supporting increased burden of disease AND</p> <ul style="list-style-type: none"> • Prescriber attests to shared decision making with respect to risks versus benefits of medical treatment AND • Additional safety criteria for prescribed agent are met (Table 1). <p>For members NOT meeting above criteria, second-line preferred agents (Gilenya, Tecfidera, Aubagio) may be approved if meeting all of the following:</p> <ul style="list-style-type: none"> • Member has a diagnosis of a relapsing form of multiple sclerosis AND • Medication is being prescribed by a neurologist or in conjunction with consultation by a neurologist AND • Member has trial and failure with Copaxone OR a preferred interferon product (failure defined as intolerable side effects, drug-drug interaction, or lack of efficacy) AND • MRI results show presence of new spinal lesions, cerebellar lesions, brain stem lesions, or change in brain atrophy AND • On clinical exam, member has signs and symptoms consistent with functional limitations lasting one month or longer AND • Additional safety criteria for prescribed agent are met (Table 1). <p><u>Non-Preferred Products:</u> Mayzent (simponimod), Mavenclad (cladribine), and Vumerity (diroxemel fumerate) must meet specific criteria listed for those agents below. All other non-preferred products may be approved following trial and failure with three preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interactions).</p> <p>Copaxone (glatiramer) 40mg may be approved for members who have severe intolerable injection site reactions to <u>brand</u> Copaxone 20mg (such as pain requiring local anesthetic, oozing, lipoatrophy, swelling, or ulceration).</p> <p>Mayzent (simponimod) may be approved if meeting all of the following:</p> <ul style="list-style-type: none"> • Medication is being prescribed by a neurologist or in conjunction with consultation by a neurologist AND • Member has a diagnosis of a relapsing form of multiple sclerosis AND • Member <u>does not</u> have diagnosis of macular degeneration AND • Member has baseline Expanded Disability Status Scale (EDSS) score of 3.0-6.5 AND • Member has no evidence of relapse in the 3 months preceding initiation of therapy AND • Member has previous trial and failure of Gilenya (fingolimod) therapy (failure is defined as lack of efficacy with 3 month trial, allergy, intolerable side effects, or significant drug-drug interaction) AND • Additional safety criteria for prescribed agent are met (Table 1) AND
---	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<ul style="list-style-type: none"> • Initial authorization will be limited to 3 months. Continuation (12 month authorization) will require documentation of EDSS reduction of 1.0 point from baseline (or reduction of 0.5 points if baseline EDSS is 5.5-6.5). <p>Mavenclad (cladribine) may be approved if meeting all of the following:</p> <ul style="list-style-type: none"> • Medication is being prescribed by a neurologist or in conjunction with consultation by a neurologist AND • Member has a diagnosis of a relapsing form of multiple sclerosis AND • Member has history of ≥ 1 relapse in the 12 months preceding initiation of therapy AND • Member has previous trial and failure of three other therapies for relapsing forms of multiple sclerosis (failure is defined as lack of efficacy with 3 month trial, allergy, intolerable side effects, or significant drug-drug interactions) AND • Additional safety criteria for prescribed agent are met (Table 1). <p>Vumerity (diroximel fumarate) may be approved if meeting all of the following:</p> <ul style="list-style-type: none"> • Medication is being prescribed by a neurologist or in conjunction with consultation by a neurologist AND • Member has a diagnosis of a relapsing form of multiple sclerosis AND • Additional safety criteria for prescribed agent are met (Table 1) AND • Member has previous trial and failure of Tecfidera (dimethyl fumarate) therapy (failure is defined as lack of efficacy with 3 month trial, allergy, intolerable side effects [if GI adverse events, must meet additional criteria below], or significant drug-drug interactions) AND • If Vumerity (diroximel fumarate) is being prescribed due to GI adverse events with Tecfidera (dimethyl fumarate) therapy (and no other reason for failure of Tecfidera is given), then the following additional criteria must be met: <ul style="list-style-type: none"> ○ Member has trialed a temporary dose reduction of Tecfidera (with maintenance dose being resumed within 4 weeks) AND ○ Member has trialed taking Tecfidera with food AND ○ GI adverse events remain significant despite maximized use of gastrointestinal symptomatic therapies (such as calcium carbonate, bismuth subsalicylate, PPIs, H2 blockers, anti-bloating/anti-constipation agents, anti-diarrheal, and centrally acting anti-emetics) AND ○ Initial authorization will be limited to 3 months. Continuation (12 month authorization) will require documentation of clinically significant reduction in GI adverse events with Vumerity (diroximel fumarate) therapy.
Table 1: Safety Criteria for Select Agents		

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<table border="1"> <tr> <td data-bbox="928 280 1129 378"> Tecfidera (dimethyl fumarate) </td> <td data-bbox="1129 280 2024 378"> <ul style="list-style-type: none"> Member has no active infections AND Member has CBC with differential conducted within the 6 months prior to initiating therapy </td> </tr> <tr> <td data-bbox="928 378 1129 751"> Aubagio (teriflunomide) </td> <td data-bbox="1129 378 2024 751"> <ul style="list-style-type: none"> Member has no active infections AND For female members of child-bearing age, have negative pregnancy test at baseline and are using a highly effective form of contraceptive when appropriate (such as long-acting reversible contraception) AND Member has transaminase and bilirubin levels with ALT < 2 times the upper limit of normal within the 6 months prior to initiating therapy AND Member has CBC with differential conducted within the 6 months prior to initiating therapy AND Member has a documented baseline blood pressure AND Member has been evaluated for active or latent tuberculosis infection by documented test results (purified protein derivative test) or blood test. </td> </tr> <tr> <td data-bbox="928 751 1129 1295"> Gilenya (fingolimod) </td> <td data-bbox="1129 751 2024 1295"> <ul style="list-style-type: none"> Member has no active infections AND Member does not have history of myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or New York Heart Association Class III-IV heart failure within six months of initiating therapy AND Member does not have a history or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome (unless patient has a pacemaker) AND Member has a baseline QTc interval < 500 ms prior to starting therapy AND Member is not receiving treatment with a Class Ia or Class III anti-arrhythmic medication AND Member has had an ophthalmologic evaluation (ocular coherence test) prior to starting therapy with follow-up within 3-4 months after therapy is initiated AND Member has had baseline CBC with differential and liver function tests conducted. </td> </tr> <tr> <td data-bbox="928 1295 1129 1450"> Mayzent (simponimod) </td> <td data-bbox="1129 1295 2024 1450"> <ul style="list-style-type: none"> Member does not have one of the following contraindications: <ul style="list-style-type: none"> CYP2C9*3/*3 genotype OR Has experienced (in the last 6 months) myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, or Class III or IV heart failure OR </td> </tr> </table>	Tecfidera (dimethyl fumarate)	<ul style="list-style-type: none"> Member has no active infections AND Member has CBC with differential conducted within the 6 months prior to initiating therapy 	Aubagio (teriflunomide)	<ul style="list-style-type: none"> Member has no active infections AND For female members of child-bearing age, have negative pregnancy test at baseline and are using a highly effective form of contraceptive when appropriate (such as long-acting reversible contraception) AND Member has transaminase and bilirubin levels with ALT < 2 times the upper limit of normal within the 6 months prior to initiating therapy AND Member has CBC with differential conducted within the 6 months prior to initiating therapy AND Member has a documented baseline blood pressure AND Member has been evaluated for active or latent tuberculosis infection by documented test results (purified protein derivative test) or blood test. 	Gilenya (fingolimod)	<ul style="list-style-type: none"> Member has no active infections AND Member does not have history of myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or New York Heart Association Class III-IV heart failure within six months of initiating therapy AND Member does not have a history or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome (unless patient has a pacemaker) AND Member has a baseline QTc interval < 500 ms prior to starting therapy AND Member is not receiving treatment with a Class Ia or Class III anti-arrhythmic medication AND Member has had an ophthalmologic evaluation (ocular coherence test) prior to starting therapy with follow-up within 3-4 months after therapy is initiated AND Member has had baseline CBC with differential and liver function tests conducted. 	Mayzent (simponimod)	<ul style="list-style-type: none"> Member does not have one of the following contraindications: <ul style="list-style-type: none"> CYP2C9*3/*3 genotype OR Has experienced (in the last 6 months) myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, or Class III or IV heart failure OR
Tecfidera (dimethyl fumarate)	<ul style="list-style-type: none"> Member has no active infections AND Member has CBC with differential conducted within the 6 months prior to initiating therapy 									
Aubagio (teriflunomide)	<ul style="list-style-type: none"> Member has no active infections AND For female members of child-bearing age, have negative pregnancy test at baseline and are using a highly effective form of contraceptive when appropriate (such as long-acting reversible contraception) AND Member has transaminase and bilirubin levels with ALT < 2 times the upper limit of normal within the 6 months prior to initiating therapy AND Member has CBC with differential conducted within the 6 months prior to initiating therapy AND Member has a documented baseline blood pressure AND Member has been evaluated for active or latent tuberculosis infection by documented test results (purified protein derivative test) or blood test. 									
Gilenya (fingolimod)	<ul style="list-style-type: none"> Member has no active infections AND Member does not have history of myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or New York Heart Association Class III-IV heart failure within six months of initiating therapy AND Member does not have a history or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome (unless patient has a pacemaker) AND Member has a baseline QTc interval < 500 ms prior to starting therapy AND Member is not receiving treatment with a Class Ia or Class III anti-arrhythmic medication AND Member has had an ophthalmologic evaluation (ocular coherence test) prior to starting therapy with follow-up within 3-4 months after therapy is initiated AND Member has had baseline CBC with differential and liver function tests conducted. 									
Mayzent (simponimod)	<ul style="list-style-type: none"> Member does not have one of the following contraindications: <ul style="list-style-type: none"> CYP2C9*3/*3 genotype OR Has experienced (in the last 6 months) myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, or Class III or IV heart failure OR 									

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<ul style="list-style-type: none"> ○ Presence of Mobitz type II second-degree, third-degree AV block, or sick sinus syndrome (unless patient has a functioning pacemaker) AND • Member has baseline QTc interval < 500 ms prior to starting therapy AND • Member has no active infections AND • Member has not had hypersensitivity reaction to Gilenya (fingolimod) AND • Baseline CBC with differential and liver function tests are conducted prior to initiating therapy. <p>Maximum Dose: 60mg per 30 days</p>
		<p>Mavenclad (cladribine)</p> <ul style="list-style-type: none"> • Member has negative pregnancy test within 30 days of request for Mavenclad AND • Men and women of childbearing potential must have plan to use effective contraception during and 6-months after therapy AND • Member does not have current evidence of malignancy AND • Member has CBC with differential drawn prior to, during, and after treatments with Mavenclad due to risk of lymphopenia and hematologic toxicity AND • Lymphocytes must be within normal limits before initiating the first treatment course and must be ≥ 800 cells per microliter before initiating the second treatment course AND • Member is not currently taking immunosuppressive or myelosuppressive therapy AND • Member has no active infections AND • Member has liver function tests drawn prior to first and second treatment course due to potential for liver injury. <p>Maximum Dose: Not exceeding 3.5mg/kg during full treatment course</p>
		<p>Vumerity (diroximel fumarate)</p> <ul style="list-style-type: none"> • Member has not had hypersensitivity reaction or angioedema as a result of Tecfidera (dimethyl fumarate) therapy AND • Member has no active infections AND • A CBC with differential will be conducted within the six months prior to initiating therapy AND • Member has liver function tests drawn prior to treatment course due to

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

		potential for liver injury. Maximum Dose: 924mg per day	
Grandfathering: Members currently stabilized on a preferred second-line product or a non-preferred product may receive approval to continue therapy with that agent.			

Symptom Management Therapies

	PA Required	<p>Ampyra (dalfampridine) prior authorization for a 3 month supply may be approved if all of the following criteria are met:</p> <ul style="list-style-type: none"> • Member has a diagnosis of MS; Member is ambulatory and has established a baseline which is defined as ambulating between 8-45 seconds Timed 25-foot Walk (T25FW) assessment OR has established a baseline activities of daily living (ADL) AND • Member has no history of seizure disorder AND • Member has no history of moderate to severe renal dysfunction (CrCl > 50 ml/min) AND • Prescriber is a neurologist or is prescribed in conjunction with a neurologist AND • The prescribed dose does not exceed 10 mg twice daily. <p>Extended coverage of Ampyra (dalfampridine) for up to one year may be approved if documentation shows improvement in ambulation (measured by T25FW assessment) or improvement in ADLs after three months of therapy.</p>	
	<p>AMPYRA ER (dalfampridine)</p> <p>Dalfampridine ER</p>		

Therapeutic Drug Class: TARGETED IMMUNE MODULATORS - <i>Effective 1/1/2020</i>

Must meet eligibility criteria*	PA Required	Eligibility Criteria for preferred agents in the class:	
ENBREL (etanercept)	ACTEMRA (tocilizumab) syringe, Actpen	Humira or Enbrel may receive approval for use for FDA-labeled indications.	
HUMIRA (adalimumab)	ARCALYST (rilonacept) injection	Cosentyx may receive approval for FDA-labeled indications following trial and failure of Humira (failure is defined as lack of efficacy of a three-month trial, contraindication to therapy, allergy, intolerable side effects or significant drug-drug interaction).	
COSENTYX (secukinumab) syringe, pen-injector	CIMZIA (certolizumab) kit	Xeljanz IR may receive approval for ulcerative colitis following trial and failure of Humira (failure is defined as lack of efficacy of a three-month trial, contraindication to therapy, allergy, intolerable side effects or significant drug-drug interaction). Xeljanz IR may receive approval with no trial and failure required for rheumatoid arthritis and psoriatic arthritis. Quantity Limits: 2 tablets per day or 60 tablets for a 30 day supply.	
XELJANZ IR (tofacitinib) tablet	ILARIS (canakinumab) vial		
	KEVZARA (sarilumab) pen, syringe		
	KINERET (anakinra) syringe		

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	<p>OLUMIANT (baricitinib) tablet</p> <p>ORENCIA (abatacept) syringe, clickject</p> <p>OTEZLA (apremilast) tablet</p> <p>RINVOQ (upadacitinib) tablet</p> <p>SILIQ (brodalumab) syringe</p> <p>SIMPONI (golimumab) pen, syringe</p> <p>SKYRIZI (risankizumab-rzaa) syringe, kit</p> <p>STELARA (ustekinumab) syringe</p> <p>TALTZ (ixekizumab) auto-injector, syringe</p> <p>TREMFYA (guselkumab) injector, syringe</p> <p>XELJANZ XR (tofacitinib ER) tablet</p> <p>*for information on IV infused Targeted Immune Modulators please see Appendix P</p>	<p>Non-Preferred Agents may receive prior authorization approval for FDA-labeled indications following trial and failure ALL preferred agents (Enbrel, Humira, Cosentyx, and Xeljanz IR) that are FDA-labeled for use for the same prescribed indication (failure is defined as lack of efficacy of a three-month trial, contraindication to therapy, allergy, intolerable side effects or significant drug-drug interaction). Agents listed below must meet the following additional criteria for approval of that agent:</p> <p>Arcalyst (rilonacept): Prior authorization approval will be given for an initial 12 weeks and authorization approval for continuation will be provided based on clinical response.</p> <p>Kineret (anakinra): May receive approval for use for familial Mediterranean fever. Approval for all other indications is subject to meeting non-preferred criteria listed above.</p> <p>Rinvoq (upadacitinib) may receive approval if meeting non-preferred criteria listed above AND following trial and failure of Olumiant (baricitinib). Failure is defined as lack of efficacy of a three-month trial, contraindication to therapy, allergy, intolerable side effects or significant drug-drug interaction.</p> <p>Siliq (brodalumab), Skyrizi (risankizumab-rzaa), or Tremfya (guselkumab) may receive approval if meeting non-preferred criteria listed above AND following trial and failure of Otezla (apremilast). Failure is defined as lack of efficacy of a three-month trial, contraindication to therapy, allergy, intolerable side effects or significant drug-drug interaction.</p> <p>Stelara (ustekinumab): Loading dose administration prior to approval of Stelara for maintenance therapy using the above criteria should be avoided and will not result in an automatic approval of Stelara maintenance therapy. Prior authorization approval may be given for an initial 16 weeks and authorization approval for continuation will be provided based on clinical response. Stelara IV vial formulation may receive approval under the pharmacy benefit if meeting non-preferred criteria listed above AND if being administered in a long-term care facility or the member's home by a home health provider (initial 16 week authorization may be placed for both IV and subcutaneous formulations at time of Stelara IV vial approval).</p> <p>Taltz (ixekizumab): Prior authorization approval will be given for an initial 12 weeks and authorization approval for continuation will be provided based on clinical response.</p> <p>Xeljanz (tofacitinib) XR: Approval will require verification of the clinically relevant reason for use of the Xeljanz XR formulation versus the Xeljanz IR formulation in addition to meeting non-preferred criteria listed above.</p>
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<i>The Department would like to remind providers that many products have patient support programs that assist patients in drug administration, education, and emotional support for our member's diseases.</i>
--	--	--

Therapeutic Drug Class: TOPICAL IMMUNOMODULATORS – Effective 7/1/2019		
--	--	--

*Must meet criteria	PA Required	
ELIDEL (pimecrolimus) ^{BNR} Pimecrolimus cream - <i>authorized generic only - Oceanside Pharm</i>	Pimecrolimus cream - <i>All other manufacturers</i> PROTOPIC (tacrolimus) Tacrolimus (generic Protopic)	Manual prior authorization review for preferred and non-preferred agents will be required for members exceeding ≥ 6 weeks of continuous therapy. Preferred topical immunomodulator products may be approved following adequate trial and failure [‡] of a prescription topical corticosteroid (verified in claims history). Non-preferred topical immunomodulator products may be approved following adequate trial and failure [‡] of one prescription topical corticosteroid AND one preferred agent. [‡] Failure is defined as a lack of efficacy with one month trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions. For members under 18 years of age, must be prescribed by or in conjunction with a dermatologist or allergist/immunologist.

X. Miscellaneous		
-------------------------	--	--

Therapeutic Drug Class: EPINEPHRINE PRODUCTS -Effective 1/1/2020		
---	--	--

No PA Required	PA Required	
<i>Generic changes effective 01/15/20</i> Epinephrine 0.15mg/0.3ml, 0.3mg/0.3ml auto-injector (generic Epipen) - <i>Mylan only-</i>	EPIPEN 0.3mg/0.3ml (epinephrine) auto-injector EPIPEN JR 0.15mg/0.3ml, (epinephrine) auto-injector Epinephrine 0.15mg/0.15ml, 0.3mg/0.3ml auto-injector (generic AdrenaClick)	Non-preferred products will be approved if the member has failed treatment with one of the preferred products. Failure is defined as allergy to ingredients in product or intolerable side effects. Quantity limit: 4 auto injectors per year unless used / damaged / lost

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	<p>Epinephrine 0.15mg/0.15ml, 0.3mg/0.3ml auto-injector (generic Epipen) -<i>Teva only</i>-</p> <p>SYMJEPI 0.15mg/0.3ml, 0.3mg/0.3ml (epinephrine) syringe</p>	
--	--	--

Therapeutic Drug Class: NEWER HEREDITARY ANGIOEDEMA PRODUCTS -Effective 10/1/2019

PA Required for all agents in this class		Medications Indicated for Routine Prophylaxis:
<p><i>Prophylaxis:</i></p> <p>HAEGARDA (C1 esterase inhibitor) 2,000 unit and 3,000 unit vial</p> <p><i>Treatment:</i></p> <p>BERINERT (C1 esterase inhibitor) 500 Unit kit</p> <p>FIRAZYR^{BNR} (icatibant acetate) 30mg/3 mL syringe</p>	<p><i>Prophylaxis:</i></p> <p>CINRYZE (C1 esterase inhibitor) 500 unit kit</p> <p>TAKHZYRO (lanadelumab) 300 mg/mL vial</p> <p><i>Treatment:</i></p> <p>Icatibant 30 mg/3 mL syringe</p> <p>RUCONEST (C1 esterase inhibitor, recomb) 2,100 unit vial</p>	<p>Members are restricted to coverage of one medication for <u>routine prophylaxis</u> at one time. Prior authorization approval will be for one year.</p> <p>Haegarda may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none"> ○ Member has a diagnosis of HAE confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, CI-INH level) AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member meets at least one of the following: <ul style="list-style-type: none"> ▪ Haegarda® is being used for short-term prophylaxis to undergo a surgical procedure or major dental work OR ▪ Haegarda® is being used for long-term prophylaxis and member meets one of the following: <ul style="list-style-type: none"> ○ History of ≥1 attacks per month resulting in documented ED admission or hospitalization OR ○ History of laryngeal attacks OR ○ History of ≥2 attacks per month involving the face, throat, or abdomen AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND ○ Member has received hepatitis A and hepatitis B vaccination AND ○ Provider attests to performing annual testing or screening (as appropriate) for HBV, HCV, and HIV <p>Max Dose: 60 IU/kg Minimum Age: 10 years</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<p>Cinryze and Takhzyro may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none"> ○ Member has history of trial and failure of Haegarda®. Failure is defined as lack of efficacy allergy, intolerable side effects, or a significant drug-drug interaction AND ○ Member has a diagnosis of HAE confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, CI-INH level) AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member meets at least one of the following: <ul style="list-style-type: none"> ▪ Cinryze® is being used for <u>short-term prophylaxis</u> to undergo a surgical procedure or major dental work OR ▪ Cinryze® is being used for <u>long-term prophylaxis</u> and member meets one of the following: <ul style="list-style-type: none"> ○ History of ≥ 1 attacks per month resulting in documented ED admission or hospitalization OR ○ History of laryngeal attacks OR ○ History of ≥ 2 attacks per month involving the face, throat, or abdomen AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND ○ Member has received hepatitis A and hepatitis B vaccination AND ○ Provider attests to performing annual testing or screening (as appropriate) for HBV, HCV, and HIV. <p>Minimum age: Cinryze: 6 years Takhzyro: 12 years</p> <p>Max dose: Cinryze: 100 Units/kg Takhzyro: 300mg every 2 weeks</p> <p>Medications Indicated for Treatment of Acute Attacks:</p> <p>Members are restricted to coverage of one medication for <u>treatment of acute attacks</u> at one time. Prior authorization approval will be for one year.</p> <p>Firazyr may be approved for members meeting the following criteria:</p>
--	--	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

		<ul style="list-style-type: none"> ○ Member has a diagnosis of HAE confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, CI-INH level) AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications <p>Minimum age: 18 years Maximum dose: 30mg</p> <p>Berinert may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none"> ○ Member has a diagnosis of HAE confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, CI-INH level) AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND ○ Member has received hepatitis A and hepatitis B vaccination AND ○ Provider attests to performing annual testing or screening (as appropriate) for HBV, HCV, and HIV <p>Minimum age: 6 years Max dose: 20 IU/kg</p> <p>Ruconest may be approved for members meeting the following criteria:</p> <ul style="list-style-type: none"> ○ Member has a history of trial and failure of Firazyr® OR Berinert®. Failure is defined as lack of efficacy, allergy, intolerable side effects, or a significant drug-drug interaction AND ○ Member has a diagnosis of HAE confirmed by laboratory tests obtained on two separate instances at least one month apart (C4 level, CI-INH level) AND ○ Member has a documented history of at least one symptom of a moderate to severe HAE attack (moderate to severe abdominal pain, facial swelling, airway swelling) in the absence of hives or a medication known to cause angioedema AND ○ Member is not taking medications that may exacerbate HAE including ACE inhibitors and estrogen-containing medications AND ○ Member has received hepatitis A and hepatitis B vaccination AND
--	--	---

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

		<ul style="list-style-type: none"> ○ Provider attests to performing annual testing or screening (as appropriate) for HBV, HCV, and HIV. <p>Minimum age: 13 years Max dose: 4200 Units/dose</p>
--	--	---

Therapeutic Drug Class: PHOSPHATE BINDERS -Effective 7/1/2019

No PA Required	PA Required	
<p><i>Generic changes effective 10/15/19</i></p> <p>Calcium acetate capsule</p> <p>FOSRENOL^{BNR} (lanthanum carbonate) chewable tablet</p> <p>PHOSLYRA (calcium acetate)</p> <p>RENAGEL^{BNR} (Sevelamer hcl)</p> <p>Sevelamer carbonate tablet (6-17 years old)*</p> <p>Sevelamer HCL <i>authorized generic -WINTHROP US only</i></p> <p>-</p>	<p>AURYXIA (ferric citrate)</p> <p>Calcium acetate tablet (generic Calphron)</p> <p>FOSRENOL (lanthanum carbonate) powder pack</p> <p>Lanthanum carbonate chewable tablet, powder pack</p> <p>RENVELA (sevelamer carbonate)</p> <p>Sevelamer carbonate powder pack</p> <p>Sevelamer hcl tablet <i>-all other manufacturers</i></p> <p>VELPHORO (sucoferric oxide)</p>	<p>*Sevelamer carbonate tablet may be approved as a preferred agent for children and adolescents 6-17 years of age. For adults ≥ 18 years of age, sevelamer carbonate tablet may be approved if member meets criteria for non-preferred products listed below.</p> <p>Prior authorization for non-preferred products in this class may be approved if member meets all of the following criteria:</p> <ul style="list-style-type: none"> ● Member has diagnosis of end stage renal disease AND ● Member has elevated serum phosphorus [> 4.5 mg/dL or > 1.46 mmol/L] AND ● Provider attests to member avoidance of high phosphate containing foods from diet AND ● Member has trialed and failed‡ two preferred agents. One trial must be from the same pharmacologic class as the non-preferred agent being requested, if applicable (for example; member is requesting Phoslo[®] must have trial with Phoslyra[®] or generic calcium acetate). <p>Auryxia[®] (ferric citrate) may be approved if the member meets all of the following criteria:</p> <ul style="list-style-type: none"> ● Member is diagnosed with end-stage renal disease, receiving dialysis, and has elevated serum phosphate (> 4.5 mg/dL or > 1.46 mmol/L). AND ● Provider attests to counseling member regarding avoiding high phosphate containing foods from diet AND ● Member has trialed and failed‡ three preferred agents with different mechanisms of action prescribed for hyperphosphatemia in end stage renal disease <p>OR</p> <ul style="list-style-type: none"> ● Member is diagnosed with chronic kidney disease with iron deficiency anemia and is not receiving dialysis AND ● Member has tried and failed‡ at least two different iron supplement product formulations (OTC or RX) <p>‡Failure is defined as lack of efficacy with 6 week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p><i>Note: Medications administered in a dialysis unit or clinic are billed through the Health First Colorado medical benefit.</i></p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

Therapeutic Drug Class: **PRENATAL VITAMINS / MINERALS** -Effective 10/1/2019

PA Required (must meet eligibility criteria)	PA Required	
CITRANATAL 90 DHA combo pack CITRANATAL ASSURE combo pack CITRANATAL B-CALM CITRANATAL DHA pack CITRANATAL HARMONY capsule CITRANATAL RX tablet COMPLETE NATAL DHA CONCEPT DHA capsule CONCEPT OB capsule M-NATAL PLUS NESTABS tablets PNV OB+DHA COMBO PACK PNV PNV-FERROUS FUMARATE-DOCU-FA tablet PRENAISSANCE PLUS capsule PRENATAL LOW IRON tablet PRENATAL VITAMIN PLUS LOW IRON PREPLUS tablet	All other rebateable prescription products are non-preferred	*Preferred and non-preferred prenatal vitamin products are a benefit for members from 11-60 years of age who are pregnant, lactating, or trying to get pregnant. Prior authorization for non-preferred agents will be approved if member fails 7-day trial with four preferred agents. (Failure is defined as: allergy, intolerable side effects, or significant drug-drug interaction)

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

TRINATAL RX 1 TRUST NATAL DHA VIRT-ADVANCE TABLET VIRT-VITE GT TABLET VOL-PLUS tablet		
---	--	--

XI. Ophthalmic

Therapeutic Drug Class: **OPHTHALMIC, ALLERGY** -Effective 4/1/2020

No PA Required	PA Required	
ALREX (loteprednol) 2%	ALAWAY (ketotifen) 0.025%	Non-preferred products may be approved following trial and failure of therapy with two preferred products (failure is defined as lack of efficacy, allergy, intolerable side effects or significant drug-drug interactions).
Cromolyn 4%	ALOCRIL (nedocromil) 2%	
Ketotifen (generic Zaditor) 0.025% (OTC)	ALOMIDE (lodoxamide) 0.1%	
LASTACAFT (alcaftadine) 0.25%	Azelastine 0.05%	
Olopatadine 0.1%, 0.2%	BEPREVE (bepotastine) 1.5%	
PAZEO (olopatadine) 0.7%	Epinastine 0.05%	
	PATADAY (olopatadine) 0.2%	
	PATANOL (olopatadine) 0.1%	
	ZADITOR (ketotifen) 0.025% (OTC)	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

Therapeutic Drug Class: OPHTHALMIC, IMMUNOMODULATORS -Effective 10/1/2019

No PA Required	PA Required	<p>Non-preferred products may be approved for members meeting all of the following criteria:</p> <ul style="list-style-type: none"> • Member is 18 years and older AND • Member has a diagnosis of chronic dry eye AND • Member has failed a three month trial of one preferred product (Failure is defined as a lack of efficacy, allergy, intolerable side effects, contraindication to, or significant drug-drug interactions) AND • Prescriber is an ophthalmologist, optometrist or rheumatologist <p><u>Maximum Quantity:</u> 60 single use containers for 30 days 5.5 mL/20 days for Restasis Multi-Dose</p>
<p>RESTASIS (cyclosporine 0.05%)</p>	<p>Cequa (cyclosporine 0.09%) solution</p> <p>RESTASIS MULTIDOSE (cyclosporine 0.05%)</p> <p>XIIDRA (lifitegrast)</p>	

Therapeutic Drug Class: OPHTHALMIC, ANTI-INFLAMMATORIES -Effective 4/1/2020

NSAIDs		<p>Non-preferred products may be approved with trial and failure of three preferred agents (failure is defined as lack of efficacy with 2 week trial, allergy, contraindication, intolerable side effects, or significant drug-drug interaction).</p> <p>Lotemax SM (loteprednol etoabonate) may be approved if meeting all of the following:</p> <ul style="list-style-type: none"> • Member is ≥ 18 years of age AND • Lotemax SM (loteprednol etoabonate) is being used for the treatment of post-operative inflammation and pain following ocular surgery AND • Member has trialed and failed therapy with two preferred loteprednol formulations (failure is defined as lack of efficacy with 2 week trial, allergy, contraindication, intolerable side effects, or significant drug-drug interaction) AND • Member has trialed and failed therapy with two preferred agents that do not contain loteprednol (failure is defined as lack of efficacy with 2 week trial, allergy, contraindication, intolerable side effects, or significant drug-drug interaction) AND • Member <u>does not</u> have any of the following conditions: <ul style="list-style-type: none"> ○ Viral diseases of the cornea and conjunctiva including epithelial herpes simplex keratitis (dendritic keratitis), vaccinia, and varicella OR ○ Mycobacterial infection of the eye and fungal diseases of ocular structures
No PA Required	PA Required	
<p>ACUVAIL (ketorolac)</p> <p>Bromfenac 0.09%</p> <p>Diclofenac 0.1%</p> <p>Flurbiprofen 0.03%</p> <p>Ketorolac 0.5%, Ketorolac LS 0.4%</p>	<p>ACULAR (ketorolac) 0.5%, LS 0.4%</p> <p>BROMSITE (bromfenac) 0.075%</p> <p>ILEVRO (nepafenac) 0.03%</p> <p>NEVANAC (nepafenac) 0.1%</p> <p>PROLENSA (bromfenac) 0.07%</p>	
Corticosteroids		
No PA Required	PA Required	
<p>Flarex (fluorometholone) 0.1%</p> <p>Fluorometholone 0.1% drops</p>	<p>Dexamethasone 0.1%</p> <p>DUREZOL (difluprednate) 0.05%</p>	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

FML Forte (fluorometholone) 0.25% drops LOTEMAX (loteprednol) 0.5% drops, 0.5% ointment MAXIDEX (dexamethasone) 0.1% PRED MILD (prednisolone) 0.12% Prednisolone acetate 1%	FML LIQUIFILM (fluorometholone) 0.1% drop FML S.O.P (fluorometholone) 0.1% ointment INVELTYS (loteprednol) 1% LOTEMAX (loteprednol) 0.5% gel LOTEMAX SM (loteprednol) 0.38% gel Loteprednol 0.5% drops OMNIPRED (prednisolone) 1% PRED FORTE (prednisolone) 1% Prednisolone sodium phosphate 1%	
---	---	--

Therapeutic Drug Class: OPTHALMIC, GLAUCOMA -Effective 4/1/2020

Beta-blockers		
No PA Required	PA Required	
Levobunolol Timolol (generic Timoptic)	BETAGAN (levobunolol) Betaxolol BETOPIC-S (betaxolol) Carteolol ISTALOL (timolol) Timolol (generic Istalol) drops	<p>Non-preferred products may be approved following trial and failure of therapy with three preferred products, including one trial with a preferred product having the same general mechanism (such as prostaglandin analogue, alpha2-adrenergic agonist, beta-blocking agent, or carbonic anhydrase inhibitor). Failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions.</p> <p>Non-preferred combination products may be approved following trial and failure of therapy with one preferred combination product AND trial and failure of individual products with the same active ingredients as the combination product being requested (if available) to establish tolerance. Failure is defined as lack of efficacy with 4 week trial, allergy, intolerable side effects or significant drug-drug interactions.</p> <p>Preservative free products may be approved with provider documentation of adverse effect to preservative-containing product.</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	<p>Timolol GFS</p> <p>TIMOPTIC, TIMOPTIC OCUDOSE (timolol)</p> <p>TIMOPTIC-XE (timolol GFS)</p>	
Carbonic anhydrase inhibitors		
No PA Required	PA Required	
<p>AZOPT (brinzolamide)</p> <p>Dorzolamide</p>	<p>TRUSOPT (dorzolamide)</p>	
Prostaglandin analogue		
No PA Required	PA Required	
<p>Latanoprost</p> <p>LUMIGAN^{BNR} (bimatoprost)</p> <p>TRAVATAN Z^{BNR} (travoprost)</p>	<p>Bimatoprost</p> <p>Latanoprost PF</p> <p>VYZULTA (latanoprostene)</p> <p>XALATAN (latanoprost)</p> <p>XELPROS (latanoprost)</p> <p>ZIOPTAN (tafluprost PF)</p>	
Alpha-2 adrenergic agonists		
No PA Required	PA Required	
<p>ALPHAGAN P 0.1% (brimonidine)</p>	<p>Apraclonidine</p>	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

ALPHAGAN P ^{BNR} 0.15% (brimonidine) Brimonidine 0.2%	Brimonidine 0.15% IOPIDINE (apraclonidine)	
Other ophthalmic, glaucoma and combinations		
No PA Required	PA Required	
COMBIGAN (brimonidine/timolol) Dorzolamide/Timolol Dorzolamide/Timolol PF	COSOPT PF (dorzolamide/timolol) Echothiopate iodide PHOSPHOLINE IODIDE (echothiophate) Pilocarpine RHOPRESSA (netarsudil) ROCKLATAN (netarsudil) SIMBRINZA (brinzolamide/brimonidine)	

XII. Renal/Genitourinary

Therapeutic Drug Class: **OVERACTIVE BLADDER AGENTS** -*Effective 10/1/19*

No PA Required	PA Required	
GELNIQUE (oxybutynin) gel, pump Oxybutynin IR, ER tablets, syrup	Darifenacin ER tablet DETROL (tolterodine) DETROL LA (tolterodine ER)	Non-preferred products will be approved for members who have failed treatment with two preferred products. Failure is defined as: lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction. Members with hepatic failure can receive approval for trospium (Sanctura) or trospium extended-release (Sanctura XR) products without a trial on a Preferred product.

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

Oxybutynin ER tablets TOVIAZ (fesoterodine ER)	DITROPAN (brand) DITROPAN XL (brand) ENABLEX (darifenacin) Flavoxate MYRBETRIQ (mirabegron) OXYTROL (oxybutynin patch) SANCTURA (trospium) SANCTURA XL (trospium ER) Solifenacin tablet Tolterodine Trospium ER capsule, tablet VESICARE (solifenacin)	
---	---	--

Therapeutic Drug Class: ANTI-HYPERURICEMICS -Effective 1/1/2020

No PA Required	PA Required	
Allopurinol tablet	Colchicine tablet	Prior authorization for non-preferred xanthine oxidase inhibitor products (allopurinol or febuxostat formulations) may be approved after trial and failure of preferred allopurinol. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.
Probenecid tablet	COLCRYS (colchicine) tablet	
Colchicine capsule	Febuxostat tablet	If member has tested positive for the HLA-B*58:01 allele, it is not recommended that they trial allopurinol. A positive result on this genetic test will count as a failure of allopurinol.
Probenecid/Colchicine tablet	GLOPERBA (colchicine) oral solution	Prior authorization for all other (non-xanthine oxidase inhibitors) non-preferred agents may be approved after trial and failure of two preferred products. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.
	MITIGARE (colchicine) capsule	

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

	ULORIC (febuxostat) tablet ZYLOPRIM (allopurinol) tablet	Prior authorization for colchicine tablets may be approved for members requiring treatment of gout flares. Colchicine tablet quantity limits: <ul style="list-style-type: none"> • Chronic hyperuricemia/gout prophylaxis: 60 tablets per 30 days • Familial Mediterranean Fever: 120 tablets per 30 days
--	---	--

Therapeutic Drug Class: BENIGN PROSTATIC HYPERPLASIA (BPH) -Effective 7/1/19

No PA Required	PA Required	
Alfuzosin	AVODART (dutasteride)	Prior authorization for non-preferred products in this class may be approved if member meets all of the following criteria: <ul style="list-style-type: none"> • Member has tried and failed‡ three preferred agents AND • For combinations agents, member has tried and failed‡ each of the individual agents within the combination agent and one other preferred agent. ‡Failure is defined as lack of efficacy with 8 week trial, allergy, intolerable side effects, contraindication to, or significant drug-drug interaction. *Cialis will be approved for members with a documented diagnosis of BPH who have failed a trial of finasteride (at least 3 months in duration) AND either a trial of a nonselective alpha blocker (therapeutic dose for at least two months) OR a trial of tamsulosin (therapeutic dose for at least one month). Documentation of BPH diagnosis will require BOTH of the following: <ul style="list-style-type: none"> • AUA Prostate Symptom Score \geq 8 AND • Results of a digital rectal exam. Cialis will not be approved for any patient continuing alpha-blocker therapy as this combination is contraindicated in this population. Doses exceeding 5mg per day of Cialis will not be approved.
Doxazosin	CARDURA (doxazosin)	
Dutasteride	CARDURA XL (doxazosin ER)	
Finasteride	*CIALIS (tadalafil) 2.5 mg, 5 mg only <i>Brand and generic</i>	
Tamsulosin	FLOMAX (tamsulosin)	
Terazosin	JALYN (dutasteride/tamsulosin) <i>Brand and generic</i>	
	PROSCAR (finasteride)	
	RAPAFLO (silodosin) <i>Brand and generic</i>	

XIII. RESPIRATORY

Therapeutic Drug Class: RESPIRATORY INHALANTS -Effective 7/1/2019

Inhaled Anticholinergics

No PA Required	PA Required	
<u>Solutions</u>	<u>Solutions</u> ATROVENT (ipratropium) solution	Non-preferred single agent anticholinergic agents will be approved for members with a diagnosis of COPD including chronic bronchitis and/or emphysema who have trialed and failed‡ treatment with two preferred agents, one of which must be Spiriva Handihaler.

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	---

<p>Ipratropium (generic Atrovent) solution</p> <p><u>Short-Acting Inhalers</u> ATROVENT HFA (ipratropium)</p> <p><u>Long-Acting Inhalers</u> SPIRIVA Handihaler (tiotropium)</p>	<p>LONHALA Magnair (glycopyrrolate) solution</p> <p>YUPELRI (revefenacin) solution</p> <p><u>Short-Acting Inhalers</u></p> <p><u>Long-Acting Inhalers</u></p> <p>INCRUSE ELLIPTA (umeclidinium)</p> <p>SEEBRI Neohaler (glycopyrrolate)</p> <p>SPIRIVA RESPIMAT (tiotropium)</p> <p>TUDORZA Pressair (aclidinium)</p>	<p>Spiriva Respimat® will be approved for members with a diagnosis of asthma who have trialed and failed‡ treatment with three preferred inhaled corticosteroids, at least two of the trials must be preferred combination inhaled corticosteroid products. Members with a diagnosis of COPD must meet non-preferred criteria for single agent inhaled anticholinergics listed above for approval of Spiriva Respimat®.</p> <p>Lonhala Magnair® will receive prior authorization approval for members ≥ 18 years of age with a diagnosis of COPD including chronic bronchitis and emphysema who have trialed and failed‡ treatment with two preferred anticholinergic agents.</p> <p>‡Failure is defined as lack of efficacy with 6 week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p>
--	---	--

Inhaled Anticholinergic Combinations

No PA Required	PA Required	
<p><u>Solutions</u> Albuterol/ipratropium solution</p> <p><u>Short-Acting Inhalers</u> COMBIVENT RESPIMAT (albuterol/ipratropium)</p>	<p><u>Solutions</u></p> <p><u>Short-Acting Inhalers</u></p> <p><u>Long-Acting Inhalers</u> ANORO ELLIPTA (umeclidinium/vilanterol)</p> <p>BEVESPI AEROSPHERE (glycopyrrolate/formoterol fumarate)</p> <p>STIOLTO Respimat (tiotropium/olodaterol)</p> <p>UTIBRON Neohaler (glycopyrrolate/indacaterol)</p>	<p>Non-preferred combination anticholinergic agents will be approved for members with a diagnosis of COPD including chronic bronchitis and/or emphysema who have trialed and failed‡ treatment with two preferred respiratory agents, one of which must be Spiriva Handihaler®. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>‡Failure is defined as lack of efficacy with 6 week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

Inhaled Beta2 Agonists (short acting)
--

<p style="text-align: center;">No PA Required</p> <p style="text-align: center;"><i>Brand changes effective 3/24/2020</i></p> <p><u>Solutions</u> Albuterol (generic) solution</p> <p><u>Inhalers</u></p> <p>PROAIR (albuterol) HFA ^{BNR}</p> <p>PROVENTIL (albuterol) HFA inhaler ^{BNR}</p> <p>VENTOLIN (albuterol) HFA inhaler ^{BNR}</p>	<p style="text-align: center;">PA Required</p> <p><u>Solutions</u></p> <p>PROVENTIL (albuterol) solution</p> <p>XOPENEX (levalbuterol) solution</p> <p><u>Inhalers</u></p> <p>Albuterol HFA</p> <p>Levalbuterol HFA</p> <p>PROAIR Respiclick (albuterol)</p> <p>XOPENEX (levalbuterol) Inhaler</p>	<p>Non-preferred, short acting beta2 agonists will be approved for members who have failed treatment with one preferred agent. Failure is defined as lack of efficacy, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>Proair HFA, Proventil HFA, Ventolin HFA: Quantity limits: 2 inhalers / 30 days</p>
--	---	--

Inhaled Beta2 Agonists (long acting)

<p>*Must meet eligibility criteria</p> <p><u>Solutions</u></p> <p><u>Inhalers</u></p> <p>*SEREVENT DISKUS (salmeterol) inhaler</p>	<p style="text-align: center;">PA Required</p> <p><u>Solutions</u></p> <p>BROVANA (arformoterol) solution</p> <p>PERFOROMIST (formoterol) solution</p> <p><u>Inhalers</u></p> <p>ARCAPTA Neohaler (indacaterol)</p> <p>FORADIL (formoterol)</p> <p>STRIVERDI Respimat (olodaterol)</p>	<p>SEREVENT ® will be approved for members with moderate to very severe COPD.</p> <p>Non-preferred agents will be approved for members with moderate to severe COPD, AND members must have failed a trial of Serevent®. (Failure is defined as lack of efficacy with a 6 week trial, allergy, intolerable side effects, or significant drug-drug interaction.</p> <p>**For treatment of members with diagnosis of asthma needing add-on therapy, please refer to preferred agents in combination Long-Acting Beta Agonist/Inhaled Corticosteroid. SEREVENT will not be approved for treatment of asthma in members needing add-on therapy due to safety risks associated with monotherapy.</p>
---	---	--

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
-------------------------	-----------------------------	--

Inhaled Corticosteroids

No PA Required	PA Required	
<p><u>Solutions</u></p> <p>Budesonide nebulules 0.25mg 0.5mg, 1mg</p> <p><u>Inhalers</u></p> <p>ASMANEX Twisthaler (mometasone)</p> <p>FLOVENT Diskus(fluticasone)</p> <p>FLOVENT HFA (fluticasone)</p>	<p><u>Solutions</u></p> <p>PULMICORT (budesonide) nebulules 0.25mg 0.5mg, 1mg</p> <p><u>Inhalers</u></p> <p>AEROSPAN HFA (flunisolide) inhaler</p> <p>ALVESCO (ciclesonide) inhaler</p> <p>ARMONAIR Respiclick (fluticasone)</p> <p>ARNUITY Ellipta (fluticasone furoate)</p> <p>ASMANEX HFA (mometasone furoate) inhaler</p> <p>PULMICORT Flexhaler(budesonide)</p> <p>QVAR Redihaler (beclomethasone)</p>	<p>Pulmicort (Budesonide) nebulizer solution will only be approved for a maximal dose of 2mg/day.</p> <p>Pulmicort Flexhaler will only be approved for female members with asthma who have a new diagnosis of pregnancy.</p> <p>Non-preferred inhaled corticosteroids will be approved in members with asthma who have failed an adequate trial of two preferred agents. An adequate trial is defined as at least 6 weeks. (Failure is defined as: lack of efficacy with a 6 week trial, allergy, contraindication to, intolerable side effects, or significant drug-drug interactions.)</p>

Inhaled Corticosteroid Combinations
--

No PA Required	PA Required	
<p><i>Brand/generic changes effective 11/01/19</i></p> <p>ADVAIR Diskus^{BNR} (fluticasone/salmeterol)</p> <p>ADVAIR HFA (fluticasone/salmeterol)</p>	<p>AIRDUO Respiclick (fluticasone/salmeterol)</p> <p>BREO Ellipta (vilanterol/fluticasone furoate)</p> <p>Fluticasone/salmeterol (generic Airduo)</p>	<p>Non-preferred inhaled corticosteroid combinations will be approved for members meeting both of the following criteria:</p> <ul style="list-style-type: none"> • Member has a qualifying diagnosis of asthma or severe COPD; AND • Member has failed two preferred agents (Failure is defined as lack of efficacy with a 6 week trial, allergy, intolerable side effects, significant drug-drug interactions, or dexterity/coordination limitations (per provider notes) that significantly impact appropriate use of a specific dosage form.) <p>Trelegy Ellipta® prior authorization will be approved if the member has trialed/failed three preferred inhaled corticosteroid combination products AND Spiriva Handihaler®. Failure is defined as lack of efficacy with a 6 week trial, allergy, intolerable side effects, significant drug-drug</p>

Preferred Agents	Non-preferred Agents	Prior Authorization Criteria (All Non-Preferred Products will be approved for one year unless otherwise stated.)
------------------	----------------------	--

DULERA (mometasone/formoterol) SYMBICORT (budesonide/formoterol) inhaler	Fluticasone/salmeterol diskus (generic Advair) TRELEGY Ellipta (Fluticasone Furoate/Umeclidinium/Vilanterol) WIXELA Inhub (fluticasone/salmeterol)	interactions, or dexterity/coordination limitations (per provider notes) that significantly impact appropriate use of a specific dosage form.
---	--	---